

A
C O U R S E
O F
E X P E R I M E N T A L P H I L O S O P H Y.

BY
J. T. DESAGULIERS, LL.D. F.R.S.
Chaplain to his Grace the DUKE of *CHANDOS*.

V O L. I.
Adorn'd with Thirty-two COPPER-PLATES.



L O N D O N :

Printed for JOHN SENEX, in *Fleetstreet*; W. INNYS and RICHARD
MANBY, in *St. Paul's Church-Yard*; and JOHN OSBORN and
THOMAS LONGMAN in *Pater-noster Row*. MDCCXXXIV.

TO HIS
ROYAL HIGHNESS
FREDERICK,
Prince of *WALES*, &c.

S I R,

TO contemplate the Works of GOD,
to discover Causes from their Effects,
and make Art and Nature subservi-
ent to the Necessities of Life, by a Skill in
joining proper Causes to produce the most use-
ful

D E D I C A T I O N.

ful Effects, is the Business of a Science, the Grounds and Principles of which, I have the Honour to lay at Your ROYAL HIGHNESS's Feet.

I have presum'd to do this, Philosophy in all Ages having been thought worthy the Consideration of Princes ; for the greatest and best, who have been handed down to us by Historians, under the most amiable and glorious Characters, have distinguish'd themselves by their Attainments in it, and their Encouragement of all Endeavours to advance and improve it.

Amongst these I may justly reckon his late MAJESTY, who was, and his present MAJESTY, who is, the Patron of a Society, erected for the Advancement of Natural Knowledge, of which one of their Royal Predecessors was the Founder.

Your

D E D I C A T I O N.

Your known Candor (GREAT SIR) will, I hope, excuse my plain Manner of treating this Subject: Tho' I take the Liberty to assure Your ROYAL HIGHNESS, that I have spared no Pains, that the Work might not be wholly unworthy of Your Patronage.

It presumes the more upon Your Countenance and Protection, as it is an Account of those Experiments, which I had the Honour to make some Years ago at *Hampton-Court*, before his late MAJESTY, by his particular Command; and as their present MAJESTIES, who likewise honour'd those Experiments with their Presence, have also given me Leave to set their Names at the Head of my Subscribers.

Were I equal to the Description of those excellent Qualities, which make You the present Delight, and future Hope of *Britain*; yet I should be guilty of the highest Indiscretion to attempt it in an Address to Your self;
since

D E D I C A T I O N.

since none are more offended with Praise than those who most deserve it. I shall therefore no longer detain Your ROYAL HIGHNESS, than whilst I beg Leave to subscribe my self, with the greatest Respect,


S I R,

Your most humble,

most obedient,

and most dutiful Servant,

J. T. DESAGULIERS.



THE
NAMES of such Persons,
As have encouraged this
WORK
BY THEIR
SUBSCRIPTIONS.

His late Majesty King GEORGE the First,
HIS PRESENT MAJESTY,
HER PRESENT MAJESTY,

Allan,

SUBSCRIBERS NAMES.

A

A Llan, Mr. Lionel, *Merchant at Rotterdam*
 Allen, Ralph Esq; *Post-master of Bath*
 Anderfon, James D. D.
 Apreece, Robert Esq;
 Armstrong, Col. John *Surveyor of his Majesty's Ordnance*
 Arnold, Robert, *Teacher of Mathematicks*

B

B Aldwin, Samuel Esq;
 Barker, Mrs. Mary, *of Broadwell in Gloucestershire*
 Baron, Rev. Charles A. M.
 Beal, John M. D.
 Beaumont, Sir George Bart.
 Beeldemaker, Gerandus
 Bentinck, Rt. Hon. William
 Bernard, John, *of Asbury, Esq;*
 Bethel, Hugh Esq;
 Big, Rev. A. M. F. W. Coll.
 Billers, Sir William, *Lord Mayor of London*
 Blackburn, William Esq;
 Bladen, Col. Martin
 Blunt, Sir John Kt.
 Bond, Dennis Esq;
 Booth, Barton Esq;
 Booyaunt, Paulus
 Bornemannus, Philip Julius, *of Holland*
 Boyd, Mr. Robert
 Brace, J. Thurlow Esq;
 Bradock, Mrs. Frances
 Branden, Peter Adrianus van den
 Bridges, Brook Esq;
 Bridges, John Esq;
 Bridges, J--- Esq;
 Brown, Mr. Thomas *Merchant*
 Brownlow, Rt. Hon. Lady E.
 Brownlow, William Esq;

Buchleugh, His Grace Francis, *Duke of*
 Bucler, Mr. John
 Buckenbrough, Rt. Hon. Countess of
 Burlington, Rt. Hon. Earl of
 Butler, Rt. Hon. Col. James
 Bute, Rt. Hon. James Earl of

C

C Apper, — Esq;
 Carr, William Esq;
 Carpenter, Rt. Hon. Lord
 Cartwright, Thomas Esq;
 Caswell, Sir George
 Chandos, His Grace James, *Duke of*
 Chamber, William Esq;
 Cheney, Rt. Hon. Lord
 Christophers, Theodore, Jan.
 Church, Thomas Esq;
 Churchill, Mr. Robert
 Churchill, Mr. Robert, *Apothecary*
 Churchill, Brig. Charles
 Churchman, Mr. Walter
 Cibber, Colley Esq;
 Clarke, Rev. Samuel D. D.
 Clare, Mr. Martin, *School-master*
 Clements, Mr. John
 Cleaveland, His Grace Duke of
 Cobb, Rev. Dr. Ward. Winch. Coll.
 Cobham, Lord Viscount
 Conduit, John Esq;
 Conway, Rt. Hon. Lady
 Cooke, Sir George Kt.
 Cope, Charles Esq;
 Corfini, Marq. Neri
 Costar, Thomas Esq;
 Costar, Mr. John
 Colbe, Mr. J. L. *Mathematical Instrument-maker*
 Coladon, — Esq;
 Collet, Dr. John
 Craufurd, Rt. Hon. Earl of
 Crop, Mr. Abraham
 Cumming, Sir Alexander Bart.

Daniel,

SUBSCRIBERS NAMES.

D

Daniel, *Capt. Hugh*
 Dappe, *Mr. John*
 Darlington, *Countess of*
 Davies, *Mr. Meredith*
 Davis, *Mr. — Shipwright at Amsterdam*
 Deering, *Daniel Esq;*
 Dela Faye, *Charles Esq;*
 Denune, *Mr. George*
 Dickinson, *Mr. Ezekiel*
 Diston, *Jo. Esq;*
 Dobson, *Rev. J. D. D. Ward. of New Coll.*
 Dobyons, *Col. John*
 Donald, *Mr. John*
 Douglass, *George M. D.*
 Dumford, *Mr. Richard*

E

Earle, *Giles Esq;*
 Earle, *Mr. Timothy*
 Edens, *Mr. Abraham Merchant*
 Eden, *Mr. John*
 Edgecumbe, *Rt. Hon. Richard Esq;*
 Edwards, *Thomas Esq;*
 Edwards, *— Esq;*
 Edwin, *Charles Esq;*
 Eglington, *Rt. Hon. Countess of*
 Elliot, *Mr. John*
 Elliot, *Sir Gilbert Bart.*
 Elton, *Mr. Abraham*
 Everhard, *Mr. John*

F

Fairfax, *Brian Esq;*
 Fisher, *Mr. Samuel*
 Fizeaux, *Mr. Jean*
 Fletcher, *Mr. Joshua*
 Folkes, *Martin Esq;*
 Foreman, *Mr. George*
 Frederick, *John Esq;*

Furly, *Mr. Benjohan Merchant*
 Furnese, *George Esq;*

G

Garnier, *Mons. — M. D.*
 Gascoigne, *Mr. J----*
 Gazola, *His Excellency Count of*
 Geelvink, *N. Secret.*
 Gibbs, *Rev. Mr. John*
 Girardot, *John Esq;*
 Golding, *Mr. John*
 Goodchild, *Mr. John*
 Gordon, *Capt. George*
 Gore, *John Esq;*
 Graham, *Richard Esq;*
 Greenway, *Rev. William A. M.*
 Greenwood, *Charles Esq;*
 Grove, *John Esq;*
 Grundy, *Mr. John Surveyor*

H

Hales, *Rev. Stephen B. D.*
 Halifax, *Rt. Hon. George Earl of*
 Hall, *Stephen M. D.*
 Halsey, *Richard Esq;*
 Hamilton, *Lady Archibald*
 Hamilton, *Lady Margaret*
 Hamilton, *Mr. Archibald Merchant*
 Hanger, *Mrs. Jane*
 Harvey, *John Esq;*
 Hardwich, *Dr. Peter*
 Harford, *Mr. Trueman*
 Hawkey, *— Esq;*
 Heath, *Ralph Esq;*
 Hays, *Charles Esq;*
 Haynes, *Mr. Thomas*
 Helot, *Mr. John, Clock-maker*
 Hepburn, *Mr. John*
 Herring, *John Esq;*
 Hewet, *Sir Thomas Kt.*
 Higgins, *Dr. Abdias*
 Hillsborough, *Rt. Hon. ---- Trevor Viscount*

b

Hooper,

SUBSCRIBERS NAMES.

Hooper, *Rev. Dr. Francis*
 Howard, *Hugh Esq;*
 Hunt, *Rev. Dr. late Master of B.*
Coll. Oxon 30 Books
 Hunt, *Rev. A. M. V. P. of H. H.*
Oxon
 Hutchinson, *Rev. Thomas, A. M.*
 Hutchinson, *John Esq;*

I

JAMES, *Mr. Thomas*
 Jaumar, *Rev. John, Chaplain to*
his Grace the Duke of Grafton
 Iberville, *Marquis de*
 Jefferies, *John Esq;*
 Jenkins, *Tobias Esq;*
 Jones, *William Esq;*

K

KENT, *Charles Esq;*
 King, *Mr. James*
 Kox, *Engil*
 Kingliffe, *Mr. William*
 Krighout, *Julius*
 Kynaston, *Mr. Thomas*

L

L Ambert, *Sir Edward*
 Law, ----- *Esq;*
 Leader, *Charles, A. M. F. N. Coll.*
 Leigh, *John Esq;*
 Lippe, *Rt. Hon. Countess de la*
 Lippe, *Rt. Hon. Count de la, Sen.*
 Lippe, *Rt. Hon. Count de la, Jun.*
 Loftus, *Rev. Mr. Bartholomew*
 Longdale, *Lord Viscount of*
 Lord, *Robert Esq;*
 Loup, *Mr. William*
 Lufnew, — *M. D.*
 Lynslager, *Capt. Henry*

M

M Acclesfield, *late Rt. Hon. Tho.*
Earl of, Lord Chancellor
 Mackworth, *Sir Thomas Bart.*
 Mackenzie, *Sir George*
 Mathew, *George Esq;*
 Mazeres, *Peter Esq;*
 Meads, *Peter Esq;*
 Milbourne, *George Esq;*
 Middlemore, *John Esq;*
 Millner — *Esq;*
 Mitchel, *Mr. Henry*
 Molewater, *Bastiaen Jun.*
 Montague, *His Grace John Duke of*
 Morgan, *Mr. John, of Bristol*

N

N Ewton, *late Sir Isaac Rt.*
 Niewaart, *Cornelius M. D.*
 Nicholas, *John Esq;*
 Nicholson, *Henry M. D.*
 Norfolk, *His Grace the Duke of*
 Nourfe, *Charles Esq;*
 Nyevelt, *Harm. Van Zuyle Secr.*

O

O Glethorpe, *James Esq;*
 Osborne, *Sir John Bart.*

P

P Aisley, *Rt. Hon. James Lord*
 Parsons, *Col. John*
 Pawlet, *Rt. Hon. William Lord*
 Pembroke, *Rt. Hon. Hen. Earl of,*
two Books
 Phillips, *Mr. Robert Bricklayer*
 Piddocke, *Mr. Thomas*
 Plumpton, *Richard Esq;*
 Popple,

SUBSCRIBERS NAMES.

Popple, Henry *Esq*;
 Price, William *Esq*;
 Price, Mr. John
 Pringle, John *M. D.*
 Prude, Mr. Henry
 Purcell, Col. Martin
 Pye, Rev. Sir Richard

Stanton, Mr. Ward
 Stephens, Mr. ---
 Steel, Mr. Eugene
 Steighertahl, — *M. D.*
 Stone, Mr. Arthur
 Strickland, Sir William
 Suffolk, Rt. Hon. Edward Earl of
 Suffex, Rt. Hon. Talbot Earl of
 Sutton, Sir Robert Kt.

R

R Edding, — *Col.*
 Rippley, Thomas *Esq*;
 Riva, John *Esq*;
 Robin, Mr. Pieter
 Rogers, Mr. Francis
 Rooke, George *Esq*;
 Rouffignac, Guy *M. D. Lecturer of*
Anatomy at Surgeons-Hall
 Rowel, Mr. John, *Glass-stainer after*
the antient manner at Wickham
 Rulhout, Sir John Bart.
 Rutty, William *M. D.*

T

T Aylor, Brook *L. L. D.*
 Tayleur, John *Esq*;
 Thomond, Rt. Hon. Earl of
 Tobin, Sir James
 Trevor, John *Esq*;
 Trimble, Mr. Francis Merchant
 Truby, Mr. Richard, *Jun.*
 Tuffinell, Mr. Samuel
 Turbot, Richard *Esq*;
 Tyrconnel, Rt. Hon. John Earl of

S

S Almon, Dr. William Henry
 Sambrook, John *Esq*;
 Samuda — *M. D.*
 Sands, Mr. William
 Senex, Mr. John, *Bookfeller*
 Shepherd, Mr. John
 Sherigley, Abraham *Esq*;
 Skreen, Richard *Esq*;
 Skipworth, Sir Fulwar
 Schelches, Mr. Andrew
 Sloan, Sir Hans Bart.
 Smith, Dr. Richard, 2 Books
 Snellen, Jan Van
 Southcoat, Philip *Esq*;
 Stampel, Mr. Peter
 Stanhope, Charles *Esq*;

V

V Alentinois, Duc de
 Vanbrugh, Sir John
 Vaughan, Gwyne *Esq*;
 Vincent, Mr. William
 Vynor, Robert *Esq*;

W

W Ager, the Hon. Sir Charles
 Walpole, Hon. Horatio *Esq*;
 Wagg, Mr. Thomas
 Wall, William *Esq*;
 Wall, Mr. Tobias
 Weddel, Capt. Charles
 Wharton, His Grace Philip Duke of

SUBSCRIBERS NAMES.

Wharton, George *M. D.*

Wharton, Edward *Jun.*

Wilkinson, Andrew *Esq;*

Woodcock, Thomas *Esq;*

Woodefon, *Mr. ----*

Woodhouse, *Mr. William*

Wyndham, Thomas *Esq;*

Wyndham, *Sir William*


Y

Y Vonet, John Paul *Esq;*

N. B. *The Subscribers are desir'd to forgive any Mistake in their Titles, or the spelling of their Names.*



PREFACE.



P R E F A C E.

ALL the Knowledge we have of Nature depends upon Facts ; for without Observations and Experiments, our natural Philosophy would only be a Science of Terms and an unintelligible Jargon. But then we must call in Geometry and Arithmetick to our Assistance, unless we are willing to content our selves with natural History and conjectural Philosophy. For, as many Causes concur in the Production of compound Effects, we are liable to mistake the predominant Cause, unless we can measure the Quantity of the Effects produc'd, compare them with, and distinguish them from each other, to find out the adequate Cause of each single Effect, and what must be the Result of their joint Action. When Mons. Des Cartes's philosophical Romance, by the Elegance of its Style and the plausible Accounts of natural Phænomena, had overthrown the Aristotelian Physicks, the World receiv'd but little Advantage by the Change : For instead of a few Pedants, who, most of them, being conscious of their Ignorance, conceal'd it with hard Words and pompous Terms ; a new Set of Philosophers started up, whose lazy Disposition easily fell in with a Philosophy, that required no Mathematicks to understand it ; and who taking a few Principles for granted, without examining their Reality or Consistence with each other, fancied they could

P R E F A C E.

could solve all *Appearances* mechanically by *Matter and Motion*; and, in their smattering *Way*, pretended to demonstrate such things, as perhaps *Cartesius* himself never believ'd; his *Philosophy* (if he had been in earnest) being unable to stand the *Test* of the *Geometry* which he was *Master* of.

It is to *Sir Isaac Newton's Application of Geometry to Philosophy*, that we owe the routing of this *Army of Goths and Vandals* in the philosophical *World*; which he has enrich'd with more and greater *Discoveries*, than all the *Philosophers* that went before him: And has laid such *Foundations* for future *Acquisitions*; that even after his *Death*, his *Works* still promote natural *Knowledge*. Before *Sir Isaac*, we had but wild *Guesses* at the *Cause* of the *Motion* of the *Comets* and *Planets* round the *Sun*; but now he has clearly deduc'd them from the universal *Laws* of *Attraction* (the *Existence* of which he has prov'd beyond *Contradiction*) and has shewn, that the seeming *Irregularities* of the *Moon*, which *Astronomers* were unable to express in *Numbers*, are but the just *Consequences* of the *Actions* of the *Sun* and *Earth* upon it, according to their different *Positions*. His *Principles* clear up all *Difficulties* of the various *Phænomena* of the *Tides*; and the true *Figure* of the *Earth* is now plainly shewn to be a flatted *Spheroid* higher at the *Æquator* than the *Poles*, notwithstanding many *Assertions* and *Conjectures* to the contrary. Our incomparable *Philosopher* has discovered and demonstrated to us the true *Nature* of *Light* and *Colours*, of which the most sagacious and inquisitive *Naturalists* were entirely ignorant; for while they sought for the *Origine* of *Colours* in the *Mixture* of *Light* and *Shadow*, *Sir Isaac Newton* found that they were congenial with the *Rays* of the *Sun*, and contain'd in *Light* it self; the *Surface* of colour'd
Bodies,

P R E F A C E.

Bodies, serving only to separate from one another those Rays, that make different Colours ; by absorbing some, and reflecting others to our Eyes, so as to produce those different Sensations, on which the pleasing Variety of colour'd Objects depends. His Opticks, besides the Properties of Light, contain a vast Fund of Philosophy ; which (tho' he has modestly delivered under the Name of Queries, as if they were only Conjectures) daily Experiments and Observations confirm ; a notable Instance of which may be seen in the Rev. Mr. Stephen Hale's excellent Book of Vegetable Staticks, which, by putting several of Sir Isaac's Queries out of all Doubt, shew how well they were founded. I pass over Sir Isaac Newton's noble Inventions in pure Mathematicks, justly admired at home and abroad ; because, tho' they have been of great use in the Discovery of the Causes of natural Phænomena, they are foreign to my present Subject, which is Physicks ; whose Knowledge I am in this Course endeavouring to convey by Experiments ; not only where Things have been discover'd that way, but even where they have been deduc'd by a long Train of mathematical Consequences ; having contriv'd Experiments, which Step by Step bring us to the same Conclusions.

The Thoughts of being oblig'd to understand Mathematicks have frighted a great many from the Newtonian Philosophy : I have heard several Cartesians say, that if the Knowledge of Geometry was necessary for their Conviction, they had rather continue in their own Way of Philosophy than be at so much Trouble ; as if a Man could deserve the Name of a Philosopher, merely because he reasons justly from Principles, when those Principles are either apparently false, precarious, or assum'd at pleasure to serve the present Purpose. It is not he, that can shew from an Hypothesis how the Celestial Motions
might

P R E F A C E.

might be performed ; but he, that demonstrates their real Causes, who gives a proper Account of the System of the World : And it is the same of other Phænomena ; for unless we can demonstrate what we explain, it is better to own our Ignorance, than to endeavour to pass our Conjectures upon the World for Solutions. If ever we come to know the Causes of the various Operations of Magnetism ; it will sooner be owing to a Comparison of the Experiments and Observations of Norman, Pound, Lord Paisley, Graham, Muschenbrock, Savery, Marcel and others (who acknowledge themselves ignorant of the Causes of those surprizing Effects) than to twenty Hypotheses of Men, whose warm Imaginations supply them with what may support their Solutions, while daily Observations and common Laws of Motion can easily confute them.

But to return to the Newtonian Philosophy ; tho' its Truth is supported by Mathematicks, yet its Physical Discoveries may be communicated without. The great Mr. Locke was the first who became a Newtonian Philosopher without the Help of Geometry ; for having asked Mr. Huygens, whether all the mathematical Propositions in Sir Isaac's Principia were true, and being told he might depend upon their Certainty ; he took them for granted, and carefully examined the Reasonings and Corollaries drawn from them, became Master of all the Physicks, and was fully convinc'd of the great Discoveries contain'd in that Book : Thus also he read the Opticks with Pleasure, acquainting himself with every thing in them that was not merely mathematical.* But since Machines have been contriv'd to explain and prove experimentally what Sir Isaac Newton has demonstrated mathe-

* This I was told several times by Sir Isaac Newton himself.

atically,

P R E F A C E.

matically, and several of his own Experiments are shewn in publick Courses; a great many Persons get a considerable Knowledge of Natural Philosophy by Way of Amusement; and some are so well pleas'd with what they learn that Way, as to be induc'd to study Mathematicks, by which they at last become eminent Philosophers. Dr. John Keill, was the first who publickly taught Natural Philosophy by Experiments in a mathematical Manner: for he laid down very simple Propositions, which he prov'd by Experiments, and from those he deduc'd others more compound, which he still confirm'd by Experiments; till he had instructed his Auditors in the Laws of Motion, the Principles of Hydrostaticks and Opticks, and some of the chief Propositions of Sir Isaac Newton concerning Light and Colours. He began these Courses in Oxford, about the Year 1704 or 1705, and that Way introduc'd the Love of the Newtonian Philosophy. There were indeed, about the same time, Experiments shewn at London by the late Mr. Hauksbee, which were electrical, hydrostatical, and pneumatical: But as they were only shewn and explain'd as so many curious Phænomena, and not made Use of as Mediums to prove a Series of philosophical Propositions in a mathematical Order, they laid no such Foundation for true Philosophy as Dr. Keill's Experiments; tho' perhaps perform'd more dexterously and with a finer Apparatus: They were Courses of Experiments, and his a Course of Experimental Philosophy.

When Dr. Keill left the University, I began to teach Experimental Philosophy, after the same Method that he had done, adding the Mechanicks (strictly so call'd, that is, the Explanation of mechanical Organs, and the Reason of their Effects) and several Optical Propositions in my Courses of Experimental Philosophy; which ever since that time I

c

have

P R E F A C E.

have endeavour'd to improve, by the Addition of new Propositions and Experiments, and by altering and changing my Machines, as I found Things might be made more intelligible to such of my Auditors as were not acquainted with Mathematicks, or more satisfactory to such as were; especially in what regards the Causes of the Motions of the heavenly Bodies, and the Phænomena of our System. About the Year 1713 I came to settle at London, where I have with great Pleasure seen the Newtonian Philosophy so generally received among Persons of all Ranks and Professions, and even the Ladies, by the Help of Experiments; that tho' several ingenious Men have since that Time with great Success taught (and do still teach) Experimental Philosophy in my (or rather Dr. Keill's) manner, I have had as many Courses as I could possibly attend; the present Course, which I am now engag'd in, being the 121st since I began at Hart-Hall in Oxford, in the Year 1710. The Satisfaction we enjoy by being any way instrumental to the Improvement of others, is so great, that I can't help boasting---that of eleven or twelve Persons, who perform Experimental Courses at this Time in England, and other Parts of the World, I have had the Honour of having Eight of them for my Scholars; whose further Discoveries become an Advantage to my self; for what would raise Envy in any other Profession, but that of a Philosopher, is receiv'd as a new Acquisition by all Lovers of Natural Knowledge, the Profit being shar'd in common, while the Discoverer has only the Honour of the Invention.

For this Reason, I never scruple making Use of Machines and Instruments contriv'd by others; nor was I ever shy of communicating, or even lending, my own, to those, who wanted to imitate them; it is enough to acknowledge the Author of any new Contrivance, which I generally do.

As

P R E F A C E.

As the greatest Part of my Auditors, at whose Desire I have printed this Course, are but little vers'd in Mathematical Sciences; the Lectures are free from difficult geometrical Demonstrations and algebraical Calculations; and the same thing is often prov'd by several Experiments; that where one does not immediately strike with a clear Conviction, another may. I only require Attention and common Sense, with a very little Arithmetick, in my Readers, to qualify them for understanding these Lectures; provided they begin the first Lecture, and go on regularly, that they may advance from the easiest Truths to those more complex ones, which are deduc'd from them; for otherwise many things may seem difficult to a Person, who should open the Book at random; especially great Part of the last Lecture of this Volume, which yet may be clearly understood by all, who have made themselves Masters of what goes before. Perhaps the Mathematicians may think me tedious and verbose in my Lectures; but such of them as have been us'd to teach, know very well, that one cannot be too plain and explicit with those that are not born with a Genius for Mathematicks (whatever good Understanding they are otherwise endow'd with) nay sometimes one must make Use of such Ways of demonstrating as are not mathematically true, to prepare them for what is a little more abstract; as I have been often forc'd to do to a large Audience, where close Attention is not very common. But I hope the rigid Philosophers will forgive me, when they find the same Things geometrically demonstrated in the Annotations; in the perusal of which, the Mathematicians perhaps will not think their Time wholly lost. However I don't mean to exclude common Readers from the Notes; for those that have carefully read and understood the Lectures, will thereby be qualified for comprehending what is in the Annotations.

P R E F A C E.

I should now close this Preface, if the Largeness of the Errata did not seem to want an Apology; but after all the Care an Author can take, mathematical Books have been of late so incorrectly printed, as to give the Reader a great deal of Trouble, and especially young Beginners, who are apt to attribute to their own Incapacity the Difficulty they find in Things, that are made unintelligible by the Printer's Fault; therefore I rather chose to offend the Eye, than puzzle the Understanding; and when I was resolv'd to rectify every Error, that might mislead the Reader, I thought it would be as well to add a Page or two, and take in every false Pointing: So that if all the Faults be first corrected with a Pen, according to the Direction in the Errata, I hope nothing will stop or puzzle an attentive Reader.

Besides, to prevent the Publick from being impos'd upon, I must not omit mentioning, that about sixteen Years ago, some Persons published a Book of Experimental Philosophy in my Name, without my Knowledge, which they endeavour'd to pass upon the World for my Lectures. It is a thin Quarto, and was at that Time call'd, A System of Natural Philosophy: And as those, who were capable of such a Thing, may, very probably, if they have any of the Books left, endeavour to sell them by giving them the same Title as my Book, I thought proper to give this Caution. But while I am complaining of others, I might be thought to ascribe to my self what is not my own, if I did not acknowledge, that most of what I have said of the Bow and Spring in the last Lecture of this Volume, as also Part of what I have said of the Fly and the Battering Ram, was copied from some Papers lent me by William Jones, Esq;

The

The READER is desir'd before he begins the Book, with his Pen to correct the following ERRATA.

Instead of

read

Page 3	Lin. 3	of the Preface, <i>abfording</i>	<i>abforbing</i>
2	13	Geometricians	Geometricians.
ib.	14	Structure, and	Structure, Powers, and
4	2	divided	divifible
6	14 and 15	9. Before the Air &c.	9. If the Receiver had been perfectly full before the Air was drawn out of it (tho' then their
10	17 and 18	its	their
13	3	write the Number 22 at the Beginning of the Paragraph	
20	3	write 38. before IF	
21	Antepenult	known,	known ;
ib.		Caufes ;	Caufes,
22	16	dele B o	
28	24	622 $\frac{1}{2}$	62, 25
36	ult.	write in the Margin	Fig. 18.
38	7	Point Q	Point O
39	27	At the End <i>dele</i> the	
53	16	overlook'd, and	overlook'd : And
56	25	Gravity of C :	Gravity of D :
60	11	Center of Motion K	Center of Gravity K
70	32	A is the	A is nearer to the
	33	then	than
71	6	probable	portable
	14	48	96
	19	24 Feet — 48 Times	48 Feet — 24 Times
	24	48 times lefs	24 times lefs
	32	4 times in	6 times in
	35	as troublefome to carry as	more troublefome to carry than
72	16	B G, B D	B G, G D
73	6	Diagonal	Hypotenuse
74	27	A B x C D :	A B + C D :
	30	A B x F E :	A B + F E
78	17	is M C	is equal to M C
83	25	fix Bodies	five Bodies
97	27	<i>dele</i> and	
105	34	raifing	rifing
	antepenult	from the	from that Part of the
107	10 and 11	<i>dele</i> whilst the Body W. rifes the Height B Z,	
108	23	C of the Arm D O,	C and the Arm D c.
111	31	the Plate,	the Plates,
112	28	THE Screw	61. THE Screw
	35	IN the	62. IN the
114	5	So is the Weight :	So is the Weight :
115	ult.	Powers, the	Powers (except the Wedge of which the Screw is made) the
118	3	Weights of their	Weights and their
125	21	Weight	Weights.
132	43	write in the Margin	Plate 13. Fig. 12.
136	12	Piece as Steel	Piece of Steel
	14	Eye c e	Eye E e
	16	supports the E	supports the Eye E e
	24	C may	C c may

Piece

E R R A T A.

	<i>Instead of</i>	<i>read</i>
P. 136 L. 25	Piece C.	Piece C c.
139 12	A B with	A B, with
150 29	Pipe E F	Pipe E H
30	Wire K <i>k</i> as far as <i>h</i>	Wire H <i>k</i> as far as D
168 8	and the Weight	and the Weight, and their Velocities
19	of which D	of which C
14	the Weight <i>w</i> ; the Weight W	the Weight <i>w</i> ; the Weight <i>w</i>
169 16	P and P.	P and P).
170 7	<i>b m</i> and <i>a t</i>	<i>b m</i> and <i>a n</i>
11	draw E C ——— to C D	draw E c — to C b
11	B <i>m b</i>	B <i>q b</i>
32	dele all the Paragraph The first &c. consisting of 8 Lines.	
171 4	and <i>m n r</i>	and <i>o C r</i>
41	The Weight	The Weight (taking in the Weight of the Wheel);
172 16	goes thro'	goes perpendicularly thro'
penult.	Weight ::	Weight of the Wheel and what it carries ::
173 29	Rub at <i>g</i> :	Rub at <i>g</i> , supposing the Wheels of the same Weight
176 28	of the Weight	of the Velocity
177 27	farther from	nearer to
28	nearer.	farther.
179 10	this Force	this Force of the Weight
181 16	D D is	D <i>d</i> is
183 21	of one th	of one 12th
24	is one 1 th	is one 12th
185 23	Base B C;	Length of the Plane A B
30	of <i>a c</i> ,	of A c
31	one 30th	a little more than one 30th
33	100 lb.	a little more than 100 lb.
186 2	Pl. 14.	Pl. 17.
188 31	dele more than	
ult.	of a Leather	of Leather
189 3	Direction a D	Direction A D
8	Wheel F	Wheel D E
30	124,16 lb.	124,61 lb.
237 36	in all 21,41 Ounces	in all 21,55 Ounces
241 31	to draw 140 lb.	to draw 240 lb.
243 18	up Hill	up Hill, when his Weight is made use of; and therefore he goes the slower.
246 28	for that the Fly	for the Fly
250 28	the Gudgeon	the other Gudgeon
252 38	then the	then let the
41	it will add	and it will add
253 12	Prim.	Prism
28	19. Fig. 1.	20. Fig. 10.
256 17	I P	F P
ult.	having brought	having brought his Legs
258 ult.	Position H L	Position h L
259 16	the Board to	that the Board should
260 antepenult.	Rolator	Rotator
261 31	about the Stone	about the Hammer
266 19	top Ropes	top Rope
275 25	Fig. 1.	Fig. 1 and 2.

revolts

E R R A T A,

	<i>Instead of</i>	<i>read</i>
P. 275 L. 31	revolts	revolves
278 19	Section	Profilis
282 6	after <i>Additions</i>	But the first Hint was from Mr. <i>Geo. Graham</i> .
9	D C; at D big enough	D C at D, big enough
37	W (<i>Fig. 6.</i>)	w (<i>Fig. 6.</i>)
283 16	of third	of the third
ult. after <i>Ec.</i>		N. B. <i>The 9th Figure is only part of the 8th Figure drawn larger.</i>
286 34	the Line A B	the Line A b,
35	the Line A b, becaufe	the Line A B, becaufe
287 15	Point D but	Point D over against him, but
291 2	<i>Fig. 2.</i>	<i>Fig. 1.</i>
	in the Margin—Pl. 24. F. 2.	Pl. 24. F. 1.
293 ult.	suppos'd it to make; the	suppos'd it to make the
296 26	<i>b a c, k,</i>	<i>b a, c k</i>
26	<i>i x,</i> which	<i>i x,</i> severally parallel to those Tangents, which
297 26	uniformly diminish'd	continually diminish'd
28	uniformly accelerated.	continually accelerated.
301 27	rectilineal	rectilineal
303 24	rise to α	rise to 3
304 penult.	join by a String to the	joyn, by a String, the
305 4	Distances of	Distances of the Centers of
6	that is, the	that is the Center of the
7	the four-Ounce Ball	the Center of the four-Ounce Ball
310 4	The 2d Figure	the 3d Figure
311 21	The Weights	9. The Weights
318 26	after—in a Second	N. B. <i>We don't consider any Elasticity that the Body might have.</i>
35	Diagonal E f	Diagonal E f
320 30	Force shou'd.	Force (supposing it now diminish'd in the Pro- portion of L 2 to L k) should
penult	by half.	by near half
ult.	but the half	but a little more than half
321 19	its Fall is	its Fall (considering Gravity only as a Blow) is
329 9	31 Miles. But	31 Miles; and partly, becaufe the Increase of the centrifugal Force destroys more of the the Action of Gravity near the Æquator than towards the Poles. But
penult.	from A to E.	from A to H
330 2 and 3	advance towards the Center of the Earth no	advance upon the Plane no
21	after ——— than A C.	N. B. A more Geometrical Solution of this may be found in the 37th Theorem of Dr. <i>John</i> <i>Keill</i> in his <i>Introductio ad veram Physicam</i> .
345 8	great than its Gravity,	greater still,
346 30	the former Arca	the former Area
349 16	one another, when.	one another, even when
20	8 Days longer	8 Days shorter
355 31	going off,	going off;
35	to A,	to A,
37	down to β ,	down to B
38	quit at B	quit it at β
357 21	Action, or equal	Action, are equal
21	A G and B.	A G and B F
361 22	the Part B $\alpha \beta$	the Part $\beta \alpha \delta$
362 26	up from l.	up to l,
368 12	and the contrary	and the same

ERRATA.

	<i>Instead of</i>	<i>read</i>
P. 368 L. 13	Southern Regions	Southern Regions in the opposite Months.
26	Water as high	Water nearly as high
28	have as high	have nearly as high
371 18	from the Center <i>b</i> round <i>o</i> to	round the Center <i>b</i> from <i>o</i> to
374 29	together, every other Stroke	together every other Stroke,
376 7	Pivion	Pinion
377 22	Barnouillii	Bernoulli
	<i>penult.</i> and <i>e D</i> : fo	and <i>e D</i> ; but more especially, because the Tangents are every where parallel to the corresponding Chords :
	Second Page 374. lin. 20. about 9 Hours	about 10 Hours.
379 20	the Body's	the Spring's
381 28	from C to A	from G to A,
30	lateral Motion,	lateral Motion of the Parallelogram in which he is plac'd
382 11	a Man carried	a Man at B carried
384 17	all the Distance	at the Distance
389 31	<i>Peribelia</i> be DE ;	<i>Peribelia</i> be D, E ;
391 5	Pl. 28. Fig. 11.	Pl. 24. F. 11.
405 24	15 $\frac{1}{8}$ Paris Feet.	15 $\frac{1}{8}$ Paris Feet.
418 2	Ball = <i>b</i> = lb,	Ball = <i>b</i> = 24 lb,
	9 as 15630.	as 15630 = 6400 X 2,4.
	23 25630	15630
436 12	28 Degrees, and of ♀ 48	about 28 Degrees, and of ♀ about 48
445 35	<i>dele</i> at Full	
	<i>Title of the Index</i> ALPHABETICAL	ALPHABETICAL

SUBSCRIBERS NAMES omitted.

Betenfon, Sir Edward, Bart.	Glover, Phillips Esq;	Page, Hon. Mrs.
Bowles, Mr. John	Holland, Mr. John	Preis, His Excellency Joachim,
Brooks, Mr. Thomas	Hog, Mr. Roger	derick
Chetwind, William Esq;	Matthew, Capt. William	Reyna Mon, Samuel Esq;
Frederick, Thomas, Esq;	Page, Thomas Esq;	Zolicofre, Theod. William Esq,

*W*HEREAS some Booksellers have declared, that as soon as my Course comes out, they will get it translated into French (as cheap as they can, no doubt,) lest my Book should be spoil'd by an hasty, and perhaps ignorant Translator, I intend to translate it my self, having already done more than half; and if any other Translation appears, I shall write my Name in each Book with my own Hand, that my foreign Auditors may not be imposed upon.



A
COURSE
OF
Experimental PHILOSOPHY.

LECTURE I.



THE Spirit of Disputing, and an earnest Desire to obtain the Victory, rather than come at the Truth, has been one of the greatest Impediments to the Improvement of natural Knowledge. And nothing occasions more wrangling, than when different Persons annex different Ideas to the same Words ; for Men can never rightly understand one another, till they mean the same Things by the same Expressions. No Disputes arise in pure Mathematics, because the Definitions of the Terms are premis'd ; and every Body that reads a Proposition has the same Idea of every Part of it : So that when any Man is so weak as to think to bring one Demonstration in opposition to another, he is to be disregarded by every one that wou'd reason like a Philo-

Lect. I. Philosopher : For we may immediately put an End to all Mathematical Controversies, by shewing, either that our Adversary has not stuck to his Definitions, or has not laid down true Premises, or else that he has drawn false Conclusions from the Principles he has laid down ; or at least own, that we do not understand some Part of his Demonstration, and desire him to explain it ; for unless we can shew where the Error or Paralogism is, we must not condemn by the Lump, but acquiesce with him in what he has prov'd.

It is true, that in mix'd Mathematics, where we reason mathematically upon Physical Subjects, we cannot give such just Definitions as the Geometricians or Logicians do : We must be content with Descriptions, and they will be of the same Use as Definitions, provided we are always consistent with our selves, and always mean the same Things by those Terms, that we have once explain'd : For to say, that others have taken the Words we use in a different Sense, can be no valid Objection ; because it may be answer'd, that then they meant not the same Things as we do. Therefore in this Course, when we make use of such Words, as have been variously understood, we shall shew the particular Meaning, in which we wou'd have them taken : And when we are oblig'd to coin new Terms, as it will often happen in the Description of Machines, we shall always explain them the first Time they are mention'd.

1. BY the Word *Matter*, we understand all that has Extension and Resistance : And because all Bodies, whether solid or fluid, are extended and do resist, therefore we say, that all Bodies are made of Matter.

2. THE *Cartesians* wou'd have Matter to consist in Extension alone ; but Extension without Resistance is nothing but mere *Space*. For tho' they affirm that one cannot have an Idea of Extension without Body ; it is contrary to Experience : Since if we take a Cube out of a cubic Box, which it exactly fill'd, we may very easily conceive the Length, Breadth, and Depth of the empty Box ; and it must be a second Idea, that will give us a Notion either of some other Body coming in to fill the Box, or of its Sides coming together by the Pressure of ambient Bodies.

3. THAT

3. THAT Matter is the same in all Bodies, is evident from its Definition; tho' the common acceptation of the Word wou'd give us another Notion; for in our common Discourse we say, *an Instrument of Wood, Brass, or Iron, or of any other Matter*, as if the Difference of Bodies consisted in their different Kind of Matter: Whereas all Matter is *homogeneous*, or of the same Nature in all Bodies, whether solid or fluid, hard or soft, more or less heavy; whether they belong to the Earth, or any other Part of the Universe; as for Example, the Matter of Cork does not essentially differ from the Matter of Flesh, or that of Gold or Diamonds. But the whole Variety of Bodies,* and the different Changes that happen in them, entirely depend upon the Situation, Distance, Magnitude, Figure, Structure, and Cohesion of the Parts that compound them. * Ann. 1.

4. THAT Mercury resists more than Water *, and Water more than Air, is not owing to the one being of a more resisting Matter than the other; but to the greater Number of Particles contain'd in the same Space in the heavier Body; and often to the stronger Cohesion of Parts; and then a lighter may resist more than a heavier Body, to a Force impress'd to separate its Parts; as Wood will resist more than Water, and a Diamond more than Gold. But where there is little or no Cohesion of Parts, even in the subtlest Fluids, we find a Resistance: For Light condens'd by a burning Glass, tho' many thousand Times rarer than Air, has a considerable Resistance, as appears from its separating the Parts of Bodies, so forcibly, and so soon, when they are plac'd in the Focus of the Glass: Nay, and when the Rays of Light are as much dispers'd as they are here on Earth, coming directly from the Sun, they have a sensible impelling Force, as appears by observing, that the Vapour arising from a Comet (which makes its Tail) is always driven toward that Side of the Comet, which is opposite to the Sun; and that happens whether the Comet be going towards, or coming from the Sun, even at Distances equal to, and greater than that of the Earth. And if there be a Medium finer than * Light (as we have Reason to believe from some Phænomena) even that Medium has a Resistance, whereby it refracts, reflects, and bends the Rays of Light near the Surfaces and Sides of Bodies. * Ann. 2.

Lect. I.

* Ann. 4.

5. THAT Quantity, and consequently Matter, is divided *in Infinitum* *, has been several Ways demonstrated by Mathematicians; and one cannot conceive a Particle of Matter ever so small, but what is still divisible; for since it is a Body, it must have a Top, a Bottom, and a Middle, unless we suppose the Top and Bottom to be the same, which is absurd; and if so, we may conceive such a small Particle to be divided in the Middle. But then we are not by such a Division to imagine the Parts to be separated from each other, any more than (when we divide a cubic Space of two Inches into eight cubic Spaces of an Inch,) those new Cubes are to be supposed remov'd from each other, or taken out of the two Inch Cube which contains them.

* Ann. 5.

6. As to the actual Division of Matter * by separating the Parts from each other, it is not possible beyond a certain Degree; because there are *Atomes*, or extremely small Parts, which are call'd the constituent or component Parts of natural Bodies, which the Wise and Almighty Author of Nature did at first create as the original Particles of Matter, from which all corporeal Natures were to arise, that are without Pores, solid, firm, and impenetrable perfectly, passive, and moveable: So that the utmost Mechanical Division that we can arrive at, does only separate some of these first Parts from one another, and alter their Contact; for mix'd and compounded Bodies are destroy'd by such a Separation, and not by breaking the original Particles to Pieces. These primary Particles being perfectly solid, must be much more hard and firm than any Bodies that can be made out of them with Pores or hidden Vacuities interspers'd, that is, so perfectly hard and firm, that they can never be worn away or diminish'd: For 'tis not reasonable to suppose that there should be any Force or Power in the ordinary Course of Nature, that can divide that into several Parts, which God in his first Creation of Things has made *One*. As long therefore as the original Particles remain entire, there may for ever be Bodies made or compos'd of them, which shall have the same Nature and Texture: But if *these* could be *broken*, *worn away* or *diminished*, then the Nature of corporeal Things, which is dependent on these, might be changed. Earth and Water compos'd of either such Particles as have been worn or broken, or of their Fragments, could not have,


at

at this Day, the same Nature and Texture, as that original Lect. I. Earth and Water, which was compos'd of these Particles when they were found and entire. Wherefore that the Nature of Things should last, and their natural Course continue the same; all the Changes made in Bodies must arise only from the various Separations, new Conjunctions and Motions of these original Particles.

7. THESE must be imagin'd of an unconceivable smallness *; * Ann. 6. but by the Union of several of them together, there are made bigger Lumps or Parts of the first Composition (as they are call'd) which have Interstices between them, because the first Particles do not touch in every Part of their Surface; and those Interstices are call'd the Pores of the first Composition. Likewise by the uniting of several of these Lumps there are form'd *Molecules*, or Lumps of the second Composition, which have Pores of the second Composition, larger than the former: And so on to the several Compositions before we come to Bodies of a sensible Magnitude. Hence it follows, that there must be a great deal of Vacuity * interspers'd in all Bodies, according as they are * Ann. 7. made up of fewer or more Compositions; and all Spaces are not equally full of Matter *. This will be plainly shewed by an Ex- * Ann. 8. periment.

EXPERIMENT I. *Plate I. Fig. I.*

8. UPON B the Brass Plate of the Air-pump (which we shall hereafter describe) set a tall Cylindric Glass-receiver A B open at both Ends of about five Inches Diameter and seven or eight Foot high. Let it be made Air-tight by Means of a wet Leather upon the the Plate B, and another upon the Mouth of the Receiver A under the Covering Plate D, to which Plate underneath is screw'd the Machine *Psps* contriv'd for letting fall Bodies *in Vacuo* at the same Instant of Time. For when the Wire W (which slips to and fro thro' the Collar of oil'd Leathers c, that the Air may no way escape) is drawn up by its Hook b, the square horizontal Plate p being brought up to a narrow Part of the Brass Springs s, s causes them to open so as to let the Plate P (moving on an Hinge) fall into a vertical Position; upon which the Bodies that were laid on it drop at the same Moment. THEN

Lect. I.  THEN upon the square Plate P lay a Down Feather, and a Guinea just by the Side of it, (See *Fig. 2.* where the Plate P is in an horizontal Situation, as it rests on the Return of one of the Springs *s*, and the Collar *c* is separated from the covering Plate and Springs, thro' both which it must be screw'd, when on the Receiver) and having exhausted the Air from the Receiver; by pulling up the Wire *w* let the Guinea and Feather drop, which falling both with the same Velocity (as at C) will come to the Bottom at the same Time: But if the Air had not been exhausted, the Guinea would have been at Bottom before the Feather had fall'n a quarter of the Height of the Receiver.

9. BEFORE the Air was drawn out of the Receiver, if it had been perfectly full (tho' then there was much more Vacuum * than Matter) yet it is evident, that there must be a great deal of Vacuity in it, after the Air is pump'd out; because the Resistance is so far diminish'd, that the Feather falls at least four Times faster than it did in the common Air. For whatever fine Air was left in the Receiver; whatever Particles of Light, or whatever subtile Effluvia penetrate the Glass, and get into the Receiver; all that Matter is much less in Quantity than the Air taken out, because the Resistance is diminish'd. For to say, that after Exhaustion the Receiver is as full as before, wou'd be as absurd, as to say, that a Gallon of Beer turn'd into Froth (so as to reach from the Bottom to the Top of the Receiver) fills it as full as eight Gallons of Beer without Froth, which it is capable to contain.

* Ann. 9.

10. GRAVITY may be look'd upon as a Property of Matter, which tho' not essential, is yet universal, and in one Sense inseparable from it; that is, all Parcels of Matter, however modified, (or all Bodies) have a Gravitation or Attraction towards one another; as will be hereafter demonstrated in Respect of heavenly, as well as terrestrial Bodies: The Tendency of heavy Bodies towards the Center of the Earth, being owing to the same Cause, that makes the Sun and Planets tend towards one another. N B. *When we use the Words Gravity, Gravitation, or Attraction; we have a Regard not to the Cause, but to the Effect; namely to that Force, which Bodies have when they are*
carried

carried towards each other, which (at equal distances) is always proportionable to their Quantity of Matter; whether it be occasion'd by the Impulsion of any subtile Fluid, or by any unknown and unmechanical Power concomitant to all Matter. Lect. I.


II. IF all the Matter in the Universe was contain'd in two equal Balls or Spheres, plac'd at some Distance from each other; these Spheres wou'd move towards one another with equal Velocity, so as to meet in the middle of their first Distance. But if the Spheres be suppos'd unequal in any Proportion; they wou'd meet in a Point as much nearer the great Ball, as the great Ball wou'd be bigger than the other.

12. THE Reason why we do not perceive this mutual Attraction in such Bodies as we daily handle, is, that our Earth having infinitely more Matter than those Bodies, attracts them so strongly as to make their mutual Tendency towards each other insensible. So it happens in respect of a Load-Stone and a Piece of Iron; which when let fall at a little Distance from one another, do not appear to move towards each other; tho' we find their Effect sensible when brought near together.

THIS will be illustrated by the following Experiment.

EXPERIMENT 2. *Plate 1. Fig. 3.*

UPON the Table TAB set the two Balls A, B, equal in Bigness and Weight, (for Ex. of two Ounces each) at the Distance of a Foot from the Hole C, and two Foot from each other. Now if the Earth was annihilated, or was remov'd to an infinite Distance, and the Table did not attract, the two Balls wou'd come to each other and meet at C; but Gravity, or the Attraction of the Earth, pressing them against the Table, they remain at rest; but to overcome that Pressure, and make them act as if the Earth was away, let a String of about 30 Inches long be tied to the Ball A, then brought thro' the smooth Hole in the Table C, then round the Pulley under the Table, and so up again thro' the Hole C, and at last fasten'd to the Ball B. Let the Weight P of four Ounces hang from the Center of the Pulley D: Then if you let go both the Balls at the same Time, they will

Lect. I. will come to each other equally fast and meet just over the
 Hole.

13. IF instead of the Ball A, you substitute another, which weighs but one Ounce (whether it be less than the other, or as big, but hollow, or of light Wood) and this Ball A be let go from the Distance of one Foot, and the heavier Ball B from the Distance of six Inches from the Hole (hanging three Ounces at the Pulley) they will both meet again at the Hole, the lightest Ball going thro' twice the Space that the other does.

14. IN the first Case, the Quantity of Matter in the two Balls, which is four Ounces, being consider'd as divided into Two, causes a two Ounce Ball on each Side to move towards the Hole, which consequently move thro' an equal Space in the same Time.

15. IN the second Case, the Quantity of Matter which is three Ounces, being divided into Two, does on one Side cause a Ball of two Ounces to move towards C, and on the other a Ball of one Ounce to move towards the other, which last moves twice as fast, because it has but half the Quantity of Matter, weighing but half.

* Ann. 10. N B. *Any Weight may hang to the Pulley, * provided it be not too light ; because it only serves to overcome that Gravity, which pressing the Balls against the Table, hinders them from moving towards each other ; as they wou'd do, if the Earth did not exist, or was remov'd to an infinite Distance.*

16. IF the Ball B be suppos'd infinitely greater than the Ball A, the Velocity of the Ball A will then become infinitely greater than that of the Ball B, so that B will stand still and A move thro' the whole Distance between A and B ; the whole Quantity of Matter belonging to both, being now wholly attributed to B alone, and A looked upon as a Point without any sensible Quantity of Matter, and consequently unable to move the Ball B by its Attraction. This is applicable to the Earth and all the Bodies about it, which in respect of them is look'd upon as immoveable, whilst they, as so many physical Points, move in the Lines describ'd by their Centers of Gravity, as they fall to the
Ground

Ground, without having any Regard to the Quantity of Matter, which they contain, whereby they attract the Earth towards them. Thus also since the Sun contains almost 230000 Times more Matter than the Earth, this last is look'd upon as a Point describing an Ellipsis about the Sun, call'd the *Magnus Orbis*; whilst the Sun, that attracts the Earth, is consider'd as immovable in one of the *Foci* of that Ellipsis: Or rather the Moon and Earth together may be consider'd as reduc'd to one Point, which is their common Center of Gravity, and which describes the Orbit abovementioned. Lect. I.


17. THE Earth in respect of the Bodies about it, and the Sun in respect of the Planets and Comets, and all the Planets in respect to the Satellites and other circumambient Bodies, do exert a greater or lesser Force of Attraction, according as the Bodies are nearer or farther off; for Gravity, * being a Virtue diffus'd Ann. 11. from an attracting Body every Way in right Lines, decreases in the same Manner as all other Virtues propagated from a Center round about. So Light and Heat become weaker, as we recede from the lucid or hot Body. This Decrease of Virtue is in a reciprocal duplicate Proportion of the Distance; that is, at twice the Distance, the Virtue is four Times weaker, and at three Times the Distance, nine Times weaker, &c. Thus for Example, if the Earth was three Times farther from the Sun, it would be nine Times less attracted, nine Times less enlightn'd, and nine Times less heated than it is at present: In like manner; if it was four Times farther, it would be sixteen Times less affected by those Qualities. So on the contrary, if it was three or four Times nearer than it is now, it wou'd be nine or sixteen Times more affected.

THIS Proportion of Increase or Decrease of Qualities diffus'd every Way, may be illustrated by the following

EXPERIMENT 3. *Plate I. Fig. 4.*

18. TAKE a Candle, so small that its Rays, that are diffus'd every Way, may proceed as it were from one Point (for if the Candle be large, its Light must be made to pass thro' a small Hole in a Pastboard) and if a Cube * of an Inch, as A, be held Ann. 12. up upon a Needle, at the Distance of one Foot from the Candle,

C
dle,

Lect. I.  dle, its Shadow will cover the Surface of a two Inch Cube B, held at the Distance of two Foot from the Candle, which last Surface is four Times larger than that of the first Cube, as appears by applying the first Cube upon the last. Hereby it is evident, that if the first Cube was twice as far from the Candle it would receive but the 4th Part of the Light; and but the 9th Part, if it was three Times as far; because when held at one Foot from the Candle, its Shadow will cover a three Inch Cube held three Times as far. In the same Manner will a Sphere of an Inch Diameter at one Foot from the Candle intercept all the Light, which wou'd fall upon a two Inch Sphere at two Foot, or a three Inch Sphere at three Foot from the Candle, their Surfaces being as 1, 4 and 9, proportionable to the Squares of their Diameters.

19. As it is easier to raise most Bodies from the Ground than to break them in Pieces; that Force by which its Parts cohere, is stronger than its Gravity. That Force, whatever be its Cause, we shall call the *Attraction of Cohesion*. This Attraction is strongest, when the Parts of the Bodies touch one another; but decreases much faster than Gravity, when the Parts that were before in contact, cease to touch; and when they come to be at any sensible Distance, this Attraction of Cohesion becomes almost insensible.


THIS Attraction is always the strongest, where the Contact is the greatest. As for Example, two Boards of Fir or Oak, being glued in the Middle along the Grain, are not so easily broken asunder in the glued Place, as any where else; because there are more Pores, and consequently fewer touching Parts, along the Wood any where else, than where the Glue is; for when a Joint is shot (as the Workmen call it) or the two Pieces of Wood made smooth, in order to join them, the Glue which is spread on the Pieces fills the Pores; and causes the Wood not only to touch where it did before, but even in the Interstices where it did not touch; because those little Spaces are fill'd with Glue, that supplies the Place of Wood. On the contrary, when the Wood is more solid, * or has fewer Pores than the Glue, it does not hold so fast where it is glued, as in the other Parts of the Wood; which may be seen in Brazil-wood, Ebony, or Lignum-vitæ, and in Metals. The Parts of Glass, which are almost round, touching but in a few Points

Points, are easily separated, and therefore it breaks easily : And Lect. I
 Fluids (whose Parts seem to be Spherical) have scarce any Co-
 hesion, except so much as serves to make Drops, whose Roundness
 plainly proves an Attraction of Cohesion: For if this Roundness
 depended upon the Pressure of the Air, the Drops wou'd not be
 round *in Vacuo*; and if it depended upon the external Pres-
 sure of any other Fluid whatever, two Drops cou'd never unite
 into one; because the Figure of any Portion of a Fluid * press'd * Ann. 14.
 every Way, by the same or any other Fluid, cannot be alter'd by
 that Pressure; whereas from a mutual Attraction of the little
 Globules * that compose the Drop, it must become round; and it * Ann. 15.
 will continue so, because then there will be the greatest Contact
 possible between all the Parts. A Drop of Water, or any o-
 ther Liquor, does indeed become flat where it touches a Table,
 or any plane Body, that does not repel it; but this is owing to
 the Attraction of the Table, and to the Gravity of the Drop, if
 the Table be horizontal.

How such an Encrease of Contact increases this Attraction,
 is more evident by the following

EXPERIMENT 4. *Plate I. Fig. 5.*

20. HAVING moisten'd or thinly smear'd over with Oil of
 Oranges * two Glass Planes A B C D, (18 Inches long, and three * Ann. 16.
 or four Inches broad) lay them upon one another in an horizon-
 tal Position in the wooden Frame H I L M, having first put a
 Drop of the same Oil upon the undermost Plane at G, and laid
 a thin Plate or Piece of Money upon the said Plane, between
 D C, to hinder the upper Plane from touching it at that End
 whilst their other Ends A B are in close Contact. The Drop
 being large enough to reach the upper Plane, will immediately
 be flatten'd and move on towards the touching Ends, continu-
 ally increasing its Diameter, as at Q and R, and likewise mov-
 ing faster as it spreads. Nay tho' the Planes be rais'd up at
 their touching Ends by means of the Prop O N, the Drop will
 continue to advance towards the touching Ends, but not so
 fast. When the Drop is at G, a small Elevation of the Planes
 will retard it; if they be rais'd a little higher, it will stop; if
 still higher, the Gravity of the Drop will act more strongly
 than the Attraction of the Planes, and make the Drop move

 Lect. I. downwards towards C D. When the Drop is at Q, the Plane will require a much greater Elevation to stop the Drop, or make it move downwards; and still a greater when it is at R. Now when the Planes are rais'd so high as to make the Drop run back; if the upper Plane be press'd by the Finger a little above the Drop so as to come nearer to the lower Plane; the Drop will not only stop, but move upwards; because it spreads more by that Pressure and touches so many more parts of the Glass than it wou'd have done, that even at this Elevation of the Planes, their Attraction overcomes the Gravity of the Drop. That it moves at first towards A B, is owing to the greater Attraction of the Planes at *e*, where they are nearer together than at *f*; and that the Velocity of the Drop encreases, is owing to its moving towards A B, where the Attraction must encrease continually * as the Planes come nearer and nearer.

* Ann. 17.

SEVERAL other Circumstances of the Attraction of Cohesion are shewn by the following Experiments.

EXPERIMENT 5. *Plate 1. Fig. 6.*

21. HAVING fix'd into a piece of Wood or Cork C C, several little Glass Tubes, the Diameters of whose Bores are less than one another, (the biggest being about $\frac{1}{4}$ of an Inch) let the Ends of those Tubes be dipp'd into any ting'd Liquor that will adhere to Glass, as red Water; and the Liquor will spontaneously rise in all of them, but always highest in those whose Diameter is least; as appears at 1, 2, 3, 4, 5, when they are dipp'd in the red Water of the Vessel A B.

IN very small capillary Tubes, the Liquor will rise very high, but then the Colour will be imperceptible; But to make the rising Liquor visible in that case, the Tube may be applied to ones Finger after it is prick'd so as to make a drop of Blood issue out, which will rise very quick and be visible tho' the Tube be as small as an Hair. See *Plate 2. Fig. 1.*

ANY porous Body will have the Effect of capillary Tubes; thus Water will rise up into a piece of Bread, or into a piece of Loaf Sugar whose lower end is dipp'd into the Liquor; but it will rise much higher into Loaf Sugar, because its Interstices are smaller than those of the Bread.



EXPERIMENT. 6. *Plate 2. Fig. 2.*

TAKE two square Planes of Glass A B C D and having moisten'd them with Water, set them upright in a Vessel of the same Water, pressing the Sides together at A B, but keeping the opposite Sides D C asunder by putting a thin Plate between them. The Water will rise between the Planes in the Curve *efg*, where it is to be observ'd that the Fluid always goes highest where the Planes are nearest, namely towards A B, just as it happens in the small Glass Tubes.


THAT none of these Phænomena arise from the Pressure of the Air, is evident by making the Experiments *in Vacuo*; which may be easily perform'd by means of the Wire *w* (*Plate 1. Fig. 2.*) which can move up and down thro' its Collar of Leathers without admitting the external Air into the Receiver, as it rises or depresses the Planes or the Tubes fasten'd to it. *The whole Apparatus for this will be hereafter described.*

EXPERIMENT. 7. *Plate 2. Fig. 3.*

23. IN a clean Glass that is not full, Liquors will be higher at A B where they touch the Glass than in the middle C; but that Elevation is hardly sensible but near the Sides, because the Attraction of Cohesion reaches but a little way. Quicksilver does the reverse in this case; and in small Tubes it does not rise so high as the Surface of the rest of the Quick-silver in the Vessel in which the Tubes are held. The Reason of these Phænomena * is, that Water is attracted by Glass * Ann. 18. more than it is attracted by it self; and that on the contrary, Quicksilver attracts Quicksilver more than Glass attracts it. ** Ann. 19.

EXPERIMENT 8. *Plate 2. Fig. 4.*

LET A B be a Cylandrick Glass Vessel fill'd up to the Line A B with Quicksilver, whose Surface is lowest at A and B the Sides of the Glass, where it rises with a Convexity. If an open Tube of a small bore D E be press'd against the inside of the Glass (to render the Experiment visible) its lower Orifice

 Lect. I. Orifice being almost at the Bottom of the Glass, the Mercury will rise in the said Tube no higher than F below the Surface A B of the Mercury in the Vessel. But to avoid all Cavils about something in the Tube that might be suppos'd to stop the Mercury, let the Experiment be made in this Manner: Let the Tube being quite full of Mercury, (and kept so by the Finger press'd on the upper Orifice D,) be put into the Vessel, whose bottom is a little convex, below the Surface of the Mercury as low as E; upon the removal of the Finger, the Mercury in the Tube will fall below F, and then rise above it, and after some Vibrations settle at F, the Point to which it rose before.

EXPERIMENT 9. *Plate 2. Fig. 5.*

24. INTO the Cylandrick Glass Jar A B of about 5 Inches Diameter and $1 \frac{1}{2}$ Inch deep pour gently about one Pound of Quicksilver, and a circular part of the Bottom as C C will remain uncover'd; then if you shake the Jar to make the Quicksilver come together, the whole Bottom will be cover'd; but if without any shake you continue to pour on more Mercury you may put in a Pound or two more before the Bottom be quite cover'd, the Pit C C becoming continually less, but deeper; and if then you bring the Mercury together to cover the bottom and destroy the Pit; a Finger being press'd against the Bottom thro' the Mercury, will, when taken away, leave a Pit as before, the Mercury there remaining convex at the Sides of the Pit; as appears in the vertical Section of the Glass and Quicksilver in the lower Figure 5. *Plate 2.*

EXPERIMENT 10. *Plate 2. Fig. 6.*

25. HAVING put in still more Mercury 'till there be no Pit left, lay a piece of Iron Wire C C about two or three Inches long and $\frac{1}{4}$ of an Inch thick upon the Surface of the Mercury, where it will swim making a dent on both Sides as at D; which happens because the Mercury attracts itself more than it does the Iron; so in the case of *Fig. 5.* it was less attracted by the Glass than by it self, and therefore made the Pit; but when once the Sides of the mercurial Pit C C were brought to touch, they never parted again of themselves.

Ex-



EXPERIMENT II. *Plate 2. Fig. 7.*

26. IF *mm* represents the Surface of the Mercury, and the Wire which swam at the Top be press'd down to the Bottom, (where *D* represents its transverse Section,) the Wedges of Mercury *bca*, that go under the Wire so as very nearly to meet at *c*, will not remain in their places, but by the rest of the Mercury they will be attracted towards *d, d*, so as to leave the Spaces *bca* without any Mercury; as appears in the Section of the Glass, Wire, and Mercury, *Fig. 8*. And that it is so in fact, appears by looking at the under side of the Glass in the upper *Fig. 8*. where the Wire or Wires (if there be more than one) become visible thro' the Bottom of the Glass, which cou'd not be unless the Mercury went away from under them; because a Cylinder can touch a Plane but in an invisible Line: And as a farther Proof of this; the Wires, tho' specifically lighter than Mercury, remain at bottom, as being press'd only downwards by the Arch of Mercury over them; which cou'd not happen if the Mercury cou'd insinuate it self under.

27. Now if the Experiment be made with a Silver Wire of the same bigness as the Iron one, when it is laid at top, the Mercury will rise up round it as at *aa*, *Fig. 9*. and this Wire will not remain at the Bottom of the Glass tho' push'd down to it, but always emerge, neither does it become visible thro' the Bottom of the Glass, tho' held down against it with the Finger; nay even when the Finger is seen on each side of it. This happens because Silver attracts Mercury more than Mercury attracts it self. But to shew that this Attraction is only strong in Contact or extremely near; let the Silver Wire be made a little foul by putting it into the Fire, and then the same thing will happen to it as to the Iron Wire; because the Attraction of Cohesion is insensible, at the Distance of the Thickness of the thin Skin that there covers the Silver.

28. THAT there shou'd be a Contact to bring Bodies to cohere very much, is evident from the foldering of Lead or Brass with a Mixture of Lead and Tin, commonly call'd soft Solder.

Lect. I. Solder. For if these Metals be not well clean'd, they can never be folder'd perfectly tight; and in Lead, after the Parts that are to be join'd have been made foul by rubbing them over with Chalk, and then with Mallows or any Green-Herb, so as to make a thin Skin; the Work-Man scrapes clean the two Edges of the Lead that are to be join'd, that the Solder may come close enough to the Metal; and when the Solder is pour'd on hot out of a Ladle, then spread with an Iron, it fastens strongly where the Metal was clean'd, but does not at all stick where the Skin remains that was made by the Chalk and Juice of the Mallows. It has been observ'd that the Air alone will foul or make a Skin over the part of the Lead that has been clean'd, and therefore Grease or Tallow is commonly rubb'd on after scraping; for the Parts of any fat or inflammable Substances being much finer than those of Air, will allow the Solder to come much closer to the Lead than a Plate of Air, or what separates from the Air to stick to the Lead.

29. IN what Proportion of the Distance this Attraction of Cohesion encreases or decreases is not yet fully known; but from some Phænomena, we have Reason to believe that it decreases in a biquadratic Ratio of the encreas'd Distance; that is, at twice the Distance it acts 16 times more weakly, and at 3 times the Distance 81 times more weakly, &c. For it becomes insensible at the least sensible Distance.

30. THERE is in Nature another sort of Attraction not so strong as the Attraction of Cohesion, but stronger than Gravity; whose Proportion of Decrease, in the Removal of Bodies Attracting, is nearly as the Cube and a quarter of the encreas'd Distance: * And this is the Magnetical Attraction. As for Example, if a Loadstone attracts a piece of Iron with a certain Force at a given Distance; at twice the Distance, the Attraction will be 10 Times weaker: And at three Times the Distance, the Attraction will be $33\frac{3}{4}$ times weaker. But as Magnetism is a particular Virtue that affects only Load-stones and Iron and Steel, we shall refer a fuller Account of it to another Place; because we are now only considering general Properties of Bodies. We shall only observe, that the Load-stone repels as well as attracts; for that Pole of the Stone, which attracts one End of a touch'd Needle, will repel the other.

* Ann. 20.

31. THERE

31. THERE are several other Instances in Nature of a repellent Power in Bodies *; and very often the same Bodies that attract one another at certain Distances, and under some Circumstances, do repel one another at different Distances, and under other Circumstances. * Annot. 21

THIS may be seen upon the Dissolution of Salts in Water. That the Parts of the Salts attract one another, appears from uniting into hard Lumps when the Water is evaporated, so that the Particles come so near to each other, as to be within the Power of the Attraction. That they repel one another at farther Distances, appears from the regular Figures into which they coalesce, when by the Evaporation of part of the Fluid in which they float, they are brought within each others Sphere of Attraction; these regular Figures depending entirely upon the Equality of their Distances one from another before this Evaporation, and this Equality of Distance being owing to an Equality of repelling Force.

32. A repelling Force is also prov'd by the Production of Air and Vapours; for those Particles which are forc'd out of Bodies by Heat and Fermentation, as soon as they are out of the Sphere of Attraction of the Bodies, do immediately recede from them, and from one another with a great Force, and avoid coming together again; so as sometimes to take up above a Million of Times the Space which they did before in a dense Body.

33. THE Attraction and Repulsion in the same Body at a considerable Distance, is evident in several electrical Experiments.


IF you rub a Piece of Amber with a dry Hand or a woollen Cloth, it will put into motion Threads, Feathers and light Bodies, attracting and repelling them at a Distance: And therefore the Name of *Electricity* has been given to that attracting and repelling Force which is excited in any other Body by the same Friction as is given to the Amber. Wax, Rosin, Sulphur, Silks, Paper, Ribbons, Hair, and Feathers, and several other Bodies have this Property: But Glass has it more than any other.

EXPERIMENT 12. *Plate 2. Fig. 10.*

34. TAKE a Glass Tube of about $1\frac{1}{2}$ Inch Diameter, and rubbing it from one End to the other with a dry Hand pretty briskly, it will attract a Feather, or any light Body, at a considerable Distance, from one to eight or ten Feet. After a Feather has been attracted and stuck to the Tube for some time, it will fly off of it self, and never come to the Tube again (which constantly repels the Feather in the Air, whenever it is brought near it) till it has touch'd some other Body; as a Finger or a Stick. And if the Finger be held pretty near the Tube, the Feather will alternately fly from the Finger to the Tube, always stretching out its Fibres the way that it is going, and that before it comes off from the Finger or the Tube. *In driving the Feather about the Room with the Tube, it must now and then be rubb'd afresh to excite the Electricity, which continually grows weaker, after the Friction is over.*

EXPERIMENT 13. *Plate 2. Fig. 11.*

35. IF several little Pieces of Leaf-gold, or Leaf-brass be laid upon a stand or small Table, and the rubb'd Tube be held over them at the Distance of a Foot or two; the Pieces of Leaf-gold will move from the Table to the Tube with great Swiftnefs: And often by the Attraction and Repulsion they will move backwards and forwards without touching either the Tube or the Table. But if two Books, or two Pieces of Wood of the same Size be set up on End on the Table on each side of the Leaf-gold, as at A, B; (*Plate 2. Fig. 12.*) so that their Distance A B be equal to the Height of one of them: Then the Tube being held between their Tops, as at D, will have no Power to move the Leaf-gold, tho' the Distance from the Leaf-gold be but six Inches, when the Tube just before attracted the Leaf-gold at a Foot or two; but if the Pieces of Wood be remov'd, without giving the Tube a new Friction, it will attract and repel the Leaf-gold as before. When the Pieces of Wood are not remov'd, the Tube will not put the Leaf-gold in Motion, till the Distance D C from the Tube to the Gold be less than half the Distance A B of the Pieces of Wood: As if this Effect could not be produc'd while the Sphere of Attraction represen-

represented by the Circle E C F (whose Center is in the middle of the Section of the Tube at D) is disturb'd by the Pieces A and B. 

EXPERIMENT 14. *Plate 2. Fig. 13.*

36. To know when the Tube is sufficiently rubb'd to make Experiments with it, you must move your Fingers ends nimbly by the Tube, as if you went to strike it in a Direction perpendicular to it's Axis, but the Fingers need not come nearer than a quarter of an Inch from the Tube: Then the Effluvia or fine Parts that fly from the Tube will snap against the Finger, or (beating back from it) against the Tube, so as to be heard with a Noise like the crackling of a green Leaf in the Fire; but not so loud.

ACCORDING to the State of the Air, the Tube requires more or less rubbing. In Weather that is hot and moist, the Tube requires a great deal of Friction before it will snap, and attract and repel the most strongly; and then at best it's Virtue will diffuse it self but a little Way: So that the same Tube, which in dry cold Weather gave Motion to the Fibres of a Feather at the Distance of eight or ten Feet, will scarce act sensibly at the Distance of two Feet when it rains in Winter.

IF the Tube be warm'd at the Fire without rubbing, it will have no Effect: It's Electricity will also be less if you rub the Tube long enough to make it very warm; and then you must let it cool before you use it again.

IT is not amiss to set the Tube or any Glas to be rubb'd before the Fire to dry it before you use it, especially if it be pretty thick, provided it is not much heated.

37. IT is remarkable, that if the Tube be rubb'd in the dark, the Effluvia will appear lucid; and when it is made to snap (as in the 13th Fig. of *Plate 2.*) there appears a Light upon the Fingers Ends, as at A; and if a little Brush be held near the Tube, as at B; or drawn along it without touching, just after it is rubb'd, Sparks of Light like Stars will appear upon every Hair of the Brush; but the same part of the Tube will not snap or give Light twice together in the same Place * without a new Friction.

* Annot. 22.

EXPERIMENT 15. *Plate 2. Fig. 14.*

IF the rubb'd Tube be brought near a Down Feather tied to the upper part of a little Stick standing upon a Foot ; the Feather will stretch out its Fibres towards the Tube ; but upon the Approach of a Finger between the Tube and Feather, they will be repell'd by the Finger ; tho' they will be again attracted by it as soon as the Tube is remov'd ; and then the Fibres will turn back again to the Stick and be attracted by it, when the Finger is taken away.

EXPERIMENT 16. *Plate 2. Fig. 15.*

39. IF you set a Glass Receiver, about five Inches wide, and twenty Inches high, over the Stick and Feather, having first dried the Receiver at the Fire or in the Sun ; upon rubbing the Glass with one or both Hands from Top to Bottom, the Feather will stretch its Fibres every way like the Radii of a Sphere, when the Hand is remov'd from the Receiver. But if whilst you rub the Receiver, or after rubbing, you only move the Hand upwards and downwards, the Fibres of the Feather will (notwithstanding the Interposition of the Glass) follow the Motion of the Hand : And if the Tube be rubb'd within a Foot or two of the Receiver, the Feather in the Receiver will likewise follow the Motion of the Hand rubbing the Tube. When the Tube has its Electricity excited by Friction, if it be brought near the outside of the Receiver, the Feather will stretch its Fibres towards the Tube ; and upon the Removal of the Tube turn back to the Stick ; though sometimes this last Phenomenon will happen at the approach of the Tube, and the Fibres will stretch out again when the Tube is taken away : Nay, sometimes there seem to be Fits of Attraction and Repulsion ; for whilst the Tube is held near the outside of the Receiver, the Fibres of the Feather will be alternately extended and contracted, without any new Friction given to either of them.

40. JUST after the Receiver has been rubb'd, if you blow towards the Feather, (See the 15th *Fig. Plate 2.*) its Fibres will fly from the Blast ; and they will also fly from an Hand mov'd briskly

briskly towards the Glas, yet so as not to touch it ; but the Ex-Lect. I. Experiment will not do twice without rubbing the Receiver a-new.

41. Most of these Experiments, if not all, will succeed even when the Air has been pump'd out of the Receiver : Only there will be this difference, upon rubbing it *in Vacuo*, that the Light excited will be of a Purple Colour, in a much greater Quantity, and all within the Glas : And whereas Bodies would be attracted before, when held near the outside of the Glas, now that Power will cease, and the Virtue will exert it self wholly inward. The same will happen to the Tube when exhausted, as also to a Glas Globe * whirl'd by Means of a Wheel, and so rubb'd by the Hand ; * Annot. 23. as is described more at large by the late Mr. *Hawksbee* in his Book of Experiments, where he has given a large Account of a great many electrical Experiments that he made.

42. I shall say no more on this Subject now ; because I shall have Occasion to consider it more fully in another Part of my Course : And the Intent of this Lecture is only to shew,—That those Properties of Bodies, such as Gravity, Attractions, and Repulsions, by which we shall hereafter explain several Phænomena, are not occult Qualities or supposed Virtues, but do really exist, and are by Experiments and Observations made the Objects of our Senses. These Properties produce Effects, according to settled Laws, always acting in the same Manner under the same Circumstances : And, tho' the Causes of those Causes * are not known, since we do not reason about these hidden Causes ; it is plain that we reject occult Qualities, instead of admitting them in our Philosophy, as the *Cartesians* always object to us. * Annot. 24.





ANNOTATIONS upon the First LECTURE.

I. [3—*But the whole Variety, &c.*]

Annotat.
Lect. I.

IF we consider the Bricks of which a Building consists, as its smallest or first Parts; we shall find that, however similar they are, their different Disposition, in respect to each other, will produce very different Parts of the Edifice: An Arch, a Wall, a Chimney, a Peer, a square or a round Pillar, a Globe or a Cube are compos'd of the same Sort of Bricks; and such as served for one Part, when pulled asunder, will as readily serve for another. So in the wonderful Edifice of the Universe, there is no need of a Difference in the Atoms or first Particles, of which the several Parts are compounded: The same Atoms being as proper to make Land as Sea, to make Gold as Clay: And, when we do not consider the Soul that actuates the Matter, a particular Disposition of the first Atoms makes all the Difference betwixt a lifeless Lump, and the Body of a curiously organiz'd Animal.

One may bring various Examples of Matter trac'd through several Bodies, whose Changes depend upon the different Texture and Position of the Parts.

When the Water of Rivers, Seas, and Lakes, is so rarified by the Heat of the Sun, as to become specifically lighter than Air (which will happen, when it takes up above 900 times the Space in Vapour that it did in Water) it will rise up so high as to form Clouds of various Colours, which float about at that Height, where the Air is of the same specifick Gravity as the Clouds.

When the Winds, by carrying off some of the Air above, cause that which is below to become specifically lighter by its Expansion; the Clouds, retaining the same specifick Gravity as they had before, do then descend, and, by the Resistance which they meet with in their Descent, are chang'd into Rain, which falling down to the Earth, does in a great Measure run back into the Rivers and Seas; but some Part of it runs into the Earth, and is imbibed by the Seed of Plants. If we consider so much of it as gets into the Grains of Wheat that are sown, the Appearance of it is much chang'd

chang'd in the green Blade of Corn, then in the Straw, afterwards in Annotat. the Ear, and in the Grains contain'd in it. The Wheat, by being ground *Lect. I.* in the Mill, puts on the Form of Flour; which being made up into Paste and then baked, is again changed into the Crust and Crumb of Bread. The nourishing Parts of the Bread (after it has been eaten by Man, and has pass'd thro' the Stomach) do out of the Intestines pass thro' the Lacteal Vessels into the *Receptaculum Chyli*, and thence up the *Ductus Thoracicus* into the left Subclavian Vein, or (as an ingenious * Anatomist has lately *Mr. St. An-* discovered) into the left internal Jugular Vein; where mixing with the *dr.* Blood, it goes along with it thro' the Heart and Lungs, where it receives vivifying Particles from the Air, and returning into the Heart, is from thence by the Action of the left Ventricle of the Heart, and the Arteries carried to the Extremities of the Body. There some Part of this new Blood circulates back again; whilst other Parts are changed into the Substance of Bones, some into Membranes, some into Hairs, some into Nails; and other Parts of it passing thro' the Glands, are turn'd into Sweat and so become Water again, as at first.

If we had consider'd such Drops of Rain as impregnated Linseed; we might have trac'd it thro' the Stalks of the Plant, Flax made of that, Thread spun from the Flax, Linnen made of the Thread, a white Pulp made of the Rags of the worn Linnen beaten up with Water at the Paper-Mills, Paper made of that Pulp thinly spread upon a fine Net-Work of Wire; and lastly the Smoke, which burn'd Paper affords, is again easily reduc'd to Water.

Solids become Fluids, as Metals do by the Action of Fire, or being dissolv'd in acid *Menstruums*; and Fluids put on the Form of Solids, as Mercury will be made hard by the Fumes of Lead: And two Chymical Liquors will immediately upon their Mixture coalesce into a firm Substance.

It is to their particular Figure that Machines and Instruments owe their Usefulness; Clocks, Mills and other mechanical Engines being only valuable, when their Parts that are made to communicate Motion have their proper fitness.

The same kind of Glass produces great Variety of Effects, according to the different Figure of its Surface; as is known to all that have seen Optic Glasses.

The Difference of Seasons which makes such changes on the Surface of our Earth, and even in the Bodies of Animals, is entirely owing to the different Situation of the several Parts of the Earth in Respect to the Sun.

Those that would read more upon this Subject may consult *Dr. John Keil's Introductio ad veram Physicam. Edit. 3. Lect. 7.*

Annotat.
Lect. I.

2. [4. *That Mercury resists, &c.*] Mercury weighs $13\frac{2}{3}$ Times more than Water, and is found to resist just $13\frac{2}{3}$ Times more; and Water which weighs between 800 and 900 Times more than Air, is found to resist just so much more. This Sir *Isaac Newton* found by making Experiments upon Pendulums of Wood and Lead in the Air, Pendulums of Lead in Water, and of Iron in Mercury. See his *Principia*, Edition 2d. Book 2. Prop. 31. and Prop. 40. where he demonstrates that the Resistance of Fluids is as their Quantity of Matter.

I made an Experiment before the *Royal Society* about two Years ago with a Ball of Gold of an Inch Diameter; which, being suspended by a String, did first oscillate in Water and then in Mercury: And it appear'd that 42 Vibrations in Water destroy'd as much of the Motion of the Golden Pendulum as three Vibrations in Quicksilver. And letting the Ball of Gold fall in a Copper Tube four Foot long, and $4\frac{1}{2}$ Inches Diameter, filled with Mercury (from an Height of three Foot ten Inches) and nicely observing the Time of its Fall, I found by comparing several Experiments, that the Resistance of the Medium (discovered by the Method taught in the said 40th Prop. of the *Principia*) agreed so exactly with Sir *Isaac Newton's* Theory, as not to differ one Tenth of an Inch in the Space that the Ball fell thro', which was three Foot ten Inches.

3 [4—*And if there be a Medium finer than Light, &c.*] See Sir *Isaac Newton's* Optics, second Edition, Book 3d. Queries 18, 19, 20 and 21.


4. [5—*That Quantity, &c. is divisible in infinitum, &c.*] Of the many Demonstrations brought to prove this Assertion, I shall only mention two that are very plain and obvious. The first taken from Dr. *s'Gravesande's* Introduction to Sir *Isaac Newton's* Philosophy, Part 1st, No. 18.

Plate 3. Fig. 1.

Let there be a Line AD perpendicular to BF, and another as GH at a small Distance from A, also perpendicular to the same Line. With the several Centers C, C, C, &c. and Distances CA, CA, &c. describe Circles cutting the Line GH in the Points e, e, &c. The greater the Radius AC is, the less is the part eG. Since the Radius may be augmented in infinitum, therefore the Part eG may be diminish'd in the same Manner; and yet it can never be reduc'd to nothing, because the Circle can never coincide with the right Line BF.

The next is from Dr. John Keil's *Introductio ad veram Physf.* Lec. 3. De Magnitudinum Divisibilitate.

Let AB (Plate 3. Fig. 2.) be a Perpendicular between the Parallels CD, EF. From the Point C in one of the Parallels draw the Line CG to a Point G on the other Parallel on the other side of the Perpendicular AB,

A B, and it will divide the said Perpendicular into two Parts at K: Ano- Annotat. ther Line drawn from C to H will divide the Part K A (of A B) into Lect. I. two Parts; and since upon the Line E F, which may be produc'd in *Infini-*  *tum*, other Points as I, &c. may be taken, new Lines may still be drawn to divide the remaining Part of A B. For whatever Line is drawn from C to any Point of the Line E F, however distant, it can never coincide with the Line C D; and therefore it will still divide the Part of A B that remains after the foregoing Division.

For other and more ample Demonstrations, See the said *Lect.* 3d. where he also shews the Absurdity of the contrary Opinion: And in the next *Lecture* he removes the Objections alledg'd against the Divisibility of Quantity; by demonstrating those very Assertions to be true, which the Objectors alledge as absurd Consequences of the Divisibility of Magnitude in *Infinitem*.

As First, *That a finite Quantity may have an infinite Number of Parts.* For if the Line A B* be divided into an Hundred Parts, all those Parts ta-* Plate 3. ken together will be equal to A B; and if it be divided into a Thousand Fig. 3. Parts, all those Parts taken together will still be equal to A B. Now, a Thousand may be encreased in any Proportion, nay, may receive an infinite Addition of Numbers; and since, however the Number of Parts is encreased, the Sum of them can never exceed the Line A B; the said Line may, without an Absurdity, be said to contain an infinite Number of Parts.

Secondly, *That it implies no Contradiction, but is agreeable to Geometry, to say, that a Finite may be equal to an Infinite*; shewing that a finite Space is equal to an infinite one, and that an infinite Solid is equal to a finite one.

Thirdly, *That there are Infinites, whose Magnitudes bear certain Proportions to one another, and that some are bigger, (nay infinitely bigger) than others.*

In the Circle A B F † take an Arc B F infinitely small, then the Chord † Plate 3. B F will be infinitely smaller than the Diameter A B; and yet it will be Fig. 4. infinitely greater than the vers'd Sine B G, found by drawing F' G perpendicular to B A. Likewise in the Circle B F A* if the Arc B F be taken* Plate 3. infinitely small, B E be its Tangent, F' G the Right Sine, B G the vers'd Fig. 5. Sine, and F H equal and parallel to the vers'd Sine; it is demonstrable, that C B is infinitely greater than B E, and B E infinitely greater than B G, and B G than H E.

But the most surprizing thing of this Kind, is, that if A E and A B † be drawn at right Angles, and the Parabolic Curves of different † Plate 3. kinds, C, D, G, H, be drawn thro' the Point A; the Angle of Contact Fig. 6. F A C, which is infinitely less than any rectilinear Angle, will be infinitely greater than the Angle F A D, and F A G will be infinitely less than F A D: And so there may be an infinite Series of Angles of Contact going on infinitely, of which every following one is infinitely greater than the former: Nay, between any two Angles, there may be inserted

E

innumerable

Annotat. innumerable Angles infinitely greater than each other. And even between
 Lect. I. any two of these Angles, there may be a Series of intermediate Angles
 going on *in Infinitum*; of which every following one is infinitely greater
 than the former. And thus Nature knows no Bounds. See the *Demon-*
strations of these Propositions in the same Book, Lect. 4.

5.] 6— *There are Atoms, &c.*] Dr. Keil in his Introduction, *Lect. 51.*
 deduces the two following *Theorems* from the infinite Divisibility of
 Matter.

THEOREM 1.

“ Any Quantity of Matter, how small soever, and any finite Space, how
 “ great soever, being given; (as for Example, a Cube circumscrib’d about
 “ the Sphere of *Saturn*) it is possible for the Matter of that small Sand
 “ to be diffus’d throughout all that Space, and to fill it so, that there shall
 “ be no Pore or Interstice in it, whose Diameter shall exceed a given
 “ Line.

From the Demonstration of this *Theorem*, he draws the following *Corol-*
lary, which serves for the Demonstration of the other.

Cor. “ Hence there may be given a Body, whose Matter, if it be re-
 “ reduc’d into a Space absolutely full; that Space may be any given Part
 “ of its former Magnitude.

THEOREM 2.

“ There may be two Bodies of equal Bulk, whose Quantities of Mat-
 “ ter being unequal in any Proportion; yet the Sum of their Pores, or the
 “ void Spaces in the two Bodies, shall be almost equal.

The Doctor applies his Demonstration to an Inch Cube of Gold, and an
 Inch Cube of Air; the Substance of which is as follows.

* Plate 3.
 Fig. 7.

The Cubic Inch of Gold, * A contains near 20000 Times more
 Matter than the Cubic Inch of Air B, but we will only suppose
 it to contain 10000 Times more. Now let the Matter in A be re-
 duced into a Space absolutely full, which we will suppose equal to the

$\frac{1}{100\ 000}$ Part of a Cubic Inch; (which may be done by the *Cor.* of the last
Theorem. Then if the Matter in B be reduc’d to a Space absolutely full,

it will only take up the $\frac{1}{1\ 000\ 000\ 000}$ Part of a Cubic Inch, because B
 contained 10000 Times less Matter than A. To compare these two full
 Spaces let us make the Denominators of the two Fractions the same;
 and

and $\frac{10000}{1\ 000\ 000\ 000}$ will exprefs the folid Space in the Inch of Gold : There- Annotat.
Lect. I.

fore the $\frac{999\ 990\ 000}{1\ 000\ 000\ 000}$ remaining Parts of that Cubic Inch will be void

Spaces ; whilst the void Spaces in the Inch of Air, (after its Matter has

been reduc'd to a folid) are exprefs'd by the Fraction $\frac{999\ 999\ 999}{1\ 000\ 000\ 000}$: And

confequently fince the Numbers 999 990 000 and 999 999 999 are almoft in the Ratio of Equality ; the void Spaces in both Bodies are nearly equal.

Though thefe *Theorems* feem inconfiftent with the Doctrine of Atoms, they do not overthrow it ; becaufe, as they are deduced from the Divifibility of Quantity, they are rather Mathematical than Phyfical. For though an Atom may be conceiv'd of an exceeding Smallnefs, yet its Diameter muft be of a determinate length, and confequently too big to answer the Conditions of the firft *Theorem*, which fupposes no firft Parts. But then even in an Atom (or firft Phyfical Part) Mathematicians may affign Parts fmaller in any Proportion, fo as to agree with the forementioned *Theorems* : For thofe Particles of natural Bodies, which cannot be divided in the Operations of Nature, do virtually contain an infinite Number of Parts ; tho' thofe Parts are never feparated from one another.

Notwithftanding, that giving Atoms a certain Bignefs limits Dr. Keil's *Theorems* in the Phyfical Senfe ; yet it appears from *Phænomena*, that there actually exift Particles fmaller enough to agree with the firft *Theorem*, if a Grain of Sand be made ufe of to fill the Sphere of *Saturn* ; and with the fecond *Theorem*, if an Inch of Gold be compar'd with an Inch of Air ; only fupposing (which is more than probable) that a Body perfectly folid is as much denfer than Gold, as Gold is denfer than Light, or Æther which is ftill rarer than Light.

And if a Grain of Sand be fuppos'd divided into fo many Parts, as to fill the Sphere of *Saturn*, without having any Pore bigger in Diameter than an Hair ; thofe Particles may be ftill bigger than thofe of Æther, if not bigger than thofe of Light.

6. [7 — *Thefe are to be imagin'd of an unconceivable Smallnefs, &c.*] Though it be furprizing to think, that Matter fhould be actually divided into Parts fo fmaller, as we have mentioned in the foregoing Note ; a few Inftances of the fmaller Parts into which it is divided by Art, or the Work of Mens Hands ; and fome Examples of its Subtility, as it is naturally difpers'd all over the Univerfe, will make the Affertion very plain to any Body that will afford the leaft Attention.

The Gold-Beaters, even with courfe Tools, reduce that Metal to fuch a Thinnefs, that fifty fquare Inches of Leaf-Gold weigh but one Grain. Now the Length of an Inch may be divided into 200 vifible Parts, as appears in

Annotat. *Plate 3. Fig. 8.* where the 20th Part of an Inch is distinguish'd into ten
 Lect. I. visible Parts by six black Strokes and five white Interstices*. Then multi-
 w multiplying 200 by 200 we have 40000, for the visible Parts of a square Inch;
 * Plate 3. which are contain'd 50 Times in a Grain of Gold; and therefore by this
 Fig. 8. Means it becomes divided into 2 000 000 of visible Parts.

If we consider the gilding of Silver, we shall find Gold in that case to contain visible Parts, even after it has been divided above ten times more; for eight Grains of Gold will gild a whole Ounce of Silver, which is afterwards drawn into a Wire 13000 Foot long; therefore one Grain gilds a Wire 1625 Foot long; and as every Foot (by what we have said above) must contain 2400 visible Parts, the whole Length of the Wire contains 3 900 000 little Cylinders, which being turn'd into Cubes, will each of them have six visible Sides; and consequently by this last Operation, one Grain of Gold, instead of being divided into Two Millions of visible Parts, will be divided into 23 400 000, which is almost twelve times as many. That one may reasonably take Cubes for the little Cylinders, appears, when we consider further, that all this Wire is beaten flat in order to wrap it round Silk for making Gold Lace; and that even after flattening, the best Microscope cannot discover the Silver through the gilding. This shews, that in this thin Skin, several Parts of the Gold still lie upon one another; though the Thickness of it (as Dr. Halley

has shewn, *Philos. Transact.* Numb. 194.) is but $\frac{1}{12450}$ Part of an Inch, or $622 \frac{1}{2}$ times less than the 200th Part of an Inch that we have taken, as the least visible Part of an Inch in Length.

Mr. Boyle, in his Book of *The Nature and Subtlety of Effluvia*, mentions, that one Grain of Copper dissolv'd in Spirit of *Sal Armoniac*, will give a strong blue Tincture to 105,157 Cubic Inches, or near two Quarts of Water. Now supposing no less a Cube of this ting'd Water to be visible, than such a one whose Side is equal to the 100th Part of an Inch (which is making eight times more Allowance than we did in respect of the Gold) it will appear upon Computation, that a Grain of Sand so small, that a Million of them may be contain'd in a Cubic Inch, does contain two Millions an Hundred and eleven Thousand and four Hundred (or 2 111 400) such Parts as the single Grain of Copper is actually divided into.

The same Gentleman, having expos'd to the open Air a certain Quantity of *Affa Fetida*, found it diminish'd in Weight but the eighth Part of a Grain in six Days. Now if we suppose, that during all that time a Man could smell the *Affa Fetida* at the Distance of five Feet, it will appear, that the Particles, into which the *Affa Fetida* is divided, cannot exceed in

Bigness the $\frac{1}{26\ 250\ 000\ 000\ 000\ 000}$ Part of an Inch.

The late Mr. *Lewenhoek*, that ingenious Searcher into Nature, gives us an Account, that in the Milt of one Cod-Fish there were more little Animals, than there are Inhabitants upon the Face of the Earth. Now by only knowing the

the Focal Length of the *Lens*, or Glass of the Microscope, we can by the Rules Annotat. of Optics find the bigness of those *Animalcula*, which cannot be so big as the Lett. I.

$\frac{1}{36\ 000\ 000\ 000\ 000}$ Part of a Cubic Inch: And, therefore, several Thousands of them might stand upon a Needle's Point. And it appears also, that if we compare these to a Whale; they will be much less in Proportion, than a Whale is, when compared with the whole Globe of the Earth. As every Animal is an organiz'd Body; how fine, delicate and subtil must be the Parts that make up one of these Animals? How small must be its Heart? How inconceivably little its Veins and Arteries? And much less must be the Globules of that Fluid which serves it for Blood, and which are still carried along in a finer Fluid.—It is worth while to consider the Smallness of those Globules; which we may do by making the following Allowance; namely, That the Particles of Blood of those Animals are as much less than their Bodies, as the globular Particles of the human Blood are less than a Man's Body.

A Man's Body is to that of an *Animalculum*, as 17 to $\frac{3}{100\ 000}$, and the Diameter of the Globules of a Man's Blood are not bigger than $\frac{1}{79\ 200}$ Part of an Inch †, (because *Lewenhoek* found the Diameter of the Vessels through which they run to be no bigger) therefore as 17 is to $\frac{3}{100\ 000}$, so is $\frac{1}{79\ 200}$ to $\frac{3}{134\ 640\ 000\ 000}$ or in Decimals $\frac{22}{1\ 000\ 000\ 000\ 000}$; therefore the Globules of Blood in these little Creatures cannot be so big as the Cube of this Number or $\frac{8}{1\ 000\ 000\ 000\ 000\ 000\ 000\ 000\ 000\ 000\ 000\ 000\ 000\ 000\ 000}$ Parts of an Inch.

But since these Numbers, express'd in Figures, do not immediately give an Idea of the Smallness of these Globules; Dr. Keil (in whose *Fifth Lecture* one may find at full Length the Demonstrations, that prove our two last Paragraphs) has shewn, that the smallest visible Grain of Sand would contain more of these Globules, than ten Thousand two Hundred and Fifty-six of the highest Mountains in the World would contain Grains of Sand.

What we have said hitherto, shews into how many small Parts Bodies are actually divided; but there are Particles of Matter, so much smaller than the Globules abovementioned, that those Globules compared with them will not only be as Mountains, but as vast Earths. I mean the Particles

† The learned and ingenious Dr: J. Jurin, Secr. of the Royal Society, has not long since found Globules of the human Blood to be bigger than what is here mentioned; and having communicated his Observations to Mr. Lewenhoek, was by him confirm'd in his Assertion; but I did not think proper to alter this Calculation, since I shall speak fully upon this Subject in another Place of this Course.

Annotat. of Light, whose inconceivable Fineness puzzles our Conception. How
 Lect. I. amazingly little must those Particles be which flow from such a Candle,
 whose Light may be seen at two Miles Distance? Since every Instant of
 Time, Particles must be darted out to fill a Sphere of four Miles in Diame-
 ter, so that a Pin's Head cannot be placed any where in that Sphere with-
 out receiving some Particles of Light. Dr. *Newenham* shews, that the
 fourteenth Part of a Grain of Wax or Tallow, (that is consumed in one
 Second of Time in a Candle of six to the Pound,) produces a greater Number
 of Particles of Light, than a Thousand Times a Thousand Millions of
 Earths, (equal to our Earth) would be able to contain Grains of Sand. See
Religious Philosopher, Vol. 3. *Contemplation* 25. *Seet*, 15, 16, 17.

7. [7—*There must be a great deal of Vacuity, &c.*] The different spe-
 cific Gravity of Bodies plainly proves this Assertion, as will be more ful-
 ly shewn in the second Lecture. And in Fluids this is evident from their
 different Resistance, which we have already shewn to be proportionable to
 the Quantity of Matter in Bodies.

8. [—*All Spaces are not equally full of Matter.*] If there was any
 such thing as a *Subtile Matter*, that wholly fill'd up the Vacuities of Bodies,
 and the whole *Æthereal Space* in which the Planets move; its Resistance
 would be such as far to exceed the Resistance of Quicksilver. In such a
Medium as that, even a perfectly solid Globe must lose half its Motion,
 before it could move thrice the length of its own Diameter; and such
 Globes, as the Planets are, would be stopt much sooner; wherefore 'tis ab-
 solutely necessary for continuing the Motions of the Planets and Comets,
 that the Places, they move in, be almost entirely void of Matter. That
 they are so, appears from the swift Motion of the Tail of a Comet,
 that does not appear to meet with any sensible Resistance in the *Medium*
 in which it moves, though it is expanded so wide, and made up of so thin
 a Vapour.

9. [9—*Much more Vacuum than Matter, &c.*] That there is more
 Vacuity in Bodies than Matter, may be clearly deduced from the Proper-
 ties of transparent Bodies; for the Rays of Light are spread every way in
 right Lines through Water, Glass, or a Diamond, with no more Difficulty
 (nay, with more Swiftness) than they are carried through the Air, whatsoever
 Side of the transparent Body be exposed to the Light; therefore there is
 always a rectilinear Passage for the Light, from the least assignable Part of
 the transparent Body to any other Part of it: And this could never happen,
 unless the Quantity of Matter in such a Body was extremely small when
 compar'd with its Bulk. Perhaps in a Diamond the solid Matter, compar'd
 to its Bulk, bears a less Proportion than the Diamond does to the whole
 Globe of the Earth: Which will not appear impossible, to those that consi-
 der what has been said before upon this Subject. Now since Gold is not
 above

above fix times denser than a Diamond, how much more Vacuity must be in it than Matter? This shews the Reason why the *Effluvia* of the Loadstone pass through Gold as easily, as through the Air; for if a Plate of Gold, or any other Metal (except Iron) be interposed between a Loadstone and a touch'd Needle, which is drawn out of its Position by the Virtue of the Stone, the Needle will in no wise be less affected than before. Nay those *Effluvia* may for a whole Day pass through the Brain, a Body so tender and of so delicate a Contexture, without affecting the Nerves with any Sensation, or disturbing the least Thought.

The Vapour of the *Aurora Borealis* (which some imagine to consist in a great Measure of the Magnetic *Effluvia* of the Earth) does freely pass through Houses and Trees, and dart through the Bodies of Animals without being felt; as appears from Observations made upon that *Phaenomenon*, when viewed from several Places at once.

10. [15—*Any Weight may hang, &c.* The same Experiment may be made, by Means of a spiral Spring in one of the Balls within a Barrel on which a String is wound up. For the end of the String being fasten'd to another Ball equal to (or double, or triple) the Ball that has the Spring; upon pulling the Balls asunder, they will come together again with Velocities reciprocally proportional to their Masses. See Plate 3. Fig. 9. Where A *represents the Section of an hollow Brass Ball, with a spiral Spring S and Barrel within it, contrived in such Manner as to pull back into the Ball the whole String AB, when it has been pull'd out by the End B. The Ball B is solid and of Ivory, but of the same Weight as the hollow Ball A. Now if the Ball B be pulled from A to the Distance AB, upon letting go both the Balls at once, they will meet at C, the middle Point between them; but if B be a Ball twice as heavy as A, upon letting both go they will meet at D, DB being but half of the Distance AD. Plate 3. Fig. 9.

11. [17—*For Gravity being a Virtue diffus'd, &c.*] When we compare Gravity to Light and Heat, we would not be understood to derive their Effects from the same Cause; or to assert, that all Sorts of Attractions in Bodies have the same Laws, since the Attraction of Cohesion*, and the Attraction of the Loadstone †, do not act in that Manner: But here we only speak of the Attraction of Gravity, whereby the Bodies about us are driven towards the Earth, and the Earth and Planets are driven towards the Sun; which Attraction is called also a *Centripetal Force*: And whatever be its Cause, its Laws abovementioned have been discover'd by Observations and Experiments, which have always concurred to confirm the Theory of Gravity. *L.I.No.19, 20, &c. †L.I.No.29.

In order to have a clear Notion of the Effects of Gravity, or the Centripetal Force, we must consider it in three Respects: Either in respect to the Quantity of Force in the Central Body that attracts others, (or towards which Circumambient Bodies tend) which is call'd the *Absolute Force*; or in respect to

Annotat. to the Velocity with which other Bodies move towards the Central one, which is call'd the *Accelerating Force*; or in respect to the Quantity of Motion in the said Bodies, when compar'd with each other, which is proportional to the Obstacle that they are able to remove, and is called the *Moving Force*.

The *Absolute Force* is proportionable to the Efficacy of the Cause that spreads its Virtue from a Center round about. So if the Earth had twice the Quantity of Matter that it has now, (whether it was twice as big, or only twice as dense) it would have twice the Absolute Force. Thus the Moon has near forty times less Absolute Force than the Earth, because it has almost forty times less Matter. Its Bulk indeed, (which is as the Cube of the Diameter) is almost fifty times less than that of the Earth; but then it is denser in the Proportion of 21 to 17. See Sir *Isaac Newton's Principia*, Book. 3. Prop. 37. Corol. 3.

* Lect. 1. The *Accelerating Force* is expressed by that Velocity generated in a given Time, with which Bodies (considered as Physical Points) move towards the central Body attracting them by its Absolute Force. And this Accelerating Force is greater or less according to the Distance of the Center of the Force, in the reciprocal duplicate Proportion abovementioned.* Thus No. 17. is the Gravity, that makes Bodies tend towards the Center of the Earth, greater in Vallies than on the Tops of high Mountains; greater at the Poles than at the *Æquator*, which is seventeen Miles higher; and greater at the *Æquator* than at greater Distances from the Globe of the Earth: For the same Body, which, near the Surface of the Earth, falls sixteen Foot in the first Second of its Fall, would fall but four Foot in the same time, if it began to fall at the Height of 4000 Miles from the Surface of the Earth, or two Semidiameters Distance from its Center. At equal Distances the † Lect. 1. Accelerating Force is the same every where: because all Bodies large or No. 8. Ex- small, heavier or lighter, (abstracting from the Resistance of the Air,) are per. 1. equally accelerated in their Fall.†.

The *Moving Force* is proportional to the Quantity of Motion, which the Absolute Force of the central Body generates in a given Time, in the Bodies that it acts upon. Though in respect of the Earth, we consider the biggest Bodies that are attracted by it (and even the Moon it self) as Physical Points; yet when we compare the Bodies with each other, we must have a Regard to their respective Quantities of Matter; for Bodies that have the same Accelerating Force, or move with the same Velocity, have their Quantities of Motion greater or less, as they have more or less Matter, or as they are more or less heavy; because the Moving Force of a Body is made up of the Sum of the Actions of the Accelerating Force of all its Parts, and consequently it is found by multiplying the Quantity of Matter into the Accelerating Force; as the Quantity of Motion in Bodies is found, by multiplying their Matter, or their Mass, into their Velocity.

Hence

Hence near the Surface of the Earth, where the *Gravitating* or *Accelerating* Force on all Bodies is the same, the *Moving Force* or Weight is as the Body: But if we ascend to Places where the *Accelerating Gravity* is less, the Weight will also be diminish'd and become as the Mass of the Body multiplied by the *Accelerating Force*. Thus if a Weight of one Pound and a Weight of four Pounds begin to fall near the Surface of the Earth, their *Moving Forces* will be as 4 and 1; for if we take 16 for the *Accelerating Force*, equal in (and common to) both, 16 times 4, (or 64) will be just 4 times as much as 16, (or 16 times 1.) But if the four Pound Weight was remov'd to the Height of 4000 Miles, or two Semidiameters from the Center of the Earth, its *Moving Force* then would be just equal to that of one Pound at the Surface of the Earth; because the *Accelerating Force* being 4 times less at twice the Distance from the Center, 4 times 4 or 16 would then express the *Moving Force* of the heavier Body.

If the one Pound Weight was plac'd at the Distance of two Semidiameters, its *Moving Force* would be 16 times less than that of the 4 Pound Weight at the Surface of the Earth.

A clear Idea of these three Forces will shew the Reason of some *Phænomena*, which would otherwise be hard to explain: As for Example, we have already said, that the Earth has near forty times more Matter than the Moon; and yet Bodies on the Surface of the Moon weigh but three times less than they do on the Surface of the Earth, though the Moon's *Absolute Force* be forty times less. But to shew that this is a necessary Consequence of what has been said, let us examine this Matter by Numbers.

Let *TAB represent the Earth, and L the Moon, AB a Diameter of the Earth, and bd a Diameter of the Moon (which are to one another as the Numbers 365 and 100) and let the Lines CE and ce be each equal to a Diameter of the Earth. Now if we suppose a Body plac'd at E, whose Weight or Gravity towards the Earth is there equal to 9,827 Pounds; the same Body plac'd at e, just as far from the Center of the Moon, as it was before from the Center of the Earth, will weigh towards the Moon L but 0,25 Pound, or a Quarter of a Pound; because the Masses or Quantities of Matter in those two Bodies, and consequently their *Absolute Forces*, are to one another as these Numbers, or as 39,721 to 1, which are in the same Proportion. *Newtoni Prin. Lib. 3. Prop. 37. Corol. 4.*

Then if the Body abovemention'd be plac'd at A, distant from the Center of the Earth but a Semidiameter, it will † weigh towards the Earth four times more than it did at E or 39,721 lib. and at a it will weigh towards the Moon 1 lib. or 4 times more than it did before, for the same Reason. Now if the Moon, without any new Addition of Matter, was so expanded or rarified, as to fill up the Sphere ma, which is equal to the Globe of the Earth, then the Point a would be on the Surface of the Moon, as the Point A is on the Surface of the Earth; and in that Case the Weight of Bodies on the Surface of the Earth would be to the Weight of Bodies on the Surface of the Moon, precisely as the Quantity of Matter in the Earth, to the Quantity of Matter in the Moon; and consequently as

F

their

Annotat.
Lect. I.

* Plate 3.
Fig. 10.

† L. I. No. 17.

Annotat.
Lect. I.



*L.I.No.17.

their *Absolute Forces*. But as the Moon is less in Diameter than the Earth, when the Body that weighed 1 *lib.* at *a* comes to be placed on its Surface at *d*, it will be nearer to the Center of the Moon than it was, in the Proportion of 182,5 to 50, or 365 to 100, and therefore * it will then weigh 13,3225 *lib.*; for as the Square of *cd* ($100 \times 100 = 10000$) is to the Square of *ca* ($365 \times 365 = 133225$) so is 1, or the Weight of the Body placed at *a*, to 13,3225, the Weight of the same Body upon the Surface of the Moon; which Number being very near the third Part of 39,371, shews, that Bodies on the Surface of the Moon weigh about a third Part of what they do on the Surface of the Earth. Which was to be demonstrated.

Hence it follows, That Bodies weigh more on the Surface of the small Planets, in Proportion to their Quantity of Matter, than on the Surface of the larger. Thus upon the vast Globe of *Jupiter*, whose Quantity of Matter or *Absolute Force* is 220 times greater than that of the Earth, Bodies weigh but twice as much as they would do upon the Surface of the Earth. And upon the immense Body of the Sun, whose Quantity of Matter is 227512 times more than that of the Earth, Bodies weigh only 24,4 times more than they do upon the Surface of the Earth.

Hence follows also, that in respect of any Planet or central Body, as for Example, the Earth, the Weight of Bodies gravitating towards its Center is greater on its Surface, than at any Distance without it, or any where within it, though nearer the Center. For if the same Body which at *A* weighed 39,371 *lib.* should be brought to *D* a Point within the Earth as near to *C* the Center of the Earth *TAB*, as *d* is to *c* the Center of the Moon *L*, it would not encrease its Weight towards *C* 13,3225 times, as it does when remov'd from *a* to *d*; but it would diminish it in the Proportion of 365 to 100, because that Part of the Earth towards the Surface between *D* and *A* attracts back the Body towards *A*.

The just Proportion of this Decrease of Gravity is determined by Sir *Isaac Newton*, *Lib. 3. Prop. 9.* to be always as the Distance from the Center, going from the Surface downwards: And as the Principles from which it is deduc'd are very evident, I shall repeat them here.

† Plate 3. If there be a concave spherical Surface, whose Particles attract accord-
Fig. 11. ing to the Laws of Gravity abovementioned, as *IHKL*,† any little Body within it will remain at rest where ever it is plac'd, the Attractions round about destroying one another. This is evident, if the Body be plac'd at *C* in the Center. And if the Body be plac'd at *P* as near again to *H I* as to *KL*, the same thing will follow; for let the Lines *I K* and *K L*, be drawn, and it will be evident that the spherical Segment between *K* and *L* will be four times greater than the Segment between *I* and *H*, because *K L* is equal to twice *H I*; therefore there are four times more attracting Particles at *KL* than at *H I*, but *H I* being twice nearer to the Body *P* attracts it four times as much, which makes amends for the fewer Particles contain'd in the lesser Segment; for the Product of the *Absolute Force* of *H I* (1) multiplied into its *Accelerating Force* (which here is 4) is equal

equal to the Product of the *Absolute Force* of *KL* (4) multiplied into its Annotat. *Accelerating Force*, which is but 1. *This will hold good in respect of any other* Lect. I.
Part of the Surface or Position of the Corpuscle: And if the Corpuscle be put
into Motion, it will go on uniformly in the concave Sphere, as if it was not
attracted at all. If instead of a Surface there was a Shell of any Thick-
ness as ABHIKL, every thing else would remain as before asserted,* * Plate 3.
provided that Shell was every where of the same Thickness and Density. Fig. 12.

If the Hollow *HIKL* be fill'd with a solid Sphere and the Corpuscle be plac'd at *P*, it will be attracted towards the Center *C* only by the Force of the inner Sphere *HIKL*; for the attractive Forces of the several Parts of the Shell destroy one another, as has been shewn before. If the Corpuscle be remov'd to *Q*, it will be only attracted by the Sphere *QR*. Let us suppose the *Absolute Force* of the whole Sphere *AB* = 64; the Lines *BC*, *PC*, and *QC*, as 4, 2, and 1; and the *Accelerating Force* of the Corpuscle at *B* to be = 1; its *Moving Force* will of consequence be 64. Now if the Corpuscle be brought to *P* (as near again to the Center of the Forces) its *Accelerating Force* will be 4, but then that Number must only be multiplied by the *Absolute Force* of the Sphere *HIKL*, which being twice less in Diameter than the Sphere *AB* is eight times less in Solidity, and therefore its *Absolute Force* will be but 8; which multiplied into the *Accelerating Force* 4, gives only 32 for the *Moving Force* of the Corpuscle at *P*. If the Corpuscle be brought to *Q*, four times nearer to *C*, its *Accelerating Force* will be 16; which being multiplied by 1, the *Absolute Force* of the Sphere *Q* (which is four times less in Diameter than the Sphere *AB*) will give 16 for the *Moving Force* of the Particle at *Q*. Now since the *Moving Forces*, whereby the Corpuscle or Particle at *B*, *P*, and *Q* gravitates towards the Center *C*, are as 64, 32, and 16, and the Distances from the Center are as 4, 2, and 1; it follows, *That going from the Surface of a Planet downwards, the Gravity decreases directly as the Distance from the Center.* Which was to be demonstrated.

To apply this to what we said before; if we conceive a Globe *CD* within the Earth *T*† equal to, and equally dense with the Moon: The † Plate 3.
 Body which at *A* weighed 39,721 *lib.* will at *D* weigh just 13,3225 Pounds, Fig. 10.
 as it would do upon the Moon's Surface.

12. [18—If a Cube of an Inch, &c.] As in the Bodies shin'd upon we only consider the Surfaces enlighten'd; the Experiment may be made with similar Pieces of Pastboard exposed to the Light: As for Example, a Circle, a Square, a Pentagon, or any other Polygon of an Inch Diameter a Foot distant from the Candle, will receive the same Quantity of Light that falls upon a Circle, Square, or Pentagon, &c. of two Inches Diameter at two Foot from the Candle, or the like Figures of three Inches Diameter at three Foot from the Candle; where we are to observe, that the Strength of the Light is diminish'd in the same Proportion, as the Area of the Figures is encreased; That is, here at the Distances of 1, 2, and 3 Feet, the

Annotat. Strength of the Light will be as 9, 4, 1; which we call a reciprocal Duplicate Proportion of the Distances.
 Lect. I.

13. [19—*When the Wood is more solid, &c.*] If we make use of a Glue whose Parts are finer in Proportion to those of the Wood to be joined together, it will hold as fast as in softer Wood; as is experienced, when hard Wood is glued with Fish-Glue dissolv'd in Spirit of Wine.

14. [19—*The Figure of any Portion of a Fluid, &c.*] When two small Drops come to touch one another, they first become oval, and then immediately spherical. Now when the Drop is of an oval Figure, the Pressure of an external Fluid acting upon it from all Parts cannot alter the Figure of the Drop so as to bring it to be round; neither can it make it flatter, as some have imagin'd, who have asserted,—That if there was no Attraction of the Particles of the Liquor, but only the Pressure of an external Fluid, an oval Drop would be more press'd against the Ends of the short Diameters, than against the Ends of the long one, or of the Drop, which would cause it to be lengthen'd. But I shall not make use of this Argument, because it is a Fallacy; as will appear to such as will be at the Pains to consult Sir *Is. Newton's Principia Lib. 2. Prop. 19.* where it is demonstrated, that if any Portion of a Fluid be press'd by the same or any other homogeneous Fluid acting from all Sides, that Portion will not have its Figure alter'd by that Pressure.

* Plate 2.
 Fig. 13.

15. [19—*Whereas from a mutual Attraction of the little Globules, &c.*] If two similar and equal Globules A and B* attract one another and touch at C, they will remain at rest, as they would if they touch'd at E or at D, or any other Point in the Circumference of either of them, because the Contact would still be the same: And therefore a very small Force will remove B from its Contact at C to E, or any other Point of the Circumference (or rather of the Surface of the Sphere) CED; because in going round one Globule it still continues to touch it as much as it did before.

† Plate 3.
 Fig. 14.

* Fig. 15.

If there be three Globules A, B, F, †touching one another in such manner, that their Centers are in the Line *ab*, they will remain in that Position; but if any one of them be mov'd out of that Position as F*, it will not remain at F, but move on toward C where the other Globule A will meet it, so that each of the three Globules will touch in two Points, being as many Points as three Spheres can touch one another in. See Fig. 16.

† Fig. 17.

If there be four Globules in the Position *g A, B, f*, they will for the same Reasons come into the Position *A, F, B, G*, supposing their Centers all in the same Plane; but if any one of them, as F, be lifted up, it will not rest till it comes to *c*, and then the Spherules will touch each of them in three Points. Hence it is, that when two Drops of Water or any other Liquor coming to touch one another, make up the Spheroid *acdbef*, they

they will not preserve that Figure, but run into the Sphere *c g d e b f*, Annotat. that there may be the greatest Contact possible between the Globules of *Left. I.* which the Sphere or new Drop is made up.

Now because it may be thought unmathematical to draw Consequences from the Figure of the Parts of Fluids, without having first demonstrated that they are spherical; I shall here subjoin another Proof of the Roundness of the Drops of Liquors, without having any regard to the Figure of the Particles of which they are compounded.

Let * *ABDE* be a Portion of an homogeneous Fluid whose Parts at- * Plate 3. tract one another, and whose Figure is not spherical. If in such a Fluid Fig. 19. we suppose a Syphon as *ACE* (or which is the same thing, if all the Fluid should be frozen, except the Canal *ACE*) whose Legs *AC* and *CE* are unequal, and meet at *C* the Center of the Fluid, towards which there is the greatest Attraction; the Fluid will run out at *A* in the Leg *AC*, till it be come down as far as *g* in the Leg *CE*, supposing *Cg* equal to *AC*. But if the Leg *AC* be lengthen'd as far as *c*, then the Fluid will only come down as far as *e* in the Leg *CE*, and at the same Time rise up to *a* in the Leg *Ca*, *Ca* being equal to *Ce*.

If such another Canal or Syphon be supposed at *BCD*, the Fluid in it will come down from *D* to *d* and rise from *B* to *b*. And since such Syphons may be supposed all over the Fluid *ABDE*, that Fluid by the Attraction of its Parts must needs be reduc'd to the spherical Figure *abde*. Which was to be demonstrated.

16. [20—*Having moistened, &c. two Glass-Planes, &c.*] This Experiment of the Drop of the Oil of Oranges, having been made by the late Mr. *Hauksbee*, in the Manner that Sir *Isaac Newton* relates it in the third Part of his *Optics* (*Query the last, towards the Middle*) that incomparable Philosopher has calculated the Force of the Attraction, and says; "That where the Oil of Oranges, between the two Glass-Planes, is of the Thickness of three Eighths of the ten Hundred Thousandth Part of an Inch, the Attraction (collected by the Rule given in the Table of the Second Part of the Second Book) seems to be so strong, as within a Circle of an Inch in Diameter, to suffice to hold up a Weight equal to that of a Cylinder of Water of an Inch in Diameter, and two or three Furlongs in length.

17. [20—*The Attraction must encrease continually as the Planes come nearer.*] There are six Properties to be observ'd in the Attraction of these Glass-Planes.

Let † the Point *O* (the Center of the Drop) be at equal Distances from † Plate 3. the Glass Planes *Qm*, *Qn*; and let the Radius *Ob* express the greatest Di- Fig. 20. stance at which the Places of Glass *Qm*, *Qn* can have any Effect upon the Point *O*. It is plain,

First,

Annotat.

Lect. I.




First, That at its greatest Distance from the Point Q , there will be no greater a Part of the Plate of Glass, than a Circle whose Diameter is cd within the Sphere of Attraction, or whose Parts are able to attract the Point O , all the other Particles being at too great a Distance. *Secondly*, The Force of the Attraction of the Points of the Circle of Glass which are nearest to the Point Q is also the greatest. *Thirdly*, The Sum of the attracting Particles plac'd towards Q , and which are contain'd in the Segment of the Circle of Glass, whose Arc has sc for its versè Sine, is greater than that of the attracting Particles contain'd in the opposite Segment, in the same Proportion as the first Segment exceeds the other, because of the Angle made by the Planes at Q . *Fourthly*, The Direction of the Attraction of the Points, that are at the same Distance from O , making all along the Line QR an Angle at O , more acute towards Q than towards R , the Point O will advance towards Q with an accelerated Velocity. *Fifthly*, As the said Point O advances towards Q , the Diameter of the Circle of Glass whose Particles can attract the Point O will be encreased, (as gd is greater than cd) and consequently the Circle also; which will cause the Drop of Oil to spread more and more upon the Planes of Glass, between which it was plac'd. *Sixthly*, The Drop or Point O will advance towards Q , with a Velocity always accelerated with greater Forces; because the Angle gOb becomes always more acute in respect to the Angle dOq in proportion as the Chord gd becomes greater than the Chord cd , which makes the Base gb always less in respect of the Base dq , whilst the Sides of the two Triangles always remain equal; and consequently the Angle gOb , is always less in proportion to the Angle dOg ; which continually encreases the Force of Attraction towards Q by the following Demonstration.

* Plate 3.
Fig. 21.

Let the * Angle abc be divided into equal Parts by the right Line bu . and about the Centers t, u , taken at pleasure upon that Line, draw the equal Circles $gbki, acml$; I say that if through the Points of Interfection, the Chords ac, lm ; gb, ik , be drawn; the Ratio of lm to ac will be greater than that of ik to gb . From the Point b draw the Tangents bfp, brq ; and from the Point u for a Center draw between the said Tangents the Circle $eond$, and join ed, no .

The Segments de, gb , are similar, as well as no, ik , as also are the Arcs ope, kfb , which makes $ik: no :: gb: de$, and $ik: gb :: no: de$. Now lm is greater than no , and on the contrary ac is less than de ; therefore the Ratio of lm to ac is greater than that of ik to gb : Which proves the sixth Property of the two Glass Planes that touch at one End, and are open'd to a small Angle at the other.

18. [23—*The Reason of these Phenomena, &c.*] Dr. James Jurin, Secretary of the Royal Society, has made a great many curious Experiments of this kind, of which he gives an Account in *Phil. Transf.* Numb. 355. where he shews in what manner the Attraction of Cohesion operates to raise and sustain Water in small Tubes, and such Spaces between solid Bodies as are analogous to small Tubes,

19. [23—*Quicksilver attracts Quicksilver more than Glass attracts it, &c.*] Annotat. Some have been apt to imagine that Quicksilver and Glass repel one another, because it does not in these and several other Experiments appear to stick to Glass; but that it is really attracted by Glass (though so much less than by it self, as to make it seem to be repell'd) will be shewn by giving an Account here of some more Experiments relating to the Attraction of Cohesion, which the abovementioned ingenious and learned Gentleman made with Glass and Quicksilver. 

EXPERIMENT 1.

Quicksilver is attracted by Glass.

If a small Globule of Quicksilver be laid upon a clean Paper, and be touch'd with a Piece of clean Glass; upon drawing the Glass gently away, the Quicksilver will adhere to it, and be drawn away with it. And if the Glass be lifted up from the Paper, the Quicksilver will be taken up by it, in the same manner as a Piece of Iron is drawn by the Loadstone, and will stick to the Glass by a plane Surface of a considerable Breadth, in Proportion to the Bulk of the Drop, as manifestly appears by an ordinary Microscope. Then if the Glass be held a little obliquely, the Drop of Mercury will roll slowly upon its Axis along the under Side of the Glass till it comes to the End, when it will be suspended as before.

EXPERIMENT 2.

If a pretty large Drop of Mercury be laid upon a Paper, and two Pieces of Glass held edge-wise be made to touch it, one on each side; upon drawing the Glasses gently from each other, the Drop of Mercury will adhere to them both, and will visibly be drawn from a globular to an oval Shape; the longer Axis passing through the middle of those Surfaces, in which the the Drop touches the Glasses.

The Particles of Quicksilver are more attracted by one another than by Glass. For the Proof of this see *Left I. N^o. 24, 25, 26,* and those other Experiments of Dr. Jurin.

EXPERIMENT 3. Plate 3. Fig. 22.

Quicksilver been pour'd into the inverted Syphon ACB, one of whose Legs A C is narrower than the other CB; the Height CE, at which the Mercury stands in the wider Leg CB, is greater than the Height CD, at which it stands in the narrower Leg CA. On the contrary, Water stands higher in the narrower Leg than in the wider.

Ex-

EXPERIMENT 4. *Pl. 3. Fig. 23.*


Annotat.
 Lett. I.

ABCD represents a rectangular Plane of a Glass which makes one side of a wooden Box. On the inside of this is another Glass Plane of the same size, which, at the End AC, is press'd close to the former, and opens to a small Angle at the opposite End BD. When Mercury is pour'd into this Box to any Height as CE, it insinuates itself between the two Glass Planes; and rising to different Heights between the Glasses, where the opening is greater or less, it forms the common Hyperbola CGF; one of whose Asymptotes EF is the Line on which the Surface of the Mercury in the Box, touches the inner Glass; the other is the Line AC, in which the Planes are joyn'd.

EXPERIMENT 5. *Pl. 3. Fig. 24.*

AB is a perpendicular Section thro' two glass Planes joyn'd at A, and open'd to a small Angle at B; C represents a pretty large Drop of Mercury, the larger the better, which (being made to descend as far as C, by holding the Planes in an erect Posture, with the End A downwards) retires from the contact of the Planes to D, upon inclining the Planes towards an horizontal Situation; and the Distance CD becomes greater or less, as the Planes are more or less inclin'd towards the Horizon.

20. [30—*As the Cube and a quarter of the increas'd Distance—the Magnetical Attraction, &c.*] That excellent Philosopher Dr. Brook Taylor made some Experiments with a touch'd Needle and a large Load-stone that is kept in the Repository of the Royal Society, which, when made at some Distance from the Stone, agreed very well with this Assertion; but near the Stone, the magnetick Virtue did not seem to act according to those Laws, which might be owing to this, viz. That that Stone rather seems to be an aggregate of Load-stones joyn'd together by a petreous Substance not magnetical. For, since that Time, by some Experiments made upon it, I found that it had 15 Poles (if I may use that Expression) or Points where the Attraction was stronger than any where else; which Experiments, and some others made upon weak Load-stones, made me imagine that every Load-stone had several Poles, or Points of Virtue on the North side, and several on the South side; whose Virtues being collected by the Iron wherewith the Stone is arm'd, made an arm'd Load-stone sustain much more Iron or Steel than the same unarm'd. But by Experiments made since, with some good Load-stones, especially, with a Stone of about six Ounces, belonging to the Right Honourable the Lord Paizley (one of the best in the World) I found that a good and homogeneous Load-stone has but two Poles.

21. [31.—*A repellent Power in Bodies, &c.*] See s'Gravesande *Introduction*, Annotat. Part I. from N^o. 40 to 44. When Light is reflected from a polish'd specular Surface of Glass, Crystal, or Metal; the Particles of Light do not strike upon the solid Parts, and so rebound from them; but are repelled from the Surface at a small Distance before they touch it, by a Power extended all over the said polish'd Surface. See Sir Isaac Newton's *Optics. Book II. Part III. Prop. 8.* Lect. I. 


The Rays of Light are also repell'd by the Edges of Bodies as they pass near them, so as to make their Shadows, in some Cases, larger than they wou'd otherwise be. See the same Author, *Book III. Part I.* where he likewise proves this repulsive Force from other *Phænomena*.

22. [37.—*The same Part of the Tube will not snap or give Light twice together, in the same Place, without a new Friction.*] By causing the Tube to snap at the Approach of the Fingers, or any other solid Body near it, the Electricity of it in that Place is also destroy'd; from which we may easily solve a *Phænomenon* mention'd by Dr. s'Gravesande in his *Introduction. Vol. II. N^o. 554. These are his Words:*

“ There is one thing remarkable and very hard to explain in this Experiment, concerning the Direction of Attrition; when you rub the Tube, one End is held in one Hand, whilst it is rubb'd with the other; which, if it be done from the Hand that holds towards the other End of the Tube, the Effect will not be sensible; but if you rub from the free End of the Tube towards the End held in the Hand, the contrary will happen. And this happens indifferently whether you hold the open or shut End of the Tube in your Hand.

To explain this, let us examine the Experiment by help of the 10th Figure of Plate 2. A is the Right Hand that holds the Tube, and B the Left Hand that rubs the Tube, which, after moving up and down several times, makes an end of the Attrition by moving in the Direction CBA, the last Stroke: Then the Tube brought near light Bodies (*Plate 2. Fig. 11.*) gives Motion to them. If the last Stroke in rubbing be made by the Motion of the Hand, from A to C, and the Hand B that went up to C quits the Tube in the Direction CD, without coming near it again, or letting the Coat-sleeve come near it, the Tube will act upon light Bodies with the same Vigour as before. But if the Left Hand, after it is come off at C, is brought down again carelessly, in the Direction CE, in a Line Parallel to the Tube, and pretty near it, either the Hand or the Coat-sleeve coming down too near, causes the Tube to snap (which is not heard without Attention) and so destroys the Virtue excited by Attrition, in the whole Length of the Tube; as the Hand A (*Plate 2. Fig. 13.*) does in the Place A where it causes a Snapping, which, as I said before, will not happen twice in the same Place, without a new Friction.

23. [41.—*A Glass Globe whirl'd, &c.*] I cannot forbear mentioning here a very surprizing *Phænomenon* in one of Mr. Hauksbee's Experiments. He
G
coated

Annotat. coated with Sealing-wax the Inside of one Hemisphere of one of his Glafs
 Lect. I. Globes to such a Thickness, as to render it perfectly opaque: Yet when this
 Glafs was exhausted of its Air, and whirl'd round; where the Hand was apply'd on the Outside to give an Attrition to the Globe, the Wax became as transparent as the Glafs it self, the rest of the Globe that had Wax remaining opaque, where it did not touch the Hand, though the Moment before it had been transparent as it pass'd under the Hand.

A View of the 25th Fig. of Pl. 3. will represent the Thing fully.

The Hemisphere ACB of the Glafs Globe GACB is made opaque, by being lin'd with Wax on the Inside, whilst AGB is clear Glafs. The Glafs is exhausted of its Air by means of the Cock D; then being set between the Pillars EF, it is whirl'd swiftly round by means of the Wheel K, whose String goes round a Pulley P fix'd to the Brass Socket, whose Shank is the Axis of the Globe. The Screw H draws forward the other Pulley I, to keep the Wheel-string always tight.

When the Hand is apply'd at C, its Inside becomes visible, through the Wax on the concave Side of the lin'd Globe, the rest of the Wax remaining opaque; so that the Hand cannot be seen by an Eye at Q, but by an Eye at O looking at it through the unlin'd Part of the Glafs at G.

24. [42.—*Though the Causes of these Causes are not known, &c.*] When Genealogists, in searching into the Original of Families, are got as far as they can, and have found the first of the Family, so call'd, because they cannot discover who were his Parents: It wou'd be very absurd to say, that because the Father of this First (which for Example we will call *John*) is not known, therefore *John* is not the Father of *Peter*, the Grand-father of *William*, and Great Grand-father of *Nicholas*, &c.—Which was prov'd before, by such Evidence as is proper in that Case. So when it appears by Observation that Gravity is the Cause of the Fall of heavy Bodies, which observe certain Laws in their Motion—That a heavy Body by its Descent moves the Axis of a Wheel, that carries round another by its Teeth, which by the Intermediation of other Wheels and Pinions, carries round a Hand upon a Dial-Plate to measure Time, or for other Uses; it wou'd be very unphilosophical to say—That our Reasoning about the Cause of the Motion of the Hand is false, as being founded upon *occult Qualities*; because we can go no higher than Gravity, whose Cause we don't pretend to know.

LECTURE II.

I. **T**HE *Momentum* or *Quantity of Motion* in Bodies (some-
times call'd simply *Motion*) is that Force with which Bo-
dies change their Place.

I DON'T mean the Stroke, Pressure, Traction, or any other Action which causes this Change of Place in a Body; but the Force which it has all the while it is moving from one Place to another.

2. THIS *moving Force* may always be known by the Effect which it is able to produce; that is, by the Stroke which the moving Body can give, or by the Resistance or Obstacle which it is able to overcome.

3. THIS Quantity of Motion, which is the Measure of the Force, is made up of the Quantity of Matter and the Velocity taken together. That is, when we compare the *Momenta*, moving Forces, or Quantities of Motion in Bodies, we multiply the Mass or Quantity of Matter in each Body by its Velocity *.

* Annot. 1.

4. *Velocity* or *Celerity*, is the Swiftnefs with which a moving Body changes its Place; and may always be known by the Space that the Body goes through in a given Time.

5. THE Quantity of Motion may be encreas'd; either by encreasing the Quantity of Matter, which is mov'd with a determinate Velocity; or keeping the same Quantity of Matter, and encreasing the Velocity; or by encreasing both. — And in the three Cases it is done by applying more Force; for here * *Force* and *Motion* mean * Annot. 2.
the same Thing.

6. THE Motion of any Whole, is the Sum of the Motion of all the Parts; and therefore (as we said before) it becomes doubled in a double Body mov'd with equal Velocity, and quadrupled in a double Body mov'd with a double Velocity.

7. IF a Man with a determinate Force throws from him a Weight of 50 Pounds to the Distance of ten Feet; he must apply twice the

Leſt. II. Force to throw a Weight of 100 Pounds to the ſame Diſtance, or to throw the 50 Pound Weight twice as far ; but if he uſes no more Force than he did before, he will throw the 100 Pound Weight only to the Diſtance of 5 Feet, and then the two Bodies will have the ſame Quantity of Motion ; becauſe 50 multiplied by 10, or 100 multiplied by 5, give the ſame Product, *viz.* 500.

EXPERIMENT I. *Pl. 4. Fig. 1.*

*Pl. 4.
Fig. 1.*

8. ABCDE is an Inſtrument contriv'd for illuſtrating what has been ſaid, and diſtinguiſhing Motion and Velocity, which ſome Authors have confounded. The rabbeted Cheeks BD, CE, are ſo contriv'd that the ſmooth cylindrick Weight K or L (the one of 8, the other of 4 Ounces) may move between them with very little Friction. Let the Spring AB be bent to a certain Degree, by ſlipping the Knot G of the String faſten'd to B upon the Catch of the Iron F ; then lay on the Weight L at B, which upon letting go the Spring, by lifting up the Knot, will be ſhot from B to the Point I, which is 24 Inches from B. If the Cylinder K be ſhot in the ſame Manner, it will go but to H 12 Inches from B.

THAT the Quantity of Motion is the ſame in both Bodies, is evident, becauſe the Spring is equally bent in both Caſes ; and that thoſe Quantities of Motion are made up of the Maſſes multiplied into the Velocities is alſo evident, becauſe $L = 4 \text{ Ounces} \times \text{BI}$ (24) its Velocity gives 96, equal to K (8 Ounces) \times BH (12) its Velocity.

BUT if you would have K driven as far as the Point I, to which L was driven ; the Spring muſt be bent with a double Force, and then K will have double the Motion that it had. N. B. *This Experiment is made uſe of rather to illuſtrate this Matter than to prove it.*

9. HENCE it follows, that any little Body may have as much Motion as a great one, be their Diſproportion what it will ; provided that the Velocities that are given them be reciprocally proportionable to their Maſſes ; that is, if the little Body has as much more Velocity than the great one as it has leſs Matter.

This is the Reaſon why, ſince the Invention of Gun-Powder, battering Rams have been diſus'd in War ; for thoſe and other heavy Machines managed by a great many Hands, and mov'd againſt a Wall with little Velocity, did no more than now is perform'd by a ſmall

a small Cannon-Ball, three or four Men only being employed to manage the Cannon. If the Ball B* weighing 36 Pounds be shot out of the Cannon C, against the Wall AHGE so as to strike it at L, it will produce the same Effect as the battering Ram R, which weighs 4112 Pounds; provided that the Cannon-Ball moves as many times swifter, as it has less Matter than the Ram. See the Calculation of it in the Notes†.

Leet. II.
* Pl. 4.
Fig. 2.

† Ann. 3.

IF a small Pistol Bullet shou'd move with the same Velocity as Light, it would strike as strongly against an immoveable Obstacle as a Cannon-Ball 700000 times as big; because Light moves 700000 times faster than a Cannon-Ball. Light is about 8 Minutes coming from the Sun, and a Cannon Ball wou'd spend ten Years in going thro' the same Space.

10. As the Quantity of Matter in a moving Body, multiplied by its Velocity, gives us the Quantity of Motion: So the Quantity of Motion divided by the Velocity, will give us the Quantity of Matter; but if it be divided by the Quantity of Matter, it will give us the Velocity. If several Bodies of different Weights move with equal Velocity, their Motions will be to one another as their Quantity of Matter.

11. HENCE may be deduced an unanswerable Argument for a *Vacuum*. For if all Bodies, abstracting from the Resistance of the Air, move downwards with the same Velocity, as has been proved*, their Motions compar'd will be respectively as their Quantities of Matter; but the *Motion downwards*, or Force which drives them to the Earth, is their *Gravity*; therefore we find the Quantity of Matter in any Body by its Gravity, which must always be proportionable to it. Now if two Bodies of equal Bulk weigh differently, as we find by Experience they do, there must be Vacuities interspers'd in the lighter: As for Example, if there be two Inch Cubes A and B†, and the Cube A be of Silver, whilst the Cube B is of Cork; it will be found that A weighs 40 times more than B, therefore B has 40 times less Matter and ought to be 40 times less in Bulk, if it had no Vacuities; for if it be answer'd that the Voids or Pores of the Cork are filled with Air and subtile Matter; that Air and subtile Matter* together with the Cork ought to weigh as much as the Silver; or else the two Cubes cannot be equally full.

* L. 1.
No. 8, 9.

† Plate 3.
Fig. 7.

* Ann. 4.

Lect. II. 12. THE whole Effect of mechanical Engines (whereby Motion is given or stopp'd, or a Resistance is overcome) depends upon what we said above, N. 9.

If a great Weight is to be sustain'd by a little one, we must contrive to give the little one so much more Velocity than the great one as it has less Matter, and then its Force being equal to that of the great one, it will sustain it, if they move in contrary Directions; because then equal Forces will destroy each other. This is done by the Contrivance of the Engines, and Manner of applying them. For if the Velocity of the great Weight be determin'd as well as its Quantity of Matter, and the Quantity of Matter in the little one; then the Velocity of the little one (which in this Case is called the Power) must be encreased in the Proportion above-mention'd. But if the Velocity of the Power be determin'd, then that of the Weight must be diminish'd in the said Proportion: The Engines for those Purposes being always so contriv'd, that the Weight or Power may be applied in such manner as to render their Velocities reciprocally proportionable to their Masses.

EXPERIMENT II. *Plate 4. Fig. 3.*

Plate 4.
Fig. 3.

13. LET AB be a Leaver or Balance divided into 20 equal Parts, whose Center of Motion is at C; if the Weight W of 200 Pounds hangs by an immoveable Hook at A, and we would sustain or keep it in *Æquilibrio* by means of the Power or small Weight P of 50 Pounds; it is plain that it has not Force enough at E to sustain the Weight W, because in the Motion of the Leaver the Point E and A will describe not only similar but equal Arcs E e and A a; so that when W moves 1 Inch, P will only move 1 Inch; and since the Velocities being equal, the Quantities of Motion or Forces are as the Masses*, the Weight W will always overpower, as having 4 times the Mass of P. But if P be remov'd to B, then it will describe an Arc similar to, but 4 times greater than that which W describes; that is, it will descend 4 times faster than W rises: If then its Velocity be 4, and that of W only 1, p or 50 multiplied by its Velocity 4 will give 200; which is equal to the Product of W, or 200 multiplied by its Velocity 1.

* No. 10.

14. BUT if the Power P had been immoveable at E, and the Weight W moveable; an *Æquilibrium* would have been had by diminishing the Velocity of the Weight in bringing it forward to D,

D, where being 4 times nearer to the Center than P, it would have had 4 times less Velocity. Lect. II.

15. IF both the Weights had been moveable, and P had been but 12 Pounds and an half, then the Velocity of P must have encreas'd by removing it to B the 16th Division, at the same time as that of W was diminish'd by bringing it to D the first Division on the other Side of the Center; for then W (200) multiplied into DC (1) which is proportionable to its Velocity, wou'd have given 200, which is equal to the Product of p ($12\frac{1}{2}$) multiplied into BC (16) its Distance from the Center, which expresses its Velocity. If therefore the Power has a little more Velocity given it, than in a reciprocal Proportion of the Masses, it will have more Force than the Weight, and consequently raise it, whereas it only sustain'd it before.

16. THUS by means of a Leaver, a Man, whose natural Strength does not exceed 200 Pounds, shall acquire so much relative Force as to raise a Stone of 2000 Pounds by applying his Leaver so as to render the Velocity of the Stone ten times less than that of his Body at the opposite End of the Leaver, which in that Case will be ten times farther from the *Fulcrum* or propping Point, than the Place where the Stone is applied. For if a Man by his natural Strength can raise 200 Pounds with a determinate Velocity, there is no Engine in the World that shall enable him to raise 2000 Pounds with the same Velocity; but he must do it with the 10th Part of that Velocity. If he must employ 10 Seconds of Time to raise 200 Pound 10 Foot; and wou'd raise a Stone of 2000 Pound Weight by a Leaver whose *Brachia* (or Lengths on each side the Center of Motion) are as 10 and 1; he must move 10 Foot at the End of the long *Brachium* of the Leaver whilst the Stone moves 1 Foot, which comes to the same as if the Stone being cut into 10 Pieces, each of them was successively lifted up one Foot by the same Man, who wou'd do it just with the same Labour as when he rais'd it all at once with the Leaver *.

* Ann. 5.

We cannot alter Nature; where we wou'd gain Strength we must lose Time; and where we wou'd gain Time, we must employ more Strength †.

† Ann. 6.

17. IF the Velocity of the Weight, as well as its Quantity of Matter, be determin'd, and likewise the Velocity of the Power; then

Lect. II. then if the Power is not sufficient to raise the Weight, there must be more Matter added to it, till the Product of its whole Mass multiplied into its Velocity be equal to the Product of the Mass of the Weight into its Velocity. If, for example, the Weight * W still supposed equal to 200 Pounds be fixed at A, and the Power *p* equal only to 12½ Pounds be fix'd at B; it will not be possible for the said Power to sustain the Weight, till its Quantity of Matter be quadrupled to make it 50 Pounds, and then 50×16 will be equal to 200×4 ; because here the Distances are as the Velocities.

* Pl. 4.
Fig. 3.

SOMETIMES it is requir'd to give a heavy Body or Weight a considerable Degree of Velocity, as when the Ancients used to throw great Stones with those kind of *Balista* which they call'd Scorpions *; and then the Power must be considerably greater than the Weight; for as it is applied nearer the Center of Motion than the Weight to be thrown, it must be heavier in a reciprocal Proportion of those Distances when the Weight is only sustain'd, and much heavier to give the Projectile Body a sufficient Velocity *. But these Things will be better understood as we come to consider the *mechanical Powers* and the Uses of simple and compound Engines; in order to which we must explain some Terms, and take notice of some Truths necessary to be known by every Engineer.

* Ann. 7.

* Ann. ibid.

DEFINITIONS.

18. A *Weight* is any Body to be sustain'd, rais'd or depress'd, push'd or drawn, or mov'd in any manner; so that the Expression *to raise a Weight* is very extensive in the mechanical Sense; as sometimes it is applied to the driving a Pile into the Ground; sometimes to the stopping of a Body in Motion, as the running of Water, &c.

19. A *Power* is whatever is made use of to raise a Weight in the Sense above-mention'd, whether the Power it self be a heavy Body, a Spring, the Motion of Water, Air, Smoke, Flame, or Pressure of Steam excited by Fire from Liquids; or the Force of any Animal, acting by its Strength or Weight, or both.

20. THE *Intensity of a Power* is its absolute Force, that is, its Force, supposing its Velocity equal to that of the Weight; for its *Moving* or *Acting* Force may be greater or less, according as its Velocity is encreas'd or diminish'd, in respect of that of the Weight. As for Example, If

IF a Man be the Power, and can raise from the Ground a cer- Lect. II.
tain Weight, that Weight will express or be equal to the *Inten-*
sity of the Power; for in this Case, whatever Engine be made
use of, that Part of the Engine, where the Weight is duly applied,
will move just as fast as that which a Man acts upon with his
whole Force. So if a Man can press upon the Point E, at the
4th Division of the Balance AB * with the Force of 200 Pounds, * Pl 4. Fig. 3.
he will just sustain the Weight W, or 200 Pounds, hanging pre-
cisely as far on the other Side of the Center or Motion.

21. THE *Line of Direction* is that Line in which a Weight or
Power acts, or endeavours to act *. * Ann. 8.


22. A Power may act in any Direction whatever; but a Weight
has but one Direction, which is towards the Center of the Earth,
in which Direction all heavy Bodies endeavour to descend, and
actually do, when no Obstacle hinders: So that the *Line of*
Direction of a Weight is a Line drawn from its Center of Gravity
to the Center of the Earth.

23. THE *Center of Motion* is that Point round which a Body or
a Machine moves, or endeavours to move when it cannot or does
not turn quite round; and in that Case, all the Points of the
Body describe Circles, or Arcs of Circles, about the Center of
Motion. *This Center may be taken any where according to the*
Make of the Engine.

24. THE *Center of Gravity* is a Point about which all the
Parts of a Body are in *Æquilibrio*. It is consider'd as the Middle
of the Weight of the Body, though often it is not the Middle of
the Body it self; and if the Body be suspended by that Point, it
will hang in any Position; otherwise the Center of Gravity will
descend as low as it can.

EXPERIMENT III. Pl. 4. Fig. 4.

25. BQ is a round Board suspended by its Center C on the
Points of the springing Calibers A. In turning the Board round,
the two Marks made upon it K and Q, describe the Circles Kk,
and Qq round the Center of Motion C, which is here the Cen-
ter of Magnitude: If the Center of Motion was taken in any
H Point


 Lect. II. Point which is not the Center of Magnitude, as at c^* , still K and Q would describe Circles about the Center of Motion, though the Center of Magnitude wou'd not then be their common Center, but itself would describe a Circle as Cd about c , the Center of Motion. If the Body suspended does not go quite round, as here the Side R stops against the Top of the Calibers at q ; instead of Circles, the Points Q, K and C describe only Arcs of the Circles Qq , Kk , Cd .


* Plate 4.
 Fig. 4. 26. IF C^* be the Center of Gravity of the Body, and the Body be suspended by it, and made to turn round that Point, it may be stopp'd in any Position of the Points K and Q, and it will then remain at rest; but if the Body be suspended by c^* which is not the Center of Gravity, then the Center of Gravity C will descend as low as it can to C.

* Plate 4.
 Fig. 6. BUT if c^* be the Center of Gravity, and placed directly over the Center of Motion K, the Body will remain in that Position, because as the Center of Gravity endeavours to descend in the Line cK , which is the Line of Direction of the Body (in which Line the Point K is supported) it presses directly upon the Point K, which is sustain'd by the Calibers; but if the Body be moved ever so little, so as to bring the Center of Gravity c towards d or e , the Body will turn and not rest till the Center of Gravity comes to M directly under the Center of Motion, the Body falling into the Position $e d f$.

EXPERIMENT IV. Pl. 4. Fig. 8.

* Ann. 9.
 † Plate 4.
 Fig. 8. 27. HENCE may be deduc'd a Method for finding the Center of Gravity of any Body mechanically*. Let AB † be a Body whose Center of Gravity is to be found. If it be suspended by any Part as A, so as to move freely upon the Pin at A, and a Plumb-Line AP hangs from the same Pin, its Center of Gravity C must be under, or rather behind that Line; because it will fall below the Center of Motion A: Let that Line AB be marked upon the Body, as in Fig. 9, and then suspend the Body by any other Part, as F; provided that the Center of Motion be not in the above-mentioned Line AB . Hang on the Plummet at F, and the Line FD under the Plumb-Line FP will cut the Line AB , and shew the Center of Gravity to be at C; for since it must be both in the Line AB and in the Line FD , it can only be in the Point C where they intersect.

N. B.

N. B. *We have not here consider'd the Thickness of the Body ;* Lect. II. *but if we suppose it a Piece of Board as the Figure represents ;*  *then we must only make the same Experiment on the other Side, and we shall find another Point C just opposite to the first Point C. The Line which joins these two Points will be the Axis of Gravity, and the middle of that Line the Center of Gravity.*

28. HENCE also follows, That let the Figure of a Body be what it will, the Body can't fall when its Center of Gravity is sustain'd ; and when any Body is in *Æquilibrio*, its Center of Gravity must be in a Line which goes thro' the Center of Motion and the Center of the Earth ; which is the Line of Direction of heavy Bodies mention'd before *. Thus in *Fig. 7.* † where the * 22. Tobacco-pipe is sustained in *Æquilibrio* upon one's Finger, the † Pl. 4. Point C just over the Finger is the Center of Gravity.

EXPERIMENT V. Pl. 4.

29. IN Bodies that are both regular and homogeneous, the Center of Gravity is just in the middle of the Body ; that is, in its Center of Magnitude ; as appears in the Body AB *, which * Plate 4. being suspended by its Center of Magnitude C, then turn'd ^{Fig. 4.} round, will remain in any Position in which it is stopp'd ; but in *Fig. 5.* * where the same Body is suspended by another Point c, it * Plate 4. will not rest but when the Point C is come down below c, or is directly above it, as we said before. But if the same Body, which was suppos'd a circular Piece of Wood (as for example of Oak) be denser in one Part than another, or be made so by letting a Piece of Lead into the Wood as at M (*Fig. 6.* *) then the Cen- * Plate 4. ter of Gravity will no longer be at C, but at K ; about which Point of Suspension only, the Body will remain in any given Position. If the Body was suspended by the Point C, it wou'd be at rest only in two Positions ; *viz.* when the Lead being carried up to l the Center of Gravity is at c *, or when the Lead is at M, and * 24. the Center of Gravity at K *. If the Body was homogeneous * 24. but not regular, the Center of Gravity wou'd also then be different from the Center of Magnitude. As for example, If you take the Pipe of *Fig. 7.* * and break it at C its Center of Gravity, * Plate 4. you will find by weighing the two Pieces successively, that there is more Matter in the half CB, than in the other half or Shank AC.

Lect. II.



30. As any Body that we consider in Mechanics, is only an Aggregate of several other Bodies or Parts: So the Center of Gravity of a Body is only the common Center of Gravity of all its Parts; and consequently if several Bodies are united in any Machine; or if there be any Combination of Bodies to be sustained, regard is no longer had to the particular Centers of Gravity of the several Bodies which make up the Compound, but only to the common Center of Gravity of the Whole. Thus a Wind-mill must be supported under the common Center of Gravity of all its Parts, and its Line of Direction must fall along the Axis of the Post round which it moves: And a Crane upon a Wharf or a Dock (where the whole Machine turns round) must have the Line of Direction in its Axis *.

* Ann. 10.

EXPERIMENT VI. Pl. 4. Fig. 10.

* 29.

* 29.

31. Let the Line AB represent an even Rod or Wire divided into two equal Parts at the Point C; its Center of Gravity will be at C *. And if two equal Bodies equally heavy are thrust on upon its Ends, so as to have their Centers of Gravity at the same Distance from C, they will be in *Æquilibrio* about the said Point which will then become their common Center of Gravity, and continue so whether the Bodies approach nearer to, or recede farther from it in Proportion to their Masses. The same will happen if the Bodies are unequal as A and b, whose Masses are to each other as Two and One, provided that the greater Body be at A twice nearer to the common Center of Gravity c than the lesser Body b: And c will still be the Center of Gravity of those Bodies, tho' they shou'd move to immense Distances from each other, provided their Distances from the said Point are reciprocally as their Masses, as we said before.

* Ann. 11.

* 45,

32. So that when two Bodies approach towards or recede from each other, with Velocities reciprocally proportionable to their Masses, their Center of Gravity will remain at rest. And if the Bodies being made fast upon the Wire, the Center of Gravity be sustain'd by a Pivot or pointed Broach, how swift soever the Bodies be made to turn round the Center of Gravity and each other, the Center of Gravity will remain at rest *, and the Bodies will describe similar Circles about it and about each other, the one never over-powering the other *. If they be carried forward in any

any


any manner by any external Force acting upon them in proportion to their Masses, their Center of Gravity will go forward uniformly in a right Line, and move just as if the two Bodies were united into one at the said Center: And if they be projected, their Center of Gravity will move in the same Curve as all Projectiles do, whether in their Motion they turn round each other or not. This is evident in the Motion of an Arrow, of Chain-shot, or Bar-shot, and of a Stick thrown from the Hand, the Center of Gravity of any of these Bodies moving in the same Manner as a single Ball wou'd do. So the Moon and Earth, in their Motion round the Sun, do neither of them describe the *Magnus Orbis*; but their common Center of Gravity describes it, in the same manner that they wou'd do if they were both united in that Point, or in the same Manner that the Earth alone is suppos'd to do it, when these Inequalities of Motion are overlook'd, and provided that their Distances from their common Center of Gravity be reciprocally proportionable to their Masses; their Distances from each other may be greater or less in any Proportion: If there were no other Bodies in our System but the Earth and Moon turning round each other, their Center of Gravity wou'd always remain at rest.

EXPERIMENT VII. *Pl. 4. Fig. 11.*

33. IF to the two Bodies A and B there be added a third at D* Pl. 4. Fig. 2. equal to one of the other, let A and B be reduced to their common Center of Gravity, and be considered as a Body equal to both placed at C; then the common Center of Gravity of C and D will be found at K*, as much nearer to C as the Mass of the Body or Bodies at C exceeds that of the Body at D. If the new Body weighed but half as much as the others, it must be removed to *d*, so as to have the Distance K*d*, quadruple the Distance KC. Now if CD be a Wire, and it be supported under K, the three Bodies, whether D or *d* be made use of, will be sustained by that means; only in taking the Center of Gravity exactly at K, we must not consider the Weight of the Wire, whose Thickness we are to suppose diminished *in infinitum* so as to look upon the Wire only as a Mathematical Line without Substance or Weight.

EXPERIMENT VIII. *Pl. 4. Fig. 12.*

34. IF upon this Center of Gravity of the three Bodies, be placed a flat triangular Body *abc* with its Center of Gravity just over

Lect. II.  over K, then a square one as *defg*, then a circular one as *bli*, all in the same manner; still the whole will be sustain'd by supporting the Point K. Whence it appears, that if the whole Weight of a Body be reduced into its Center of Gravity, it will (act as a heavy Body, that is) *gravitate* just in the same manner as it wou'd before.


35. If these three Bodies, united to or acting upon one another in any manner proportionable to their Masses, be carried round their common Center of Gravity; that Point will be at rest, for the same Reason as it will happen in respect of two Bodies; because any Number of Bodies may in this respect be reduced to two. Thus in our System, where the Sun and all the Planets move round their common Center of Gravity, that Center is at rest in the middle of the System. Tho' we commonly consider the Sun as immovable in the middle of the System; because as it has vastly more Matter than all the Planets together, that Center will always be very near the Sun's Center.

36. It often happens that the Center of Gravity of a Body, or of a System of Bodies, is not within the Body itself, or any one of the combined Bodies; yet we are to have the same Regard to its Support, Descent, or Motion in any Direction, as if it was. As for example, let us suppose the Bodies A and *b** to be at the Distance *Ab* from each other, and that *ab* is no longer a Wire but *Ab* is a Line representing their Distance; we shall find their Center of Gravity to be at *c* without the Bodies. And if instead of the Wire CD we suppose the Body D* joined to the two Bodies A and B by the Wires AD and BD, the Center of Gravity at K will be neither in the Bodies nor Wire; so that if we wou'd sustain the said Bodies we must support some one of them, or some Part of the Wire as G, which being made the Center of Motion, the Center of Gravity K will (if the Bodies hang freely) come just under it; or if we support the Point H, the Bodies will be at rest by reason of the Center of Gravity being just over H the Center of Motion. In the Ring ABH* the Center of Gravity is in no Part of it, but it may be supported by any other Point as O or E, &c. Upon this Account the Ring of *Saturn* has its Center of Gravity in the Body of *Saturn*. And tho' the common Center of Gravity of the Sun, Moon, and Earth is within the Sun's Body, yet the common Center of Gravity of the Moon and Earth is in neither of the Bodies, but between them.

* Pl. 4.
Fig. 10.

* Pl. 4.
Fig. 11.

* Pl. 4.
Fig. 13.

35. SINCE OE is the Line of Direction, it is evident that in a-Lect. II.
ny Position of a Body or of a Combination of Bodies, *If the Point of Suspension or Center of Motion be in any Part of the Line of Direction, the Body or Bodies will remain in that Position; otherwise the Center of Gravity will descend as low as it can, and by its Motion alter the Position of the Body.* 

EXPERIMENTS IX and X.

36. SEVERAL odd *Phænomena* depend upon this Principle; as for example, the double Cone or Spindle ACBD* being laid on at E* Pl. 4. upon the lower Part of the Rulers EF will of itself move towards Fig. 14. FF, tho' those Ends are raised up to the Heights FG by the two Screws FG; and by that means will seem to move upwards: And the Cylinder M*, whose Center of Gravity is in the Mid-way between K and O, will actually roll up the inclin'd Plane AC, provided it be hinder'd from sliding by the Rope Rr. How high FG** F. 14. may be in proportion to the Bigness of the Spindle ABCD; or BC* in proportion to the Cylinder M, is demonstrated in the* F. 15. Notes*.

* Ann. 12.

37. BUT before we mention any more of these *Phænomena*, it will not be amiss to shew how the common Center of Gravity of two or more Bodies may be found:

IN two Bodies it is in a Line which joins their particular Centers of Gravity, and this Analogy gives it.

As the Mass of (or Quantity of Matter in) the two Bodies:

Is to the Mass of one of the Bodies::

So is the Distance of the Centers of Gravity of the Bodies:

To the Distance of the common Center of Gravity from the Center of Gravity of the other Body.

FOR example, if we suppose the Bodies A and B* to weigh* Pl. 4. 2 Pounds each, and to have their respective Centers of Gravity Fig. 19. 4 Foot distant from each other; then,

As 4 the Mass of the two Bodies:

Is to 2 the Mass of the Body B::

So is AB, or 4, the Distance of their two Centers of Gravity:

To AC, the Distance of the common Center of Gravity, from the Center of Gravity of A, which is 2 Foot.

OR

Lect. II. OR to express it short by Algebraical Notations $A+B : B :: AB : AC$; but if we take A of 2 Pounds and b of 1 Pound, and Ab , equal to 3 Feet, the common Center of Gravity will be at c within 1 Foot of A , or twice nearer its Center than the Center of b ; because $A+b (3) : b (1) :: Ab (3) : Ac (1)$. Thus is the common Center of Gravity of the Moon and Earth found, when once we know their Masses and Distances. The Earth weighs about 40 times more than the Moon, and the Center of the Moon is about 61 Semi-diameters of the Earth distant from the Earth's Center; therefore the common Center of Gravity of both is distant from the Center of the Earth almost one Semidiameter and a half, or near 2000 Miles above the Surface of the Earth; for as the Mass of the Earth and Moon (41): to the Mass of the Moon (1) :: so is their Distance (61): to the Distance of the common Center of Gravity of both from the Center of Gravity of the Earth, that is, $1\frac{1}{2}$ Semidiameter of the Earth.

Fig. II.

38. IF there be more Bodies than two, as for example, the three Bodies A , B , and D^* ; first find the common Center of Gravity of two of them by the foregoing Rule, and it will be at C ; then say,

As the Mass of the two Bodies considered as united at C , together with the Body D :

Is to the Body D ::

So is CD the Distance of the common Center of Gravity of A and B from the Center of Gravity of C :

To CK , the Distance of the common Center of Gravity of the three Bodies from the common Center of Gravity of the two first.

OR in short $A+B+D : D :: CD : CK$. And if there be a Combination or System of any Number of Bodies; their common Center of Gravity may be found, Step by Step, in the same

* Ann. 13. Manner*.

39. HENCE follows, that one may alter the Place of the Center of Gravity of a System of Bodies, by adding one or more Bodies to it, or by taking one or more of the Bodies away. And the Center of Gravity of a single Body may be removed at pleasure, by adding to or taking from its Mass. And this is of singular Use in Machinery or that Part of Mechanics that relates chiefly to Engines, because of the several Powers that are combin'd
in

in a Machine, and the several Positions that they must have in re-
spect to each other in their Motions. Lect. II.

EXPERIMENT XI. *Plate 5. Fig. 1.*

40. AB* is a rolling Lamp that has within it the two moveable * Pl. 5.
Circles DE and FG, whose common Center of Motion is at K, ^{Fig. 1.}
where their Axes of Motion cross one another, in which Point
also is their common Center of Gravity. If to the inward Circle
you join within the Lamp KC made pretty heavy and movea-
ble about its Axis HI, and whose Center of Gravity is at C, the
common Center of Gravity of the whole Machine will fall between
K and C, and by reason of the Pivots A, B, D, E, H, I, will
be always at Liberty to descend; and therefore let the whole Lamp
be roll'd along the Ground or mov'd in any manner, the Flame
will always be uppermost, and the Oil cannot spill. Thus is the
Compass hung at Sea; and thus shou'd all the Moon-Lanterns be
made that are carried upon a Pole before Coaches or Carriages that
go in the Night.


EXPERIMENT XII. *Pl. 5. Fig. 2.*

41. WHEN inclin'd Bodies, such as the oblique Cylinders ABED*, * Pl. 5.
abed, are set upon an horizontal Plane; they will fall the Way ^{Fig. 2.}
that they incline, if in their Motion that Way, their Center of
Gravity can descend without first rising*. Thus *c*, which is the * 24.
Center of Gravity of the Body *abed*, will descend in the Arc *ck*,
whose Center is the Point *e*, which is the Center of Motion of the
Body when it falls. But *C* the Center of Gravity of the Body
ABED moving round the Center of Motion *E* in the Arc *CK*,
cannot come down to *K* without first rising up to *F* which is a-
bove *C*; therefore the Body can't fall by its own Gravity, because
the Center of Gravity will not rise of itself*. * 24, 28.

42. HERE it is observable that *co* the Line of Direction * of the * 20.
Body *abed* falls without its Base: And *CO* the Line of Direction
of ABED falls within its Base; whence it follows, that inclin'd
Bodies set upon an horizontal Plane will fall, when their Line of
Direction does not go thro' their Base; but they will stand when
the Line of Direction falls within their Base*.

^{Ann. 14.}

Lect. II.

* Plate 5.
Fig. 3.* Pl. 5.
Fig. 2.

THIS is the Reason why an inclin'd Tower, such as that of *Pisa* or *Bologna*, does not fall, tho' its Top hangs so far over the Base as to appear dangerous to those that walk at *o** near its Foot, and don't know upon what Principle it is safe.

43. IF *HI ed* the lower Part of the Body *ab ed** be equal, similar, and alike inclined, to the Body *ABED*, it will not fall for the Reason above-mentioned, its Center of Gravity being then at *Q*; but as soon as the upper Part or Cylinder *ab IH* is set upon it, and made fast by the Pins *ff*, it brings up the common Center of Gravity to *c*, and then the two (which are now become one Body) fall towards *o*; but if *HI ed* being held fast, the upper Part *ab HI* be set upon it, that Part will stand fast without the Help of the Pins *ff*; for as its Center of Gravity is at *G*, its Line of Direction will go thro' the Plane *HI* which is now the Base of the Body *ab IH*. But upon letting go *HI ed*, the upper Part will bring down the lower, and they will fall together, beginning to move round the Center of Motion *e*, the common Center of Gravity *c* descending towards *k**.

* Ann. 14.

EXPERIMENT XIII. *Pl. 5. Fig. 4.** Pl. 5.
Fig. 4.

* 26.


* Ann. 15.

44. IF in the Hand *H** be held upright a Needle or sharp pointed Broach as *C*, the Fork *D*, whose Center of Gravity is at *D*, being set upon the Point of this Needle will be supported*, though it will be difficult to place it right; because the Point of the Needle is so small a Base, that it will require a nice Hand to bring the Center of Gravity so directly over that Base as to make the Line of Direction go thro' it*. But if another Fork, as *B*, be stuck into the Handle of the first, and a third of equal Weight with the second, as *A*, be stuck upon the Points of the first, all the three will be sustained.

* 39.

* 33.

THE Line *AB*, which goes thro' the Centers of Gravity of the Forks *A* and *B*, being bisected at *c*, shews that Point to be the common Center of Gravity of those two Bodies: The Addition of the Fork *D* alters the Center of Gravity*, and causes the Point *C* (which is as near again to *c* as to *D*, the Center of Gravity of *D*) to become the common Center of Gravity of the three Bodies*. In this case the Bodies will be all sustained by means of the Needle under *D*, because the Center of Gravity is as low as it can be: Only with this Difference; that in the Case of the single Fork *D*, the Center of Gravity (which was then over the Center of Motion) wou'd

wou'd descend upon the least Shake, and throw down the Fork; *Leſt. II.* but now no Shake, but what is ſtrong enough to make D jump off from the Point of the Needle, can cauſe the Bodies to fall, for if the Center of Gravity be rais'd out of its Place, it will always return to C the loweſt Point it can deſcend to *.  * 24.

45. SINCE the Line of Direction runs through the ſupported Point under D, it follows, *That a Body, or System of Bodies will be ſuſtained* (that is, their common Center of Gravity won't deſcend) *when any Part of the Body which is in the Line of Direction is ſupported; but will fall when no Part which is in the ſaid Line is ſupported.*

46. IF by a Force imprefs'd upon the Bodies A or B*, or both, * Pl. 5. they be made to turn round each other and round the Center c, Fig. 4. in a Circle whoſe Diameter is AB; they will ſtill be ſupported as before; whether they turn faſt or ſlowly round*; and whether D* 32. be heavy or not, in which latter caſe the common Center of Gravity will return to c; and whether D, C, or c be the Point ſupported: Nay, if the Hand that carries the Bodies ſtands ſtill, or moves in a right or a curve Line, the ſame will hold; that is, the Action of the Bodies upon each other (or in reſpect to each other) will not be alter'd tho' they are carried along with their Center of Gravity: Neither will the Alteration of the Plane in which the Bodies move (that is here the riſing of A and ſinking of B, or the riſing of B and ſinking of A (as they turn) have any Effect upon the Motion of their Center of Gravity. Thus whether the Moon and Earth move faſt or ſlow about each other, and their common Center of Gravity; and whether the Plane of the Moon's Orbit be more or leſs inclin'd to the Plane of the Ecliptick, and whatever may be the Change of that Inclination, the Motion of their common Center of Gravity (which deſcribes the *Magnus Orbis*) will be no way affected.

EXPERIMENT XIV. *Pl. 5. Fig. 5.*

47. IF ſuch a Body as AB* be ſet upon the Peſtetal NDP, it will fall, * Pl. 5. becauſe its Center of Gravity can deſcend (in the Arc C q) or, which Fig. 5. is all one, no part of it that is in its Line of Direction CO is ſupported; but if the two Awls L, M, be ſtuck into the ſaid Body, their common Center of Gravity being at k (in the Mid-way between L and M) will bring back the common Center of Gravity of all the three

Lect. II. Bodies to K, and then K ϕ will become the Line of Direction of the Bodies, in which the Point K being supported the Bodies cannot fall *.

* 45.

EXPERIMENT XV. *Pl. 5. Fig. 6.*

* Plate 5.
Fig. 6.

48. LIKEWISE the Body AB * being set upon the Pedestal AE would fall, its Center of Gravity moving round the Point M in the Arc C q . But if an heavy Body, as D, be join'd to it so as to bring the Center of Gravity to K, then CO will cease to be the Line of Direction; and in the Motion of the Bodies round M the Center of Motion K must describe the Ark K k moving upwards, which cannot be *; therefore AB will be supported by adding another heavy Body to it.

* 24.

49. WHEN Bodies are laid upon an inclin'd Plane, they will come down, though the Line of Direction falls within their Base. In such a Case their Center of Gravity will not move in their Line of Direction (which only happens when Bodies fall freely) but will move in a Line parallel to the Plane, the Body sliding all the while. But if the Line of Direction of the Body falls out of its Base, which is applied to the Plane, the Body will tumble or roll

* Ann. 13. along the Plane *.

EXPERIMENT XVI. *Pl. 5. Fig. 7.*

* Plate 5.
Fig. 7.

HENCE follows that the same Body, that in one Position wou'd slide along an inclin'd Plane, will roll down in another. Thus the Body ABCD*, when set upon the Plane $c M N$, will slide in the Position $abcd$, because its Center of Gravity k cannot fall in the Line of Direction of $k \phi$ (the Body being stopp'd by the Plane) nor move in the Arc kc about d for a Center of Motion; because in this last Case the Center of Gravity must rise, which cannot be *: And therefore the Center of Gravity will descend in the Line $k s$. But if the same Body shou'd be laid on in the Position $abcd$, it wou'd tumble towards M, its Center of Gravity descending in the Arc K q . For this Reason a Column may be drawn up an Hill when laid along in a Waggon, which wou'd tumble backwards if set upright in the same Waggon: And a Load of Hay wou'd be overthrown going along the side of an Hill, where the same Waggon would go along safely, if loaded with

* 24.

with an equal Weight of Iron ; only because in the Load of Iron ^{Left. II.} the Center of Gravity is low ; but very high in the Load of Hay *. ^{Ann. 17.} What has been said in the three last Paragraphs * will be farther ^{47, 48,} confirm'd by the following Experiments. ^{49.}

EXPERIMENT XVII. *Pl. 5. Fig. 8.*

50. UPON the Table T t T * which has a Slit from X to α , set the * Plate 5. little Image DM in such manner that the Saw c (which is fasten'd ^{Fig. 8.} at one end to the Hands of the Image, and has a Weight W fix'd at the other end) may pass through the Slit X α , and the Image will stand in an upright Posture: Then if the Head of the Image be brought down to A or B, it will imitate the Motion of Sawing, and vibrate several times in the Arc A o B whilst the Weight W does in the same manner describe the Arc VWV, the Center of Motion of the whole (that is, of the Image, Saw, and Weight) being at M; the common Center of Gravity K does likewise describe the Arc LKL, till (after having several times descended from L on either side) it comes to settle at K, just under the Center of Motion. If the Image had no Saw, it would stand upright when set on the Table, because its Center of Gravity C would then be just over the Center of Motion M*; and some Part of the I- * ^{26.} image which is in the Line of Direction o O would be supported *, * ^{45.} but the least Alteration of Position that should move c from over M would throw down the Image *. Then if the Saw c be * ^{26.} added; since its Center of Gravity is at c, the common Center of Gravity of the Man and Saw will be at L, and in that Case the Image with its Saw will fall towards X; but if by means of a curv'd Wire the heavy Weight W be join'd to the Saw; the common Center of Gravity of the Man, Saw, and Weight, will be at K*, and the Line of Direction will again be o O, therefore the * ^{38.} Image will stand in its upright Position. If now the Image be inclin'd forwards or backwards, it will after several Vibrations return to its first Position, because the Center of Gravity always endeavours to descend to K, in doing which it will bring the Image upright. N. B. *This Experiment succeeds best with two little Points in the Heels of the Shoes to bear on the Table.*

Lect. II.

EXPERIMENT XVIII. Pl. 5. Fig. 8.

* Plate 5.
Fig. 8.

51. UPON the Stick SS^* (which of it self would fall from the Table, because its Center of Gravity hangs over) suspend the Pail P , fixing another Stick pq , one end in a Notch at p , and the other against the inside of the Pail close to the Bottom, and the Pail without any other Help will be supported on the Stick SS , which will not fall from the Table though the Pail be afterwards fill'd full of Water, provided the Bale or Handle of the Pail be pretty near the Table, and the Stick pq long enough to push the Pail a little out of the upright.

* Plate 5.
Fig. 9.

WHEN the Stick SS^* is horizontal on the Table TdB , c is the common Center of Gravity of the two Sticks $SS PQ$, the Pail DQE , and the Water contain'd in it, all which taken together are to be look'd upon as one Body whose Line of Direction is OO ; and as the part of SS which is a little behind the Bale B is in the Line of Direction, and sustain'd upon the Edge of the Table, the whole Body abovemention'd cannot fall; for if it did, the part BS must rise at the End S , into the Position Bs , and PS descend into the Position ps , which cannot happen unless the Pail rises into the Position dqe bringing up the common Center of Gravity to c in the Arc CcD , which is impossible * from Gravity alone, without the Action of an extrinfecal Agent. But if the Pail DE be lifted up under the Table, and the Stick SS inclin'd above it, so that the whole Machine comes into the Position $ss pe qd$: If then it be left to it self, and the Stick ss is so smooth as well as the Table under B that there is little or no Friction between them, the whole Machine will slide down and so fall from the Table; ss moving in the Direction $sp s$, and the common Center of Gravity c in the Line cE Tangent to the Arc DcC . Here it is observable that as oo is now the Line of Direction, no Part of the Body in the said Line is supported *.

* 45.

N. B. The Experiment is represented in Figure 8.

ANNOTATIONS upon the Second LECTURE.

1. [3. — *By its Velocity.*]

SOME Authors have confounded *Velocity* with *Motion*, tho' it is Annotat. but an Adjunct of it, imagining that one Body must have as much Le&t. II. Motion as another, whenever it moves as fast, and more Motion when it moves faster; but this is only true when the Bodies have equal Quantities of Matter, or when we compare the Motion which a Body has at one time with the Motion that it has at another. The general Definition indeed (*viz.* that *Motion is the Translation of a Body from one Place to another*) takes no Notice of the Quantity of Matter; but when we compare moving Bodies, we have a Regard to their Quantity of Matter, for Velocity alone, without considering how much Matter is mov'd, will never enable us to determine the Force, which we call the *Quantity of Motion*. For example, shou'd a Dog and an Horse moving with equal Swiftnes run against a Wall of the Thickness of a single Brick; the Dog wou'd be beaten back, whilst the Horse that carried 40 times more Matter wou'd beat down the Wall, &c.

2. [5. — *Force and Motion mean the same Thing.*] See Sir Isaac Newton in the Beginning of the first Book of his *Principia*, Def. 2. and 8.

3. [9. — *Less Matter than the Ram.*] If the Battering Ram R** Pl. 4. be 28 Inches in Diameter, and 180 Foot long, made of several Pieces of Fig. 2. Timber, as for example, of Oak joyn'd together, it will contain 750 Cubic Feet of that Timber, which at 50 Pound a Cubic Foot will weigh 37500 Pounds: If the Head of it made of Cast Iron weighs a Ton and a half, that will be 3360 Pounds; then if the five Iron Hoops about it are one Inch thick, two Inches deep and 94 Inches in Circumference, they will weigh about 50 Pound each, which with two Pounds of Nails allow'd to keep them tight to their Places will make 252 Pounds: Now all these Weights added together will give us 41112 Pounds for the whole Weight of the Ram, which if it be mov'd by 1000 Men employ'd only to make it strike against the Point L of the Wall AHIGE (supposing it to swing suspended by its Center of Gravity from a moveable Gallery or only a Tressel) each Man will move a Weight of 41 Pounds. The Quantity of Motion produc'd by this Action, when the Ram moves one Foot in a second, may be express'd by the Number 41112; which Motion or Force compar'd with the Quantity

Annotat. tity of Motion in the Iron Ball B shot out of the Cannon C, will be found
 Lect. II. equal to it: For a Cannon Ball is known to move as fast as Sound for the
 Space of above a Mile, and if you multiply 36 Pound the Weight of the
 Ball by 1142 (the Number of Feet which Sound moves in one Second) you
 will have the Number 41112 to express the Force or Quantity of Motion in
 the Ball B striking at L. And if, after a few Strokes given by the Batter-
 ing Ram, the Mortar or Cement is so loosen'd that the Piece of Wall
 ADDFE is at last by a Stroke of the Ram carried forward from F to K
 and so beaten down; the same thing will be performed by one Cannon
 Ball after the same Number of equal Strokes given before by others, as we
 had supposed by the Battering Ram; and then the Quantity of Motion in
 the Piece of Wall ADDFE carried from F to K will be just equal to the
 Shock of the Ram, or of the Bullet B.

This shews how advantageous the Invention of Gun-powder is; since we
 are thereby enabled to give such a prodigious Velocity to a small Body, that
 it shall have as great a Quantity of Motion as a Body immensely greater,
 and therefore perform as much by its Percussion; tho' we use but few Hands
 in the Management of it, for three Men are able to manage a Cannon which
 shall do as much Execution as the Battering Ram abovemention'd. *Those
 that would have a more ample Account of the Battering Rams and warlike Ma-
 chines of the Ancients may find them describ'd in several Authors, especially in*
 * Poliorch. Justus Lipsius*. *The Ram which I have consider'd here is taken at a Mean,
 Lib. 3. being bigger than some and less than others that we read of.*

4. [11.—*Subtile Matter, &c.*] The Cartesians, in order to maintain
 their Plenum, suppose a certain *subtile Matter* to fill all Spaces between,
 and Pores within Bodies, and that this *subtile Matter* by being continually
 divided, becomes so fine a Dust as to have neither Weight nor Resistance,
 and yet that it is the Cause of Gravity; but any one that considers this as-
 sertion will find it inconsistent with itself. For first if such a Matter fill'd
 all the Interstices between the Parts of Bodies, it wou'd render them equally
 full: *Secondly*, Wherever it fill'd any Space that had no other Bodies in it,
 it must be more solid than Gold and harder than Diamonds, and consequent-
 ly could not be a fine Dust as is imagin'd; for a Body, when solid, differs
 from the same Body in Dust, only because its Parts are separated from one an-
 other, so as to have a great many Voids between: As a Pound of solid
 Gold differs from as much Gold Dust, only because the Parts of the Dust
 are more separated, mix'd with Vacuities, and do not touch in so many Points
 as the solid Lump; the Dust easily becoming a solid Lump when the Vacui-
 ties are driven away by Fire, which turns the Dust into a Fluid that after-
 wards (upon the Removal of the Fire) is chang'd into a solid Lump with-
 out losing any of its Weight. *Thirdly*, As the Cartesians affirm — That of
subtile Matter the Earth and Air (heavy Bodies) are compounded, it is
 absurd to suppose the Matter of which they are made to be without Weight;
 since the Gravity of any whole Body is made up of the Gravity of all its
 Parts taken together. *Fourthly*, When the Cartesians make their *subtile*
 Matter

Matter the Cause of Gravity, they seem to forget what they have said of it before, namely, *that it has no Resistance*; for if a Fluid, by its Motion about the Earth, impells all the Bodies near it to fall down to the Earth, it cannot be void of Resistance, because whatever *impells* must *resist*: And to say, that solid Bodies can move thro' this subtile Matter without suffering any Resistance, but that the subtile Matter when it runs against solid Bodies drives them out of their Places, is an Absurdity unworthy of a Philosopher. Yet because the *Cartesians* cannot deny the Experiment of a Piece of Gold, and a Feather falling equally fast in a Glass Receiver out of which the Air has been pump'd * (and the Experiment has been made in a Receiver 10 Foot* L. 1. 8. 9. high; see *Phil. Transf.* No. 354) rather than give up their *Plenum*, they say, that when the Air is pump'd out, the Receiver is as full as before, but that it is fill'd with such a subtile Matter as makes no manner of Resistance: So that upon the whole, the *Cartesian* Account of the Cause of Gravity, and the Non-resistance of their subtile Matter, are inconsistent with each other. Many more are the Contradictions which they fall into by endeavouring to solve *Phænomena* by the Powers and Motions in all Directions which they attribute to this subtile Fluid; but I shall say no more on this Subject, till I come to speak of the Motions of the heavenly Bodies.

5. [16. *At once with the Leaver.*] It was upon this Principle that *Archimedes* propos'd the lifting of the whole Earth, in case that there could be found a *fix'd Point*, or Place to support his Instrument, *ὅς τις πᾶσι τοῖς κοσμοῖς κινῆται*. By which he meant—That the least Power, by encreasing its Velocity, might raise the greatest Weight; and that in this Respect there are no Bounds, where we can get a Place for a *fix'd Point*, and a due Distance for the Power. Now, tho' raising the Earth is a Proposition purely Mathematical, and not to be reduc'd to Practice; yet for Curiosity Sake we'll consider it a little here.

If we take the common Center of Gravity of the Earth and Moon for the *fix'd Point* (*Fulcrum*, or propping Point) of our Leaver, which we will suppose 240000 Miles long, that is, reaching from the Center of the Earth to the Center of the Moon; then the Moon, or a Weight equal to it made use of as a Power, will be able to support the Earth at the other End of the Leaver; and if it be removed but an Inch farther, it will raise it up.* Here* No 15. the Distance of the Center of Gravity of the Earth from the *fix'd Point* of the Leaver is 6000 Miles, and that of the Center of Gravity of the Moon or Power is almost 40 times as far; and if the Moon be supposed to move with the same Velocity that it wou'd fall to the Earth by the Force of Gravity, if its projectile Force did not hinder it; then the Earth would be moved one Inch out of its Place, by the Moon's moving about 40 Inches.

Now if, instead of the Moon, we make use of a Power whose Intensity is equal only to a Weight of 200 Pound, as for Example, the Force of a Man, as *Archimedes* proposes: Then still supposing the Earth at the Distance of 6000 Miles from the *fix'd Point*, that *Brachium* of the Leaver to which

K

the

Annotat. the Power is applied, must be lengthened in the Proportion that the Weight
 Le& II. of the whole Earth bears to 200 Weight. In this case the End of the
 Leaver will reach quite out of our System among the fix'd Stars, above fifteen
 thousand Millions of Millions of times farther than the Distance of *Saturn*:
 And if *Archimedes* (or the Power) be suppos'd to press upon the Engine
 with the Velocity of a Cannon Ball. He wou'd be in Motion at the End
 of the Leaver above 26 and near 27 Millions of Millions of Years to raise
 the Earth one Inch in that Time, and wou'd go thro' a Space above 39
 thousand Millions of times greater than the Periphery of *Saturn's* Orbit.

For the Sake of such as wou'd examine this Calculation, we subjoin the Numbers that we made use of.

The mean Diameter of the Earth is equal to 19 688 725 Paris Feet
 (*Newton. Princ. Lib. 3. Prop. 20. Pag. 387.*)

Supposing the Diameter to the Circumference as 7 to 22; and multiplying
 the Diameter by the Circumference, you will have 1 218 315 660 966 250
 Square Feet for the Surface of the Earth: Which last Number, multipli-
 ed by the sixth Part of the Diameter, will give 3 997 847 001 180 744
 647 897 $\frac{1}{2}$ for the Cubic Feet contained in the whole Earth.

Now if we suppose a Cubic Foot of Earth to weigh 100 Pound, mul-
 tiplying by 100 we shall have the Weight of the whole Earth in Pounds,
viz. 399 784 700 118 074 464 789 750.

Then, As 200 Pound (or the Intensity of the Power):

Is to 399 784 700 118 074 464 789 750 (or the Intensity of the Weight): :

*So is 6000 Miles (the Distance of the Weight or of the Center of the Earth
 from the fix'd Point):*

To 11 993 541 003 542 233 943 692 500 Miles (or the Distance of the Power).

This last Number not only expresses the Distance of the Power, but the
 Number of Miles that the Power must move to raise the Earth one Mile;
 because the Velocities of the Power and Weight must be reciprocally as their
 Masses. But if we wou'd raise the Earth but one Inch, we must divide by
 66 360 (the Number of Inches in a Mile) and we shall have the Miles gone
 thro' by the Power whilst the Earth moves 1 Inch, *viz.* 189 291 996 583
 668 465 020 Miles.

To have a clearer Idea of what has been said, let us compare the Distance
 of the Power and Space that it must go thro', with some great Distance
 that we know, and with the Space describ'd by some Body that we can ob-
 serve; as for example, with the Distance of the Planet *Saturn*, and the Space
 which it describes in going thro' the whole Circumference of its Orbit.

Saturn, at its mean Distance from the Earth (which is equal to its Di-
 stance from the Sun) is above 9 times and a half (or 9,51 times) farther from
 the Sun than the Earth, which last we suppose 81 Millions of Miles from the
 Sun; and therefore will be express'd by this Number of Miles (*viz.*)
 770 310 000, by which dividing 11 993 541 003 542 233 943 692 500,
 we shall have a Number which shews us that the Power must be applied
 farther

farther from the Fulcrum or Center of Motion than *Saturn's* Distance, Annotat. 15 569 745 951 035 731 times. Then if the Space that the Power must go thro' (or 189 291 996 583 668 465 020 Miles) be divided by 4 841 948 571 the Number of Miles in the Periphery of *Saturn's* Orbit, the Quotient will shew us that the Power will go thro' a Number of Miles 39 094 177 438 times greater.

Again, if we suppose the Power, or *Archimedes* pushing forward the farthest End of the Leaver; we shall find, that tho' he should move as fast as a Cannon Ball, he wou'd spend 26 978 123 942 460 Years in moving the Earth one Inch. For if we suppose a Cannon Ball to move with the same Velocity as Sound, or a Mile in 4 Seconds and an half (as Experiments have confirm'd it) it must move 800 Miles in an Hour: And as one Year contains 8766 Hours, 800 times that Number is the Number of Miles which a Body with the Velocity of a Cannon Ball would move in a Year; by which Number (7 012 800) if you divide 189 291 996 583 668 465 020 (the Number of Miles that the Power goes thro') you will have the Number of Years which it must take up to go thro' the said Space, namely 26 978 123 942 460 Years, which was to be proved.

Bishop Wilkins in his *Mechanical Powers* mentions a compound Engine of Wheels, whereby the Earth might be rais'd, without supposing the Power to be applied at any considerable Distance from the Earth; but what he says depends entirely upon the same Principle; for even in that Case, the Power (if equal to what we have suppos'd) must go through a Length equal to as many Miles as we have mention'd above, in order to raise the Earth but one Inch; though the Line in which it mov'd should be but a Circle of a Foot in Diameter; for let the Machine be made in any manner whatever, the Difference of the Velocities of the Power and Weight will always be reciprocally as their Masses (or their Intensities) considering the Power as a Quantity of Matter in Motion when they balance one another; and the smallest Addition of Velocity (still suppos'd in these Calculations) will make the Power overpoise.

6. [16.—Where we would gain Time, we must employ more Strength.] For want of rightly considering this, a great deal of Time and Money has been fruitlessly spent in mechanical Works, by such as imagine that Force might be generated by the Figure of a Machine; whereas Mechanics teach us not to make, but to apply Powers, such as we find them in Nature; for we deceive our selves, if we think that by means of any Engine whatever, One Man shall do the Work of Two in the same Time, supposing the Men to employ the same Strength. But yet the Science of Mechanics is not to be rejected as useless; for to the practical Arts that are deriv'd from that Science we are indebted for a great many of the Necessaries as well as Conveniencies of Life. In the Performance of several Works, where we have sufficient Strength, we often want Time; and sometimes where we have Time to spare we want Strength. In such Cases the Skill of a good Mechanick is to be exerted in directing the Application of the Powers according to Time. Thus in making Harbours, and carrying on Dignes, Moles

Annotat. or Banks, where at every Tide the Sea may damage the Work, and a
 Lect. II. Spring-Tide over-set it; the greatest Number of Hands *must be employ'd*
 ~~~~~ *that can work by one another.* In some Cases, as raising Blocks of Marble  
 or other heavy Goods out of Ships to lay them upon a Wharf, many  
 Hands cannot be employ'd; either because they cannot well stand by one an-  
 other about the same Block; or because they cannot lift all at once; or  
 when they have got up their Burthen, they cannot conveniently walk with  
 it; or if they could, the Planks over which they go, or Ladders which they  
 must climb, could not support them and the Weight; then an Engine must  
 be us'd (as for Example, a Crane) where one Man shall do the Work of  
 10 or 20 Men, but he shall be 10 or 20 times longer in performing it. Yet  
 the Engine is absolutely necessary; because without it the Work could not  
 be done, therefore a sufficient Time must be employ'd, without which a  
 great Strength would be of no Use. Thus likewise in Building we must use  
 Engines to raise great Stones and large Pieces of Timber, where so much  
 more Time is employ'd according as the Force of the Men working at the  
 Engine is less than the Force that would be requir'd to raise the Stone if  
 the Hands were applied directly to it; but then the Engine takes up less  
 room, and the rest of the Men may be otherwise employ'd. In draining  
 Mines we are always confin'd in Time, because the subterraneous Springs  
 supply the Water whilst a Force is employ'd to draw it out; and in such a  
 Case the Power (that is, the Intensity of it) must be superior to the Quanti-  
 ty of Water to be rais'd in a certain Time; that is, the Power must be a-  
 ble, without any Engine, to draw from the Bottom to the Top of the Pit  
 (suppose an Horse drawing up a Bucket fasten'd to a Rope that runs over  
 a single Roller) a Weight greater than the Weight of the Quantity of Wa-  
 ter that runs in, during the Time that the Power goes through a Space e-  
 qual to the Depth of the Pit. Engines are applied for the Conveniency of  
 delivering the Water, and not to gain any Degree of Force; for we always  
 lose some — So much as is employ'd to move the Parts of the Engine,  
 which cannot be apply'd to one another without a Friction that spends  
 some of the Power; so that the best Engine is that which consists of the  
 fewest Parts\*. And he that by Machinery pretends to gain by bringing  
 up a greater Weight, or a larger Vessel of Water with the same Power,  
 does not consider, that it will rise so much the slower, and give Time to  
 the Springs to supply more Water in Proportion; or if, by adding any Part  
 to the Engine, the Power (as for Example, an Horse or Horses) is made  
 to go easier, then less Water will be rais'd at a Time. *This shou'd* give a  
 caution to those that have any Concerns in Mines, Water-works, Mills, or  
 other Manufactures; that they might not be impos'd upon by Engine-makers  
 that pretend to (and often fancy they can) by some new invented Engine  
 outdo all others, and make one Horse do as much as three or four.  
 This proceeds from their being unskill'd in Mechanical Principles, the

\* All that relates to raising Water shall be particularly consider'd, when we come to treat of Hy-  
 draustic

Knowledge of which would keep them from attempting Impossibilities. It Annotat. were to be wish'd that our Engine-makers, who often abound with Inven- Left. II. tion; and are generally quite ignorant of Mathematics, wou'd apply themselves to that Science; at least to know so much as wou'd direct them in their Works; or that some of our best Mathematicians wou'd not think it below them to direct Workmen, and consider Engines a little more than they do, which wou'd render their Speculations more useful to Mankind. There are some, who, being too clumsy, and wanting a nice Hand to make Experiments, are unwilling to own it, and therefore ridicule and despise Mechanical Performances; forgetting that the incomparable Sir *Isaac Newton*, whom They with all other Philosophers admire, has made as many as (if not more Experiments than) any Man living; and look'd upon Geometry as no farther useful than as it directs us how to make Experiments and Observations, and draw Consequences from them when made; so that the Improvement of Philosophy must be the Result of mix'd Mathematics, that is, of Mechanics and Geometry. A Man that shou'd learn to fence by Book, wou'd be as much at a Loss if he was call'd to fight, as another that shou'd prefer brutal Courage to the whole Art of Fencing; only with this Difference, that the latter wou'd be much more likely to kill his Adversary; as Men quite illiterate have often produc'd wonderful Engines. The Machine at *Marly* which was made by an ordinary Man of *Liege*, who was entirely ignorant in Mathematics, has a great many excellent Contrivances; but does not raise all the Water that it shou'd do, because the Workman did not know how to calculate, so as to give the Power of the River *Seine* its utmost Advantage.

When great Works or Manufactures are carried on in such a Manner that a great Part of the Intensity of the Power is uselessly spent, and but little of it employ'd in doing real Services; as for Example, when there are unnecessary Frictions arising from the ill Contrivance of the whole Engine or the wrong Figure of some of its Parts, or the bad Performance of the Workman; or if Men or Horses, &c. exert but a small part of that Strength which they might apply without Weariness or Inconveniency: Then the Skill of a good Engineer may be advantageously applied in changing the Form or altering the Parts and Motions of a Machine. An Instance of this may be seen in the winding of Thread or Silk, in which Business if 50 Men being employ'd they should only move a Weight or overcome a Resistance equal to half a Pound, as they carry their Hand round (whereas one Man can easily raise 25 Pound, with the same Velocity as his Hand moves during the Space of 10 Hours in a Day) a Machine may be contriv'd whereby one Man applying his whole Strength shall do the Work of the fifty Men in the same Time. Thus in any other Case Machines may afford us great Profit, by rendering effectual that Force or Intensity of the Power, or Powers, which was before misapply'd or not employ'd at all. This has been perform'd at *Derby* in a very ingenious Manner by Messieurs *Thomas* and *John Lombe*, who have employ'd the Force of a Water-wheel to the working of *Italian* Silk, so as to lose or mispend no Part of the Power.

As.

Annotat. As the Proprietors of this curious Machine are not willing to have a Description of the whole Engine or any of the Movements of it made Publick, I shall only here give the general Account of it, viz. That there are

26 586 *Wheels.*  
 97 746 *Movements.*  
 73 728 *Yards of Silk wound every time the Wheel goes round, which is three times every Minute,*  
 318 504 960 *Yards of Silk in one Day and Night, and consequently*  
 99 373 547 550 *Yards of Silk in one Year.*

*One Water-wheel communicates Motion to all the rest of the Wheels and Movements, of which any one may be stopp'd separately and independent on the rest. One Fire-Engine conveys Air to every individual Part of the Machine, and one Regulator governs the whole Work.*

\* Polior.  
 Lib. 3.  
 Dial. 4.  
 † Pl. 1.  
 Fig. 6.

7. [17.—*Scorpions, &c. give a sufficient Velocity, &c.*] Scorpions were Machines to throw Arrows, Fire-Balls, or great Stones. A Description of them may be found in *Vitruvius* and the above-mention'd *Lipsius*\*, from whom I have taken the Figures represented in Plate 6. The 1st Figure represents one of these Machines charg'd. The Point A of the longest Brachium AC, which in the natural Situation is kept uppermost by the Boxes of Stones or Weights BB, having been brought down to A (by the Rope RR and Loop a, drawn by Help of the Wheel W and Pinion at I round the Rollers M and L) is kept from rising up again by the Pin HH, made a little taper: Then the Loop a being taken off from A, and the Sling S being charg'd with the Ball or Stone T, the Scorpion is ready to be discharg'd; which is done by a smart Blow of an Hammer on the End H; of the Pin, or a sudden drawing it out with a Rope, for then A being no longer kept down rises with great Velocity by the Descent of the Weights BB, and one of the Loops of the Sling slipping off of the Point A, made conical for that Purpose, the Stone flies out as represented in the second Figure, which is another Scorpion little differing from the former. All the Difference is, that in this last Figure, as the discharging End A is the Axis of Motion DD, then the same End in the former Figure, the Pulley \* L is applied in such a Manner as to cause the Handle I of the Pinion leading the Wheel W, to move as easy again as in the former Case: And as to the Effect, supposing the Weights BB, equal in both Scorpions; the latter will throw a Ball of greater Weight, but then it will have less Velocity than the former Projectile. In both Cases the Scorpion turns upon the Pivot C, and the whole Frame HI round the upright Shaft Cc, that the Machine may be directed any Way. The Hook H in the second Figure does the Office of the Pin H in the first.

\* Pl. 6.  
 Fig. 1.

However powerful these Machines were, and however numerous, they are not to be compared with a Battery of Cannon, either for Force or Expedition. Bishop *Wilkins* had not preferr'd them to our Artillery, as some others

thers have done, if he had examin'd them less superficially, and calculated their Force. Annotat. Lect. II.

To prove this, we will here consider the Force of one of these Scorpions, even supposing it much larger than cou'd conveniently be carried about (for, that they were probable, appears from the Account that *Cæsar* had great Numbers of them in his Camp) and we shall see how much short it will fall of the Force of a Cannon.

Let \*AD the Tail of the Scorpion, or the End that throws the Stone\* Pl. 6. or Bullet, be supposed 24 Foot long; and the shorter *Brachia* DB which Fig. 1. carry the Boxes BB, 8 Foot long each. When the Boxes are fill'd with Stones so as to weigh, for example, 1000 Pound each, and the Tail brought down to A in order to throw the Stone T; the utmost Velocity that can be given to it can never exceed 48 Feet in a Second, because the Weights at B cannot fall faster than 16 Feet in a Second, tho' they shou'd not move the Machine in their Descent; but as we may reasonably suppose that throwing up the End A loaded with the Projectile T, must retard them one half, it will follow that the Body T will be thrown forward only at the Rate of 24 Feet in a Second, which is about 48 times slower than the Motion of a Cannon Ball; and therefore the Effect of the Scorpion will be 48 times less than that of a Cannon throwing a Ball of the same Weight. Besides, there must be more Men to manage the Machine, and a great deal more Time spent in drawing down A by means of the Wheel W, than in charging a Cannon; for if we suppose the Force requir'd to raise the Boxes loaded with Stones, and to overcome the Friction, to be equal only to 2500 Pound, and the Engineer who turns the Handle at I to move his Hand thro' a Space of  $3\frac{1}{2}$  Feet in a Second (which is the utmost he can do if the Force he applies be equal to 25 Pounds, he must spend 7, 6 Minutes or above  $\frac{1}{8}$  of an Hour to carry the Handle thro' a Space of 1600 Feet, in order to raise the Boxes 16 Feet in Height: This, besides the Time spent to put in the Pin H and to fit the Stone T into the Sling S, will so retard the Operation, that the Scorpion cannot discharge its Projectile above 4 times in an Hour, whereas a Cannon may be fir'd with great Ease twice as many times in an Hour.

Now if we consider how much such a Scorpion as we have describ'd must weigh, it will appear as troublesome to carry as a Cannon. The upright Shaft C must be 30 Foot long; and that it may have sufficient Strength, we will suppose it of  $13\frac{1}{2}$  Inches in Diameter, which will make it contain 30 cubic Feet of Timber, which if of Oak will weigh about 50 Pound a Foot; let the Body and Tail of the Scorpion ABB with the Boxes BB, contain 40 Foot more of the same Timber; the Frame KH 60 Foot, and the Wheel W with the Pinion on the Handle I, the Pullies, Ropes and Iron-work of the Machine weigh as much as 60 cubic Feet more of Oak: All this together will make 90 cubic Feet, which multiplied by 50 Pound will give 4500 Pound, a Weight which will render the Machine very inconvenient, even tho' it be taken to Pieces. We may therefore suppose the Scorpions much less than what I have describ'd: So that there can be no Comparison between their Effect and that of our Artillery. Whoever will be at the Pains

of

Annotat. of calculating the Force of any other of the Machines used by the Ancients,  
 Lect. II. will find that they fall very short of the Effects of Gunpowder: Especially if  
 we consider what Force it exerts in the springing of a Mine, whereby prodigious Rocks are rent in Pieces, and such Quantities of Earth and strong Walls lifted up, that all the Machines made use of in a Roman Army, if they cou'd be apply'd at once to one Part of a Fortification, wou'd produce nothing like this new Invention of a portable Powder which contains such an immense Force in so small a Compass.

\* Pl. 7.  
 Fig. 1.

8. [21.—*Line of Direction, &c. endeavours to act.*] Though an heavy Body as A\*, may by a Force or Forces impress'd upon it, be made to move in any Direction, yet (as it always retains its Tendency towards the Center of the Earth, which, if no Obstacle hinder'd, would carry it the nearest Way) its *natural* Line of Direction always goes thro' the Center of the Earth; and if, whilst the Body is going down by its own Gravity, it be hinder'd by one or more inclin'd Planes, as BG, BD, and carried down afterwards in a Curve (+6) in its Descent on account of the Alteration of the Direction of its Motion, or carried up again along another Plane as DE; we are not to call the Line 1 2 3 4 6, or 1 2 3 4 5 which it describes, its *Line of Direction*; but when the Body comes from the Point 1 successively to the Points 2, 3, 4, its Lines of Direction are 1 c, 2 c, 3 c, 4 c, which are all directed to the Center of the Earth, and by reason of their great Distance from it, may be consider'd as parallel, or indeed as one and the same Line.

\* Pl. 7.  
 Fig. 2.

Now the Line of Direction of a Power varies according to the Application of the Power, whatever Line the Body acted upon moves in. Thus when the heavy Body A\* (whose Line of Direction is c C) suspended by a Rope, is held by the Hand at H, the Line of Direction of the Power is the same with that of the Weight; but if the Rope be brought over the Pulley B, the Lines of Direction of the Power may be any of the Lines BG, BF, BE, BD, whilst the Line of Direction of the Weight still continues the same. Nay, if the Body A\*, being acted upon by the Power at I, be made to move in the Line c D, along the Plane MB, c D is not the Direction of the Power but AI. And if the said Body be raised from E to F by means of the Wedge KFL drawn under it (whilst a Board or immoveable Plane at HG keeps it from going out of the Line EF) the Line of Direction of the Power will be LB: But if the Body having been plac'd at K upon the horizontal Plane L, the Wedge or inclin'd Plane F be suppos'd immoveable, and the Plane HG to move from K to G, and to push up the Body in the Line KG, then does the Power acting in the Line of Direction KG cause the heavy Body to rise the Height EF, whether it moves in the Line KG or directly up in the Line EF to get that Height.

\* Pl. 7.  
 Fig. 3.

Hence it follows that the Velocity of a Power is not to be consider'd in the same manner as the Velocity of a Weight (unless when a Power ascends or descends directly from or towards the Center of the Earth) for the *Velocity of a Power is the Space that it goes through in a certain Time,* which

which may be greater or less to perform the same Operation, according to Annotat. the manner in which it is applied; as the Power B \* moves only the Le<sup>ft</sup>. II. Length LK, when it raises the Weight up through the Line EF, by drawing the Wedge LFK through EF; but if it pushes it up along FK sup-<sup>\* Plate 7.</sup> pos'd immoveable, it must move the whole Length of the Diagonal FK of<sup>Fig. 3.</sup> the Triangle LFK. But whether the Weight rises from K or from E to the Point F, its Velocity must only be call'd EF, because whatever Line it runs thro', it only rises (or removes farther from the Center of the Earth the Height EF. So likewise the Velocity of A \* is only the Line i c, when<sup>\* Plate 7.</sup> the Body runs thro' a much longer Space, namely from i to c through the<sup>Fig. 1.</sup> Points 2, 2, 4 along the Planes BG, GD. Therefore the Velocity of a Weight is always to be measur'd by the Line of its upright Ascent or downright Descent, which shews how much it is got nearer to or farther from the Center of the Earth.

*We are to observe that this Definition relates chiefly to that Part of Mechanics which considers the Actions of Bodies upon one another by the Application of Instruments.*

9. [27.— *A Method for finding the Center of Gravity, &c. Mechanically.*] If a Board, of equal thickness all over, be laid upon the Edge of a Triangular Prism P p \*, or upon the sharp Edge of any streight Body placed in an<sup>\* Plate 7.</sup> horizontal Situation, so as to be in *Æquilibrium*; whatever Bodies are laid<sup>Fig. 4. & 5.</sup> upon such a Board in such manner that they do not alter its *Æquilibrium*, must have one Plane of Gravity (that is, a Plane in which their Center of Gravity is) directly over the Edge which supports the Board. Another Position of the Body, if the *Æquilibrium* be still preserv'd, will give another Plane of Gravity, the Section of which Plane will give an Axis of Gravity, or a Line which has the Center of Gravity in it. A third Position of the Body may be obtain'd so as to find a third Plane of Gravity, which shall cut the other two at right Angles, or any large Angle, and the Intersection of the three Planes will give the very Point which is the Center of Gravity. If the Body, whose Center of Gravity you would have, be long and flexible, so as not to lye across upon the Board above-mention'd, a second Board must be laid upon the first with a Pin in its Center, so that it may turn quite round without altering the *Æquilibrium*; and then long Bodies laid upon this last Board may easily be mov'd on the Board so as to find their different Planes of Gravity. Thus one may find the Center of Gravity of an humane Body,<sup>Plate 7.</sup> or of any Animal. In relation to the humane Body, it is observable, that whe-<sup>Fig. 4. & 5.</sup> ther a Man be fat or lean (nay in a Skeleton) the Center of Gravity is always near the same Place, viz. in the *Pelvis*; between the Hips, the *Os<sup>s</sup>a Pubis*, and the lower part of the Back-bone. Raising up the Arms and Legs will raise up the Center of Gravity a little; but still it is always so plac'd that the Limbs move freely round it, the Center of Gravity at the same time moving much less than if it was in any other Part of the Body. A Statue, though it represents a Man, has not its Center of Gravity in the same Place as a Man; for if it be hollow, the Hollows are not in the same Places as in a Man's Body, and the Center of Gravity in a  
L solid

Annotat.  
Leſt. II.



ſolid naked Statue is higher than in a Man. I mention this, becauſe in ſetting up a Statue, and fixing it (eſpecially in a Place expoſed to the Wind) great regard is to be had to place the Center of Gravity over the Middle of the Baſe; or, if the Attitude of the Statue does not permit it, then the Statue is to be ſecur'd moſt ſtrongly on that Side which is fartheſt from the Center of Gravity.

Mathematicians, in order to ſettle Rules for finding the Center of Gravity of Bodies, firſt give Methods for finding the Center of Gravity of 2 or more Lines, then of the Periphery of Figures, then of Planes; for tho' Lines and Surfaces do not exiſt ſeparate from Bodies, yet they conſider Lines as ſlender homogeneous Bodies, and Planes as extremely thin Solids; and, from that Conſideration, more regularly proceed to find the Center of Gravity of Solids. Dr. Wallis has fully treated of this Subject in his Mechanics, in his Chapter *de Inveſtigazione Centri Gravitatis*, and Monſ. Ozanam in the third Chapter of his *ſtatics*, in the fourth Volume of his *Course of Mathematics*.

I ſhall give here ſome of the moſt eaſy and uſeful Methods, and muſt refer the more curious Readers to the Authors above-mention'd, and other Mathematicians that have particularly conſider'd the Center of Gravity.

\* 29.  
\* Plate 7.  
Fig. 6.

If a Line be conſider'd as an homogeneous Wire infinitely diminifh'd, its Center of Gravity will be in its Middle Point\*; as the Point I in the Line AB\*. Let there be another Line as CD\* in any Poſition in reſpect of AB; then if from the Center of Gravity I of AB, a Line (ſuppoſed without Gravity) be drawn to K the Center of Gravity of CD, you will have their common Center of Gravity at G by this Analogy,

\*\* 38.

$$** AB \times CD : CD :: KI : IG.$$

And if the Second Line had been leſs than AB, as for example, if FE had been taken inſtead of CD, the common Center of Gravity would have been at H; becauſe \*\*  $AB \times FE : FE :: KI : IH$ .

\* 39.

If there be three Lines (whether they include a Space, ſo as to make the Periphery of a Triangle, or not) their common Center of Gravity may be found after the ſame manner as that of three Bodies\*. So likewise may be found that of 4 or more Lines, and conſequently of Polygons.

\* Plate 7.  
Fig. 7.

It is to be obſerv'd that the Center of Gravity of plane Surfaces is not the ſame as the Center of Gravity of their Peripheries, unleſs when they are regular. Thus in the Triangle\* ABC, which is not equilateral, the Center of Gravity of the Periphery will be found at H, nearer to the Angle B than I which is the Center of Gravity of the Triangle. For (by 39) D and E being the Centers of Gravity of the two Lines AB and BC, F is found to be their common Center of Gravity, and H the common Center of Gravity of the three Lines AB, BC, and CA.

To find the Center of Gravity of a Triangle, draw a Line from the Middle of any Side of its oppoſite Angle as GB: Let GI be taken =  $\frac{1}{3}$  of the ſaid Line, and the Point I will be the Center of Gravity of the Triangle. Now as every rectilineal Figure may be divided into Triangles, the common Center of all the Triangles will be the Center of Gravity of the Figure. Let

Let \* AB and CD be two Quantities (whether Surfaces or Solids) whose particular Centers of Gravity are at F and G, the Center of Gravity of their Sum, or of AD, will be at \* E; but if you would have the Center of Gravity of the Difference of two Quantities, their particular Centers of Gravity being known; (as for Example, the Center of Gravity of CD which is the Difference of the two Quantities AB and AD, whose particular Centers of Gravity are F and E;) draw FE and produce that Line towards G, and you will find the Point G the Center of Gravity requir'd, by this Analogy  $CD : AB :: FE : EG$ . That is, *As the Difference is to the least Quantity; so is the Line FE: to the Line EG, or the Length of the Production of the Line FE.* Annotat. Left. II. \* Plate 7. Fig. 8. 39.

The Center of Gravity of a Cone is in its Axis, at the distance of one fourth Part from the Base; as for Example, in the Cone ABC\*, whose Axis is DC, the Center of Gravity will be at F, FD being equal to  $\frac{DC}{4}$ . \* Plate 7. Fig. 9. But in a Conic Surface the Center of Gravity will be distant from the

Base  $\frac{1}{3}$  of the Axis; that is, DF will be equal to  $\frac{DC}{3}$ .

If ABIK be a truncated Cone, to find its Center of Gravity, let the Cone be compleated, which will then be ACB; then having found the Center of Gravity of the Cone ABC (namely, the Point F) and the Center of Gravity of the Cone ICK (which is the Point E) joyn those Centers together by the Line FE; then considering that ABKI is the Difference of the two Quantities ACB and ICK, you will find its Center of Gravity by the Rule above-mention'd, which will be at G; for  $ABIK : ICK :: EF : FG$ .

If a Bucket be made of Copper, Tin, or Wood, in the Shape of a truncated Cone, the Center of Gravity of such a Vessel will not be in the same Place when it is empty as when the said Vessel is full; which Consideration is useful in several Cases of Mechanics in general, and Hydraulics in particular. For by that means Vessels made of the Shape above-mentioned, which being suspended and moveable upon Pins or an Axis (passing between the Center of Gravity of the empty and the Center of Gravity of the full Vessel) shall turn with the Bottom upwards when empty, will be drawn directly up with the Bottom downwards when full; or on the contrary have their Mouth upwards when empty, and turn over and empty themselves as soon as they are quite full.

Let ABED be the Section of a truncated hollow Conic Vessel, whose Mouth is AD. Its Center of Gravity by the Rules above-mention'd will be found to be at c; but because the Bottom or Bottom-plate BE is of some Weight, the Center of Gravity will be brought down to C. The Center of Gravity of the full Vessel (which will be a solid truncated Cone) will be at K. If therefore the Axis of Suspension be placed between those two Centers, as at O [Fig. 11.] such a Vessel when empty may be drawn up and down hanging with the Mouth downwards, but come up when full with the Mouth upwards. This is of Use in a Chain of Buckets going \* Plate 7. Fig. 10. 11.

Annotat. round an Axis or Rag-wheel to draw Water from a Depth and deliver it  
 Lect. II. in a Trough above.



Plate 7.  
 Fig. 11.

Plate 7.  
 Fig. 13.

Pl 7.  
 Fig. 14.

Plate 7.  
 Fig. 15.

But if *abcd* be such a Vessel, only with this Difference, that the Bottom is fix'd to the narrow Part *ed*, and the Mouth is at *ab*. The Center of Gravity of the empty Vessel (without considering the Bottom) is at *c*, but by the Weight of the Bottom brought to *C*. The Center of Gravity of the full Vessel will be at *K*. If such a Vessel be suspended between those Centers, as at *O* (Fig. 13.) it will continue with the Mouth upwards when empty; but turn over as soon as it is full. Such a Bucket may be of Use to raise Water by a Machine made with a couple of Buckets fix'd to a Beam which moves upon a Center unequally distant from its Ends, in such manner that a Bucket fix'd to the shorter Brachium raises up a Bucket at the other End, so as to make it empty its Water in a Cistern above — But a short Description and Figure will make the Thing more plain.

*AA*, Are the two Spouts running from a Brook or Spring of Water into the two Buckets *D* and *E*, *D* containing about 30 Gallons, and being call'd the *losing Bucket*, and *E* the *gaining Bucket* containing less than a quarter part of *D*; as for Example, 6 Gallons.

*DE* is a Leaver or Beam moveable about the Axis or Center *C*; which is supported by the Pieces *FF*, between which the Bucket *D* can descend when the contrary Bucket *E* is rais'd up. *DC* is to *CE*, as 1 to 4.

*GL* is an upright Piece, through the Top of which the Leaver *KI* moves about the Center *L*, sometimes resting on the Prop *H*, and sometimes raised from it by the Pressure of the Arm *CE* on the End *I*.

The Bucket *D*, when empty has its Mouth upwards, being suspended as above-mentioned. The End *D* with its Bucket is also lighter than the End with the Bucket *E*, when both are empty. By reason of the different Bore of the Spouts, *D* is fill'd almost as soon as *E*, and immediately preponderating sinks down to *D* (Fig. 15.) and thereby raises the contrary End of the Leaver and its Bucket *E* up to the Cistern *M*, where it discharges its Water, but immediately the Bucket *D* becoming full pours out its Water, and the End of the Leaver *E* comes down again into its horizontal Situation, and striking upon the End *I* of the loaded Leaver *IK* raises the Weight *K*, by which means the Force of its Blow is broke. If the Distance *AB*, or Fall of the Water be about six Foot, this Machine will raise the Water into the Cistern *M* 24 Foot high. Such a Machine is very simple, and may be made in any Proportion according to the Fall of the Water, the Quantity allow'd to be wasted, and the Height to which the Water must be raised.

" Some Years ago a Gentleman shew'd me a Model of such an Engine, varying something from this, but so contriv'd as to stop the running of the Water at *A*, *A*, when the Leaver *DE* began to move.  
 " According to this he told me that he had set up an Engine in *Ireland*  
 " which raised about half a Hoghead of Water in a Minute 40 Foot high,  
 " and did not cost 40 Shillings a Year to keep in Repair; and that it  
 " was not very expensive to set it up at first.

10. [30.— *A Windmill must be supported, &c. and a Crane.*] This is Annotat. not strictly true in Practice; because a Regard is to be had to the Force Le&t. II. with which the Wind pushes the whole Windmill back with part of its Pressure, whilst it turns the Sails with the rest; and therefore the Line of re&tion passing through the Center of Gravity must fall before the Axis of the Post nearer the Sails. Likewise in a Crane (I mean such a one as wholly turns round with the Weight) regard is to be had to the Weight which is to be lifted by it; and the Center of Gravity of the Crane plac'd so much back from the Weight, that the Line of Direction may only pass through the Middle of the Shaft, when the Weight (which brings the Center of Gravity forward) is hanging upon the Crane.

11. [32.— *The Center of Gravity will remain at rest, &c.*] See Sir Isaac Newton's Principia, Coroll. 4th, of the Laws of Motion, 2d Edition: Page 17.

12. [37.— *Demonstrated in the Notes.*] Let the Spindle or double Cone of Plate 4. Fig. 14. be represented here seen endwise (Plate 8. Fig. 1.) A F Plate 4. is one of the rising Rulers upon which the Body is to roll, A G the horizontal Line, B the Vertex of one of the Cones. Let F G the lower part Fig. 14. of the Screw S be made equal to  $e E$ , which is somewhat less than the Semidiameter of the common Base of the two Cones; or (which is the same thing) let E F be another horizontal Line passing a little under B the Axis of the Cones and B F will be the Way of the Center of Gravity of the Body, which Line having a Declivity towards S, the Center of Gravity of the Body must descend and consequently bring the Body along, more or less swiftly, as that Declivity is quicker or slower. Plate 8.

The Cylinder of Figure 15 Plate 4 is made of light Wood, with a small Plate 4. Cylinder of Lead at K going quite through it near the curve Surface and Fig 15. parallel to the Axis of the great Cylinder, to the Intent that the Center of Gravity of the compound Body may be removed from the Axis M into the Line K O; and then the Cylinder must be so laid on the inclin'd Plain A C, that the Center of Gravity of the said Cylinder may descend whilst it is rolling towards R, which will make it go up the Plane till the Center of Gravity is fallen as low as it can: Supposing always a String fix'd to the upper part of the Plane, and going round the Cylinder to keep it from sliding when the Plane is not horizontal, as in the Figure.

As the Length of the Cylinder does no way relate to its rising on, or being supported by, the inclin'd Plane; in considering the Motion of the Cylinder on the Plane differently posited, we shall in the second Figure of Plate 8. only consider the Sections of the Cylinder, Plane, and Horizon. Plate 8. P T A is the Section of the Wooden Cylinder, C A that of the leaden Fig. 2. one, C the Center of Gravity, M the Center of Magnitude, and P Q the Section of the Plane at first supposed Horizontal.

I say, first, If P a be taken upon P Q equal to P T A the half Circumference of the Cylinder, the Point a will be the farthest Place to which the Cylinder will roll. When

Annotat. When the Diameter  $PA$  going thro' the Center of Gravity  $C$  is perpendicular to the horizontal Plane, as in the Figure, the Cylinder will stand still, because the Center of Gravity is directly over the Center of Motion at  $P^*$ , but as soon as  $C$  is ever so little inclin'd towards  $Q$ , the Body will roll till the Point  $A$  comes to  $a$  describing the Semi-Cycloid  $Aa$ , whilst all the Points of the Semicircle apply themselves successively to the Line  $Pa$ , which is the Base of the Cycloid. That the Cylinder will go quite to  $a$ , is evident by observing  $Cc$  the way of the Center of Gravity, which is not in its lowest Place till it is come to  $c$ , and must afterwards rise towards  $\pi$  if the Body roll'd on farther; and therefore if by the Velocity acquir'd the Body should go on towards  $Q$ , the Center of Gravity in going down again from  $\pi$  will bring back the Body to  $a$ ; the Diameter  $PA$  being again perpendicular to the Horizon, but in the inverted Position  $ap$ . *Q. E. D.*

I say secondly, that if the Plane be inclin'd to the Horizon in any Angle whose *right Sine* is  $MC$  the Distance of the Center of Magnitude from the Center of Gravity, the Semidiameter of the Cylinder being *Radius*; the Cylinder laid upon such a Plane will neither ascend nor descend when the Center of Gravity is directly over the Point  $T$ , where the Cylinder touches the Plane, provided it be kept from sliding by a String going under it  $*$ , in the manner represented in the 15 Fig. of Plate 4.

\* L. 2.  
Plate 4.  
Fig. 15.

Turn the Cylinder till the Center of Gravity is at  $K$  in the same horizontal Line with the Center of Magnitude; or (which is the same thing) till the Semidiameter  $MA$  becomes  $M\alpha$ ; from  $K$  drop the Perpendicular  $KT$ , which cuts the Circle at  $T$ , and draw the Radius  $MT$ , to which the Plane  $\pi\pi$  being made perpendicular, you will have the Angle  $\pi DQ$  made by the Plane with the Horizon, equal to the Angle  $MTK$ , whose *Sine* is  $MK$  equal to  $MC$ . For by producing  $M\alpha$  to  $L$  it is evident (by 8. 6. of *Euc.*) that the Angle  $MTK = KLT$ , but (by 29. 1. *Euc.*)  $KLT = LDQ$ .


In this Situation it is evident by Construction, that the Center of Gravity cannot descend, because the Line of Direction is supported at  $T^*$ , where the Plane touches the Cylinder.

\* No. 45.

For if the Body was roll'd up any higher on the Plane to bring  $K$  towards  $\pi$ ,  $T$  the touching Point would advance faster towards  $\pi$  than  $K$  does, and therefore the Line of Direction would cut the Plane below  $T$  towards  $D$ , so that the Center of Gravity would descend and bring back the Cylinder to bear on  $T$ . The other way the Body would roll down upon moving  $K$  ever so little towards  $D$ , the Line of Direction then advancing faster towards  $D$  than the Point of Contact  $T$ . This may be made plainer, by considering the Cylinder as a Balance; as for Example, if  $Mw$  be a Balance, sustaining on its End  $M$  a Weight, equal to the Weight of the Cylinder, without the Lead, and on the other End  $w$ , a Weight equal to the Excess of Weight of the Lead, above the Bulk of Wood whose Room it takes up. Let  $K$  be their common Center of Gravity, found as has been taught  $*$ : Then considering  $KT$  as an inflexible perpendicular Prop, sustaining

\* 38.

taining the Balance at K, the Balance will continue in *Equilibrio*, whilst the Prop is supported by bearing on T, the Place where the Plane touches the Cylinder. If the Plane should make a greater Angle with the Horizon, the Point T being remov'd farther towards L, it would be the same, as if the Prop should endeavour to support the Balance between K and *w*, in which Case the Weight M wou'd preponderate, and carry the whole Cylinder towards P; but if the Plane makes a less Angle with the Horizon, T will go towards D, and the Balance then being propp'd between K and M, the Weight at *w* will preponderate and carry the Cylinder towards L. Q. E. D.

Annotat.  
Left. II.  


C O R O L L A R Y.

Hence follows also, that there must be an Angle  $dDQ$  less than  $\pi DQ$  which will be the Inclination of the Plane, on which the Cylinder above-mention'd can roll to its greatest Height on the Plane. For if the Angle  $\pi DQ$  be diminish'd a little, the Cylinder will roll towards  $\pi$  going upwards; and if the Plane  $DQ$  be rais'd a little, so as to make a small Angle with the Horizon, the Cylinder will rise on that Plane, tho' it will not roll so far upon it as in the horizontal Situation. If this Angle be encreased, the Cylinder will rise higher, as its Way on the Plane shortens, but not beyond a certain number of Degrees of Elevation, at which Elevation the Cylinder will go no higher above the Horizon, tho' its Way measured on the Plane will continually shorten till it is contracted to a Point when the Inclination is in the Angle  $\pi DQ$ . Likewise as the Angle  $\pi DQ$  (by lowering the Plane  $D\pi$  round the Center) diminishes, the Cylinder will go higher as it rolls farther on the Plane, till the Angle is diminished to a certain number of Degrees, after which the Cylinder will rise less; but its Way measured on the Plane will encrease, till the Plane comes to be horizontal. Therefore there is an Angle of the Plane  $dDQ$ , which is a *Maximum*, as to the Rise of the Cylinder on the Plane.

I say, *thirdly*, that the Inclination of the Plane being given on which Plate 8. the Cylinder will rise to the greatest Height (or any other Inclination of a Fig. 3. Plane on which it can rise at all) the Length which the Cylinder will go on the Plane will be equal to the Length  $Tv = TV$ , which is equal to half the Circumference *minus* the Arc  $AV$ , which Arc  $AV$  contains twice the number of Degrees of the Plane's Inclination, together with the Degrees of twice the Difference of the Angle at the Center of two rectangular Triangles which have  $MN$  the Sine of Inclination for Radius, but their Secants are  $MR$  the Semidiameter of the Cylinder, and  $MC$  the Distance of the Center of Gravity from  $M$  the Center of the Cylinder. Moreover,  $Tv$  being the Length of the Cylinder's Progression on the Plane, the Height of the Plane at  $v$ , *viz.* the Perpendicular  $vZ$ , will be the Cylinder's Rise above the Horizontal Line.

Annotat.  
Lect. II.

P R E P A R A T I O N.

$PQ$  is the Horizon and  $Pq$  the Plane of given Inclination. Since  $Pq$  cuts the Circle  $PVA$ , the Cylinder cannot rise upon it, therefore we must take another Plane parallel to it. The Diameter  $PG$  being drawn perpendicular to the Horizon, and  $RT$  thro' the Center of Gravity  $C$  or  $S$  parallel to  $PG$ , draw the Diameter  $TA$ , making the Angle  $PMT$  equal to the Inclination of the Plane, and draw  $\pi\pi$  perpendicular to that Diameter at  $T$ , and  $\pi\pi$  will be a Tangent Plane parallel to the former; (for  $PMT$  being equal to  $MTR$ , because of the Parallels  $PG$ ,  $RT$ , and the common Angle  $RTv$  being taken away from the two right Angles  $MTv$  and  $RTZ$ ,  $vTZ$  the Angle of Inclination of the new Plane  $\pi\pi$  will appear equal to the given Angle  $PMT$ ) draw  $MR$ ; thro'  $C$  draw  $Mf$ ; make the Angle  $XCM$  equal to  $MCT$ , produce  $XC$  to  $V$  and draw  $MV$ . Because  $XCv$  is one straight Line and at the same Distance from the Center  $M$  as  $RT$ , therefore  $CR$  and  $Cv$  are equal (by 7. 3. of *Eucl.*) since the Triangles  $CMd$ ,  $CMe$  are equal (by 6. 7. 8. 1. *Eucl.*) the Angle  $RMV$  is bisected. Draw  $MN$  the Sine of the Angle of Inclination, and the Degrees of the Arc  $Rv$ , or the two Differences mention'd will be found by comparing together the Angle of Inclination and the Triangles  $RMN$  and  $CMN$ .

The Cylinder being laid on the Plane in the Position describ'd in the Figure, will not only remain at rest, if the Center of Gravity be at  $S$ , but will return, to that Position when moved out of it towards  $\pi$  or towards  $v$ , because in each of those Cases the Center of Gravity must rise, and therefore  $ST$  must be the Distance of the Center of Gravity from the Plane, measured upon the Line of Direction of the Center of Gravity when nearest to the Plane, over that part of the Plane, where the Cylinder rolling upwards or downwards will stop its Motion. Now if the Center of Gravity be brought to  $C$ , it is evident that  $CR$  will be equal to  $SR$ , and that  $Cv$  will also be equal to it, because by Construction it is equally distant from  $Mf$  a Line going thro' the Center, and consequently  $= CR$ ; also that no other Point of the Circumference will be equally distant from  $C$ . Then if the Center of Gravity from the Point  $C$  in the Line  $RT$ , where it is exactly over the Point of Contact, be ever so little moved towards  $v$  by pushing the Cylinder that way, the Cylinder will roll on the Plane, and if it advances the Length of the Semicircumference till the Diameter  $TA$  is inverted and becomes  $at$ , the Center of Gravity will be got to  $c$  the Chord  $TR$  being now  $tr$ ; but since  $r$  does not touch the Plane, the Center of Gravity must descend again and bring back the Cylinder till  $u$  (which was the Point  $V$ ) returns on the Plane to  $v$ , where  $\alpha Kv$  will be the Chord which was mark'd  $XCv$  in the first Position of the Cylinder,  $K$  being the the Center of Gravity and  $Kv$  which is equal to  $Cv$  being in the Situation of  $ST$ , and the whole Cylinder bearing on  $v$  just as it did on  $T$ , when we suppos'd  $S$  the Center of Gravity,  $TA$  and  $TR$  being turn'd into  $T\alpha$  and

$T\rho$ .

**T**e. Therefore the Distance  $Tv$  on the Plane will not be equal to the Semicircumference, but want of it the Arc  $au = \alpha v = AV$ . Consequently the Length run on the Plane by the Cylinder will be equal to the Arc  $TV$ , which may be found by a Thread applied to the Cylinder from  $T$  to  $V$ . Annotat. Lect. II.

Now the Number of Degrees of this Arc  $TV$  may be found in the following manner. Plate 8. Fig. 3.

The Angle of Inclination is  $\angle TZ = MTN = MRN = TMP = AMG = GMR$ .

In the rectangular Triangles  $MRN$  and  $MCN$ ,  $NM$  the Sine of Inclination of the Plane is the Radius common to both;  $MR$  the Semidiameter of the Cylinder is the Secant in the Triangle  $MRN$ , and  $MC$  the Distance of the Center of Gravity from the Center of the Cylinder is the Secant in the Triangle  $MCN$ . The Angle  $RMN = CMN = RMC$ ; therefore  $RMV = 2 RMC$ , which may be found by Trigonometrical Tables, and consequently it is known. But the Angle of Inclination being given, its double is given, therefore  $AV$  and  $TV$  are known. *Q. E. I.*

To apply what has been said, let us suppose the Center of Gravity of the Cylinder to be distant from its Center of Magnitude  $\frac{2}{3}$  of the Radius, that is to be at the Point  $C$  in the Figure \*.

\* Plate 8.

The Angle  $plg$  of the Plane, on which the Cylinder cannot rise is thus found, by Analogy. Fig. 3.

As  $Mb$  is to  $Mk :: (3 : 2 ::)$  So is the whole Sine: To the Sine of the Angle  $Mlk = lMP = plg = 41^\circ. 48' +$ , &c.

The Maximum of the Angle of the Plane on which it will rise most is in this Case found to be  $26^\circ$ , and the Height of that Rise is  $vZ = 42, 85$  such Parts of which the Circumference of the Cylinder contains 360.

**N. B.** Hence it appears that the Maximum is not in the middle of the Angle  $plg$ , or at  $20^\circ. 54'$ , as one might at first imagine.

Now let us suppose the Angle of Inclination of the Plane to be  $15^\circ$ ; the Length which the Cylinder will roll on the Plane, and the Height to which it will go above the Plane's Base will be found not only in the manner abovemention'd; but also two other Ways, which for Variety-sake I set down here.

## P R E P A R A T I O N.

The Center of Gravity being in the Point  $C$ , at  $\frac{2}{3}$  of the Radius of the Cylinder  $MC$  from its Center  $M$ ; making the Arc  $fV = Rf$ , and drawing thro' the Center of Gravity  $C$  the Line  $VCX$ , the Segment  $VbX$  is = to the Segment  $RzT$ , and consequently when the Cylinder has trac'd on the Plane  $Tv$  the Arc  $TzV$ , it will rest and be *equipois'd* (on the Plane) upon the Point  $V$  for the same Reason that it is so upon the Point  $T$  before its rolling.

As the Angle of the Plane  $\angle TZ$  is  $= 15^\circ$ , that Angle is equal to the Angle  $PMT = bMR$ ; therefore there only remains the Angle  $RMV$   
M
to

Annotat. to be found, or its Half  $RMf$ ; and the two following Ways give the same  
 Lect. II. Value for it.



1. Since the Sine  $GR$  for the Angle  $\angle MR$ , when the Radius is the Radius of the Cylinder, is equal to  $yC$ , the Sine of the Angle  $\angle Mf$  for the Radius of  $oCk$ , the Circle of the Center of Gravity: I say,

As the Radius of the Center of Gravity:

To the Radius of the Cylinder :: (or as 2 to 3.)

So is  $no = 2588190$ , the Sine of  $15^\circ$  for the Radius of the Center of Gravity:

To  $3882285 = GR$  or  $yC$ , the Sine of the Angle  $\angle Mf$ , which is found to be  $22^\circ. 50' +$ .

From which therefore subtracting the Angle  $\angle bMR = 15^\circ$ , there will remain  $7^\circ. 50'$  for the Angle  $RMf = fMV$ . Therefore the whole Angle  $RMV$  is  $= 15^\circ. 40' +$ ; to which adding  $30^\circ$  for the two equal Angles  $\angle bMR$ ,  $\angle PMT$  each of  $15^\circ$ , the whole makes  $45^\circ. 40' +$  to be subtracted from  $180$ , equal to half the Circumference  $bV \approx TP$  of the Cylinder, and there remains the Arc  $VT$  for the Length run by the Cylinder rolling on the Plane  $Tv$ .

2. The other Way to find the Angle  $RMf$  is thus,

In the Triangle  $RM C$  are given  $RM$ ,  $MC$ , and the Angle  $MRC = RMb = 15^\circ$ ; therefore I say,

As the Radius of the Center of Gravity:

To the Radius of the Cylinder :: (or as 2 to 3, or  $MC:MR::$ )

So is the Sine of the Angle  $MRC$  of  $15^\circ = 2588190$ :

To  $3882285$  the Sine of the Angle  $RCM$ , or  $RCf = 22^\circ. 50' +$ .

And that Angle  $RCf$  being equal to the two opposite Angles  $MRC$ ,  $MCb$ , or  $RCf$ , if from  $22^\circ. 50'$  be subtracted  $15^\circ$  for the Angle  $MRC = RMb$ , there remains the Angle sought  $RMf = 7^\circ. 50' +$ , as before.

Having therefore subtracted  $45^\circ. 40' +$  from  $180^\circ$ , there will remain for the Arc  $VZT$   $134^\circ. 20'$  for the Way of the Cylinder on the Plain  $Tv$ , that is,  $134\frac{1}{3}$  of such Parts of which  $360$  make the Circumference of the Cylinder, by which one may find the Height  $vZ$  to which it rises on the Plane, thus,

As the whole Sine:

To  $134\frac{1}{3} ::$

So is the Sine of the Angle  $vTZ$ .

To  $vZ = 34, 77 -$ . *Q. E. I.*

\* See the last

Edition of

Sir Isaac

Newton's

Principia B.

3. Prop. 12.

13. [39 — Common Center of Gravity may be found, &c.] By this Means may be found the Center of Gravity of our System, in any Position of the Planets \*. Let us first suppose them all on one Side and in a Line as  $SCA$ , going thro' the Sun's Center  $C$ : Or to speak like an Astronomer,

let

let all the superiour Planets be in Opposition, and the inferiour ones in Annotat. their inferiour Conjunction. *First*, the common Center of Gravity of *Mer-Lect. II.* *cury* and the Sun will be very near the Sun's Center; because the Quantity of Matter in the Sun is more than a Million of times greater than in *Mercury*, and *Mercury* is not 825 Semidiameters of the Sun distant from the Sun's Center. *Secondly*, *Venus* (supposing it, as very probable, to have about the same Quantity of Matter as the Earth) taken into Consideration, will bring the common Center of Gravity of the three Bodies but a little forwarder, viz. to  $\varphi$ , because its Mass is to the Sun's but as about 1 to 169282, and its Distance from the Sun's Center but about 145 Semidiameters of the Sun. *Thirdly*, The common Center of Gravity of the Earth, and the three Bodies abovemention'd, will be brought but to  $\Theta$ , or a little forwarder. *Fourthly*, The common Center of Gravity of *Mars* and the other four Bodies will be brought still a little nearer to the Surface of the Sun, as to  $\delta$ , but not half way from the Sun's Center to its Surface. *Fifthly*, The Quantity of Matter in *Jupiter* being to the Quantity of Matter in the Sun, as 1 to 1067; and *Jupiter's* Distance from the Sun compar'd to the Sun's Semidiameter in a Ratio something greater; the common Center of Gravity of *Jupiter* and the Sun will be a little way without the Sun's Surface; and therefore the common Center of Gravity of *Jupiter* and the other five Bodies will come to  $\psi$ , a little farther out. *Lastly*, As the Matter in *Saturn* is to the Matter in the Sun as 1 to 3021, and the Distance of *Saturn* from the Sun is to the Semidiameter of the Sun in a Ratio something less; their common Center of Gravity, without the other six Bodies, would be in a Point as  $h$ , a little within the Sun's Surface, and therefore the common Center of Gravity of all the seven Bodies will be at  $I$ , still a little farther out from the Surface of the Sun, but hardly a whole Diameter distant from the Center of the Sun. When *Jupiter* and *Saturn* are on different Sides of the Sun, this common Center of Gravity will always be within the Body of the Sun, let the other Planets be in any Position, because of their Nearness, and the small Quantity of Matter which they contain. It is this common Center of Gravity of our System which is at rest, and not the Center of the Sun; for the Sun has a kind of wabbling Motion about this Center. *The little Difference made by the Comets and Satellites of the primary Planets is not worth mentioning here.*

14. [43 — *Line of Direction falls within their Base.*] The Tower of *Pisa* is a round Tower 138 Foot high, whose Top over-hangs the Base 15 Foot, as represented in the fifth Figure: And the Tower of *Bolonia* is square, 130 Foot high, its Top over-hanging the Base only nine Foot. See the sixth Figure. Plate 8. Fig. 5 and 6.

Animal Motions are always subject to these Rules, which we observe without thinking on them. When we stand upright with our Feet, as represented in *Fig. 7.* the Line of Direction goes thro' the Point  $C$ , and passes between our Feet at  $D$ , and we may move our Heads from  $F$  to  $E$  or  $G$ , and our Bodies forwards, backwards, or side-wise, as far as  $I$  or  $H$ , Plate 8. Fig. 7.

Annotat.  
Lect. II.


  
\* Plate 8.  
Fig. 10.

Plate 8.  
Fig. 7.  
and 9.  
\* Plate 8.  
Fig. 9.

\* Plate 8.  
Fig. 8.

without danger of falling or stirring our Feet, as long as the Line of Direction traverses no farther than I A or H B, and falls any where within the Space A B; which in this Situation of our Feet makes a pretty large Base. But if we set one Foot before the other, as in *Fig. 10* \*, a little Push side-wise will make the Line of Direction (which went thro' C) fall out of the Base to the right or left, towards E or B; in which Case a Man must fall, if he does not quickly remove his Feet to the Position of *Fig. 7* or *9*. When we stand upon either Leg we must bring our Body so much over the Foot A B or D E \*, that the Center of Gravity being directly over it, the Line of Direction may go thro' c or K; and in walking, the Line of Direction must travel thro' every Place where each Foot is set down; going successively thro' the Points E, A, D, B, whilst the Center of Gravity goes thro' the Points G, C, F, &c. so that unless a Man, in walking streight forward, sets one Foot directly before the other, the Line of Direction will not describe a streight Line upon the Plane where the Man walks, but an indented Line, that is Angles, to the right and left, whilst the Body of the Man goes on in a wadling Motion. This we see in the walking of fat People, and all others that straddle in their Gate \*. The Line of Direction

\* It is not strictly true that any Man in his common Walk sets one Foot so exactly before the other, as to carry on the Bottom of his Line of Direction in a streight Line, as represented in *Fig. 10* and *11*. Because if a streight Line be drawn with Chalk, it is difficult to walk streight along it; but the plainest Proof is the Observation of two upright Sticks, of about the Height of a Man, the one painted white and the other black, and set up about ten Yards beyond one another, in the same Line that a Man walks towards them; for in such a Case, tho' he keep one Eye shut, the last Stick will appear sometimes on the right and sometimes on the left of the First; and the more so, the nearer the Man comes to the Sticks. Rope-dancers\*, indeed, go in a streight Line; but it is what they have learn'd by Art, and inur'd themselves to by long Practice; yet they must even after all have Helps to keep their Center of Gravity over the Rope. They generally fix their Eyes on some distant Point in the same Plane as the Rope. They have commonly a long Pole loaded at the Ends with the Balls of Lead B, b, by the Motion of which they can alter the Position of the common Center of Gravity of their Body and the Pole; as for Example, the Center of Gravity of the Rope-dancer C A being at A, his Line of Direction should go thro' a, off from the Rope; but by moving the Pole towards B,

\* Plate 8.  
Fig. 13.

the common Center of Gravity of the Man and Pole is brought to C, in which Case the Line of Direction C D goes thro' the Rope. Those who are well skill'd in this Art will sometimes use their Arms only, instead of a Pole; and it is very common for several of them to dance with a Flag, with which they strike the Air the same way that the Center of Gravity goes when the Line of Direction does not go thro' the Rope; and, by the Reaction of the Air, the Center of Gravity is brought back to its proper Place.

*Those that would enquire farther into this Matter, may consult J. A. Borelli, in his Book De Motu Animalium, Chap. 18, 19, 20 and 21. In the said 21st Chapter he gives an Account of the Motion of an Horse, part of which (as it is very curious) I shall repeat here.*

The Ancients observing that Horses and other Quadrupeds, in Galloping, lift up their two fore Feet, and then their hind Feet, as soon as the fore Feet are set down, did imagine, that in Walking, as well as Pacing and Trotting, an Horse has two Feet off of the Ground at one time; and accordingly in their Bras or Marble Statues, they have represented their Horses with two Legs off of the Ground diagonally opposite, as the right before and left behind, or left before and right behind. The modern Statuaries have also fall'n into the same Error, because in the quick walking of an Horse the Eye cannot well distinguish,

rection going thro' the Points A, B, C, D, E, describes a streight Line in Annotat. Fig. 11, where the Feet are set before one another; but when the Motion of one Foot is in a parallel Line with the Motion of the other, an indented Line is describ'd by the Center of Gravity above, and the Line of Direction as it cuts the Ground at A, B, C, D, E, Fig. 12. Ducks, Geese, and the greatest part of the Water-fowl, whose Legs are set wide asunder for the Conveniency of their swimming, and turning quick in the Water, have always a wadling Motion upon Land; but a Cock, a Stork, an Ostridge, and most other Birds that are not web-footed, walk almost directly forward, without wadling (especially when they walk slow) having their Legs so plac'd as to put one Foot before the other with greater Ease. Thus Quadrupeds seldom or never waddle, because they have commonly three Feet upon the Ground at a time: So that however the Base receiving the Line of Direction alters from a quadrangular to a triangular Figure; that Part of it, in which the Line of Direction falls, is always in, or near the same Line.

When a Man stands in a firm Posture \*, A B, the Distance of his Feet, is the length of a quadrilateral Figure, whose Breadth is nearly the Length of the Feet, and D is the Point under the Center of Gravity C, where the Line of Direction falls. Let the Lines A C and B C be drawn, then let those two Lines and D C be continued to the Points E F G, so as to make the Triangles E C G and A C B equal and similar. As long as the Line F D (or a Plane going thro' it) cuts the whole Body of the Man into two equal Parts, the Center of Gravity will be at C, and C D will be the Line of Direction. But if the Body be inclin'd towards the left Hand H, the Center of Gravity will move from C to H, the Line of Direction will become H B, and the right Foot being easily remov'd from A may be carried on beyond B, by which Means the Man will go on towards the left:

distinguish, and therefore he has shewn from Mechanical Principles, that the Motion of raising two Feet at once in Walking cannot be consistent with the Wisdom and Simplicity of Nature. To Borelli therefore I refer the Reader, who wou'd know what the Motion is not, and only copy from him, here, what the Motion is.

\* Plate 8. Let us consider an Horse \*  
Fig. 14. as an oblong Machine sustain'd by the four Legs, as four Props or Columns, resting on the Points A B C D, which make a rectangular quadrilateral Figure; then the Line of Direction will fall perpendicularly on E, a Point in or near the Center of the quadrilateral Figure, which will make the Station or Standing of the Horse the most firm. The progressive Motion begins by one of the hind Feet, as for Example, the left hind Foot C, which by strongly pressing back the Ground moves forward the Center of Gravity, and conse-

quently carries on the Line of Direction from E to G, as it self moves from C to F; this done, immediately the Foot B is rais'd, and carried forward as far as H, which Motion of the Foot is easy, because the Line of Direction first falls within the Triangle A B D, secondly within the Trapezium A B F D; that is, the Body of the Horse is sustain'd by three or by four Columns. Lastly, the three Feet A, D, F remaining firm, and taking in the Line of Direction at G, immediately the left fore Foot B is carried forward to H; and, by the Impulse already made, the Center of Gravity is also carried over I, namely the central Point of the Rhomb A H F D. The Motion of the two left Feet being completed, the Impulse and Motion of the right hind Foot D begins, and then that of the right fore Foot; and so on in the Manner above describ'd, as the Animal moves forward.

In

Annotat.  
Lect. II.

Plate 8.  
Fig. 8.

In like Manner by inclining towards I the Line of Direction will be removed to IA, and the Man go to the right. When a Man stands upon one Foot, it is with some Difficulty. For Example, let the Line of Direction be CD; by the Motion of the Blood, and Lungs, and other Animal Motions, the Center of Gravity will be apt to vacillate or totter towards F or G on either Side about the Center of Motion D, where now the Base is but small. If the Line of Direction comes to B the Man must fall forwards, backwards if to E; and tho' A be under the Heel of the Foot, yet in the Motion of the said Line of Direction from D to A the Body will be apt to go too far towards E, and so bring the Line of Direction beyond the Base. This will more probably happen in the side Motion of the Body; so that the Body will be in danger of falling, unless the right Foot be put down towards that Side where the Body inclines. Birds stand upon one Foot much more easily than Men, because their Line of Direction being much shorter, and the Base of one Foot a large Rhomboidical Figure made by the four Claws, the Line of Direction cannot go out of that Base, unless the Center of Gravity rises, which is impossible without a violent Motion \*.

\* 28, 43.

When a Porter carries a Burthen upon his Shoulders, he must stoop, because if he should stand upright, the common Center of Gravity of the Man and Burthen would be so far brought back that the Line of Direction would fall behind the Feet. For the same Reason, when a Woman with Child is very near her Time, she bends backwards as she goes, by Reason of the Burthen before, which otherwise would cause her to fall forwards.

Plate 9.  
Fig. 1.

15. [49. — *Roll along the Plane.*] If the Ball \* FE be laid upon a smooth horizontal Plane AB, it will stand still, tho' it touches but in one Point as O, because the Line of Direction CO goes thro' the said Point; but if the Plane be ever so little inclin'd to the Horizon, as in the Position CD, the Ball will continually roll forward towards D, because then the Line of Direction will always fall before the touching Point.

Plate 9.  
Fig. 2 and 3.

If such a solid Body as G, contain'd under twelve rectangular Parallelograms, and two opposite, parallel and equal Dodecagons, be laid upon the inclin'd Plane BAC, as it will slide from A to C, the Center of Gravity moving in the Line eg parallel to the Plane DC, and *io* one of the parallelogram Planes of the Body always touching the said Plane AC. But if the Plane be more inclin'd, as in the Position DE, it will appear, by drawing the Arc ef with the Distance *ie* about the Point *i* (the only touching Place of the Body then) that the Center of Gravity can descend; and when the Point *o* is applied to the Plane DC, the Line of Direction will fall beyond the said Point *o* towards E; and therefore the Body will roll or tumble towards E. In the same Manner all the angular Points, or rather the Edges of the Plane Surfaces, will successively apply themselves to the Plane DE till the Body has roll'd quite down.

16. [49 — *In the Load of Iron the Center of Gravity is low; but very high in the Load of Hay.* Annotat. Lect. II.

K \* is the common Center of Gravity of the Load of Hay and the Carriage P M, whose Line of Direction is K E, whilst the Plane P M (on which the Carriage is drawn) is horizontal; but if C D be an horizontal Line, the Plane P M will be inclin'd to the Horizon in the Angle B P D, and the Line of Direction being chang'd from K E to K P (because K P is the only Perpendicular from K to C D) will fall out of the Base Q M towards C, and consequently the Waggon will be thrown over that way; which also appears by drawing round the touching Point Q with the Distance Q K, the Arc K R shewing the Way of the Center of Gravity, which can in this Case descend without first rising. For the same Reason, if the Carriage was drawn along an Horizontal Plane, whose Section is represented by C D or P N, and the Wheel M should meet with a Rub of the Height of N M, the Load of Hay would also be overthrown upon that Account. Plate 9. Fig. 4.

But as a Load of Iron \* lies much lower upon its Carriage, the Center of Gravity must also be lower, and therefore the Line of Direction will fall within the Base on the same Inclination of the supporting Plane, as would occasion it to fall out of the Base in a Load of Hay, as is evident from the Figure. Let C D also here represent an Horizontal Line, and Q M the Road or the Base supporting the Carriage, the Angle of Inclination M Q N in this Figure being equal to M Q N in Fig. 4. The Line of Direction K P will here fall within the Base Q M, and cannot fall out of it till the Angle of Inclination is encreas'd to B Q Y, by making X Y the Horizontal Line; or, which is the same Thing, unless the Wheel M meets with a Rub of the Height of y M much higher than the Rub which would overturn the Load of Hay. Plate 9. Fig. 5.

---



---

## LECTURE III.

Lect. III. 1. *Simple Machines* or *Organs*, call'd by some *Mechanical*  
*Faculties*, or *Mechanical Powers* \*, are such Instruments  
 \* Annot. 1. as are of one Piece (or consider'd as such) by means of  
 \* L. 2, 19, which the \* *Powers* describ'd in the last Lecture act upon  
 20. *Weights*, in order to give or stop Motion; to overcome, make  
 or stop Resistance.

2. All Engines (however compounded) for the Uses of Life, are made up of various Combinations of the simple Machines. Sometimes all of them may be found in one Engine; sometimes two or three of them; and sometimes only one of the Mechanical Powers multiplied.

3. The *Simple Machines* are the seven following, viz. the *Balance*, the *Leaver*, the *Pulley*, the *Axis in Peritrochio* (or Axle in the Wheel) the *Inclin'd Plane*, the *Wedge*, and the *Screw*. N.B. *Authors differ in their Enumeration of the Mechanical Powers, some making them to be Six, excluding the Inclin'd Plane from those which I have nam'd. Others have reduced them to Five, considering that the Screw is only a Wedge carried round a Cylinder: And others again have made the Leaver and Balance to be the same Power, from their near Resemblance. But since the same Principle is only applied differently (as may be shewn by reducing all the Mechanical Powers to the Leaver, or explaining all their Operations by that of the Leaver \*) and we are to give an Account of the Instruments contriv'd for that different Application, the Mechanical Powers will appear to be Seven.*

\* Annot. 2.

4. Before we explain these Powers or Organs severally, there are some general Things to be consider'd relating to all Engines, which will facilitate our Calculations concerning them, and render the Execution of them in Practice, as perfect as the Nature of the Materials of which they are made will allow.

S U P.

S U P P O S I T I O N S.

5. THO' the Earth be spherical, yet we are to suppose it flat, when we consider Mechanical Engines; because the largest Machine covers so small a Part of the Earth's Surface, that to allow it any sensible Roundness in so small a Compass, would be to allow too much. Lect. III.


6. THO' none of the Bodies which we handle are perfectly hard, or truly of the Figure which we intend to give them; yet we are to suppose every Thing perfect in all our Engines; as for Example, that straight Bodies, as the *Beam of a Balance*, a *Leaver*, &c. are Mathematical *Rigids*, and without Thickness, or Lines altogether inflexible; that the *Mechanical Powers* (whether Simple or in Composition in complex Engines) are without Weight, whatever Materials they are made of; that Bodies are perfectly hard and smooth; that the Parts of Engines move one another without Friction; that Cords are extremely pliable; that the Center Pins of Pullies, or Axes of Motion of them, of Balances, Leavers, Axes in *Peritrochio*, &c. are only Mathematical Lines.

7. THO' the \* Lines of Direction of all heavy Bodies tend \* L. 2. 22. towards the Center of the Earth, and consequently converge together in a small Angle; yet we must consider them as parallel, because they are so, as to Sense; their Point of Convergence being near 4000 Miles off. For this Reason the Walls of a Building, when exactly adapted to the Plumb Rule, are nearer at Bottom, than at Top; tho' in Practice we must always look upon them as parallel.

8. NOTWITHSTANDING the Falsity of the Suppositions, we shall not be led into any Error by them; because, by a second Consideration, we are to have regard to the Imperfection of Engines and Materials, and the Quantity of *Stickage* or Friction; which differ according to the Number and Combination of Parts and Nature of the Materials, of which the several Engines consist: And having made use of the best Methods we can to discover the Imperfections abovemention'd, in each particular Machine; we are to take care to allow

N

enough

Lect. III. enough to be deducted from the Calculation made concerning  
 an Engine suppos'd Mathematically true.

N. B. *Several Methods of finding the Quantity of Friction in Engines will be consider'd hereafter* \*.  
 \* Ann. 3. and L. 4.

### DEFINITIONS.

9. WHEN equal or unequal Quantities of Matter are so applied to a Mechanical Organ, or Engine, that their *Momenta*, or Quantities of Motion, or moving Forces, destroy one another; they are said to be in *Æquilibrio* \*.  
 \* L. 2. 12.

10. WHEN Powers, whose Intensities are equal or unequal, are so applied to an Engine as to destroy each other's Actions, they are also said to be in *Æquilibrio* \*.  
 \* L. 2. 12.

11. ALL Bodies that act against each other by means of Engines, may be consider'd as *Powers* and *Weights*, already described in Lect. 2. N°. 18, 19.

12. BODIES in *Æquilibrio* are said to *æquiponderate*.

13. WHEN Powers and Weights have their Velocities reciprocally as their Masses, or as their Intensities \*, their *Momenta* are equal; therefore they are in *Æquilibrio* †.  
 \* L. 2. 20. † 9.


14. IF the *Momentum* of a Power be greater than that of the Weight, or (on the contrary) the *Momentum* of the Weight be greater; that Power, or Weight is said to *præponderate*, or over-power.

15. WHEN either the *Velocity* (the Mass being equal) or the *Mass* or *Intensity* of a Weight or Power (the Velocity being equal) or both Mass and Velocity together are greater in a Power or Weight than in the opposite (or counter-acting) Weight or Power; the *Momentum* of the former will be greater than that of the latter \*.  
 \* L. 2. 15.

17. N. B. *This will happen, let the Difference be ever so small; tho' then the Resistance made by the Friction will hinder the Effect from being visible.*

Of

## Of the B A L A N C E.

16. THE essential Parts of a Balance are (1.) the Beam, as Lect. III. AB \*; (2.) the Axis of Motion, consider'd only as a Point or  Center of Motion C, which divides the Beam into two Parts; \* Plate 9. Fig. 6 and 7. (3.) those Parts call'd the Arms or *Brachia*, as AC and CB, which are either equal, as in Fig. 6, or unequal, as in Fig. 7. (4.) the Points of Suspension, as \* A, B, in the 6th, and A, \* Plate 9. Fig. 6. B, K,  $\kappa$ , in the 7th Figure. Fig. 7.

17. WHEN Weights hang freely from the Points of Suspension, they gravitate neither more nor less for hanging nearer to or farther from the said Points.


### EXPERIMENT I. Pl. 9. Fig. 6.

18. LET the Weight Q, with its Rope QA hanging at the End A of the Balance AB, be equal to the Weight P, together with its Rope DB. Hang the Weight P at any of the Loops G, F, E, D, of its Rope, and it will at any of them make an *Æquilibrium* with the opposite and equidistant Weight Q.

19. WHEN the Beam of the Balance is equally divided by the Center of Motion (as in the 6th Figure) with Scales hanging freely from the Points A, B, instead of the Weights Q and P, it is the *Libra*, or common *Pair of Scales*.

20. THIS Instrument serves to compare together Bodies which have equal Quantities of Matter, tho' sometimes differing in Bulk; for when the Commodities to be bought or sold are placed in one Scale, so as to keep the Weights in the opposite Scale in *Æquilibrio*, the *Momenta* are equal; and since the Velocities are equal on account of the equal Distances AC and CB \*, the Quantities of Matter must be likewise equal; and this \* 12. is shewn by the horizontal Position of the Beam hanging freely on its Center of Motion, which is plac'd a little above its Center of Gravity. See the 4th Annotation. Ann. 4.

HENCE follows, that the Distance (that is the acting Distance) of any Weight is not to be measured from the Center of Motion of the Balance to the Center of Gravity of the Weight;

Leſt. III.  Weight; and therefore the Lines  $Cn$ ,  $Cm$ , do not expreſs the Diſtances of the Weights  $P$  and  $Q$ ; but their Diſtances are properly  $CB$  and  $CA$ , the leaſt Diſtance from their Lines of Direction  $Nn$  and  $Mm$  to the Center of Motion  $C$ : And therefore when the Weights hang freely upon an Horizontal Balance, the Diſtance of their Points of Suſpenſion from  $C$  may be call'd their Diſtance and meaſur'd upon the Beam; but if the Balance be in an inclin'd Poſition as  $ab$ , not  $bC$  and  $aC$ , but  $dC$  and  $eC$  will be the Diſtances of the Weights  $p$  and  $q$ , being Lines perpendicular to their Lines of Direction  $po$ ,  $qa$ , and going thro' the Center of Motion, conſequently the leaſt Diſtances from it to the ſaid Lines of Direction.

\*Pl. 9. Fig. 7. 21. THE Balance, whoſe *Brachia* are unequal \*, as  $AB$  (Fig. 7.) is the *Roman Statera*, or our Steel-yard; or the little Inſtrument which the *Chineſe* (who take all their Money by Weight) always carry about 'em, call'd the *Dotchins*. See Num. 13. of Leſt. 2. Plate 4. Fig. 3. This Inſtrument ſerves to compare together, at one Operation, Bodies that have equal or unequal Quantities of Matter, but in weighing heavy Goods, it is not ſo exact as the Scales.

\*Pl. 9. Fig. 8. THE \* Balance of Fig. 8. may ſerve as a Steel-yard or as a Pair of Scales, on account of the ſeveral Diviſions on each *Brachium*.

### T H E O R E M.

22. *One or more Weights ſuſpended on one Brachium of a Balance, will be in Æquilibrio with one or more Weights ſuſpended on the other Brachium; provided that the Sum of the Momenta (or the whole Quantity of Motion) of the Weights on one Side of the Center of Motion, be equal to the Momenta on the other Side of it.*

### EXPERIMENT II. Pl. 9. Fig. 8.

23. UPON the *Brachium*  $AC$  hang three ſingle Pounds at the Diviſion 8, a ſix Pound Weight at Number 5, a three Pound Weight at Number 1, and a nine Pound Weight at Number 3. Then on the *Brachium*  $CB$ , hang a two Pound Weight  
at

at 2, a twelve Pound Weight at 5, and a two Pound Weight at Le&t. III. 10. Then the Balance will hang in *Æquilibrium*.

24. SINCE the Velocity of Weights hanging to a Balance depend upon their Distances from the Center of Motion, each Weight being multiplied by its Distance from that Center will give its *Momentum*: Therefore  $8 \times 3 (= 24) + 5 \times 6 (= 30) + 3 \times 1 (= 3) + 3 \times 9 (= 27)$  make up the Sum of 84 for the *Momentum* or Quantity of Motion on the *Brachium* A C: And  $2 \times 2 (= 4) + 5 \times 12 (= 60) + 10 \times 2 (= 20)$  make the like Sum 84 for the *Momentum* on the *Brachium* C D: And consequently these equal *Momenta* acting in contrary Directions must produce an *Æquilibrium* \*. If all the Weights on the *Brachium* A C were reduc'd into one, viz. a twenty-one Pound Weight hanging at the 4th Division, it would keep in *Æquilibrium* a Weight equal to all the Weights on the *Brachium* C D hanging at a quarter of a Division beyond the 5th, because  $4 \times 21 = 5\frac{1}{4} \times 16 = 84$ , as has been shewn in the second Lecture \*. \* L. 2. 13.

### P R O B L E M.

25. *Equal or unequal Weights being suspended at the Ends of a Balance of known Length and Weight. How to find the fix'd Point or Center of Motion about which the said Weights will be in Æquilibrium.*

#### EXPERIMENT III. Pl. 9. Fig. 7.

A B is a Balance weighing four Ounces, and of 12 Inches in Length, at whose Ends A and B are suspended Weights of four and eight Ounces. Find out the common Center of Gravity of the said Weights \* which will be directly under the Point K at Numb. 4. Let K be made the Center of Motion, and then the Center of Gravity being at its lowest Place \*, the Problem will be solv'd, if the Balance A B has no Gravity. But as the Balance weighs four Ounces, the *Brachium* A K will overweigh the *Brachium* K B, and so destroy the *Æquilibrium*. Now a second Operation like the former, will perfectly solve the Problem, after the following Preparation. Suspend the Weight E equal to the two former Weights or 12 Ounces at the Hook under K at the common Center of Gravity of the two Weights; which

Lect. III. which reducing their Weights into the Center of Gravity, they will act as before \*: Reduce also the Weight of the Balance into its Center of Gravity, by hanging the Weight D of four Ounces (or equal to the Weight of the Balance) at  $\kappa$  under 6 the Center of Magnitude, which is also the Center of Gravity, because the Balance is regular and homogeneous. Then shall we have the short Balance 6 4 or  $\kappa$  K without Weight, and the Point C or true Center of Motion will be found by this Analogy,  $E + D$  (16 Ounces) : D (4) ::  $\kappa$  K (or a Length of two Inches) : KC (or half an Inch.) This Point will be directly over the Center of Gravity of the Balance and all the Weights; and taking away the Weights D and E the *Æquilibrium* will remain, the Alteration of the Center of Motion from K to C making amends for the unequal Weight of the *Brachia* of the Balance.

\* L. 2. 34.

### P R O B L E M.

26. *A given Weight hanging at one of the Ends of a Balance of known Weight, to find the fix'd Point about which the Balance and Weight will be in Æquilibrium.*

### EXPERIMENT IV. Pl. 9. Fig. 9.

HAVING suspended the given Weight D equal to four Pounds (for example) at the End A of the Balance AB, which weighs also four Pounds; since the Weight of the Balance is consider'd in this Operation, we must suppose the whole Weight of the Balance reduc'd into its Center of Gravity \*, as if the Weight E of four Pounds hung at C the Center of Gravity of the Balance, which is at its middle Division 6: then we shall have a new Balance (*viz.* AC) without Weight, at whose Ends hang the Weights D and E, whose fix'd Point will (by the last Prop.) be found at 3; or more generally by this Analogy.

\* L. 2. 43.

As  $D + E$  (or the Weight of the Body and the Weight of the Balance):

Is to E (The Weight of the Balance) ::

So is CA (Half the Length of the Balance):

To A 3 (The Distance of the fix'd Point from the given heavy Body.)

P R O.

P R O B L E M.

27. To make a deceitful Balance or Pair of Scales, whose *Beam* will hang in *Æquilibrium* without the Scales, or with the empty Scales; and yet shall also be in *Æquilibrium* when unequal Weights are plac'd in the Scales; so as to cheat in any Proportion intended in making the Balance at first. Lect. III.

THIS Problem is solv'd by making a *Statera* or *Steel-yard* with the Appearance of a common Beam, as in the following

EXPERIMENT V. Pl. 9. Fig. 10.

To the Beam A B (23 Inches long, whose *Brachium* CB, of 11 Inches in Length, keeps in *Æquilibrium* about the Point C the *Brachium* CA, of 12 Inches in Length, by being made so much thicker, or having so much more Matter, as may make amends for its being shorter) suspend the Scales D, E in such manner that D, which weighs one Part in twelve less than E, shall hang at the longest end of the Beam, and they will keep each other in *Æquilibrium* \*. Then placing 12 Pound Weight at G in the Scale E, it will keep in *Æquilibrium* no more than 11 Pounds of F, the Commodity to be sold if plac'd in the Scale D. Because then, F will be to G, in a reciprocal Proportion of B C to A C \*.

\* 9. 13.

\* 9. 13.

Tho' such a Balance may be so nicely made as to deceive the Eye; the Cheat is immediately discover'd by changing the Weights and the Commodity F from one Scale to another; for then the Owner of the Balance must either confess the Fraud or add to the Commodity he sells, not only what was wanting, but also as much as he intended to cheat him of, and a Fraction of that added Weight proportional to the Inequality of the *Brachia* of the Balance. That is, in this Case, the Buyer instead of 11 Pounds offer'd him for 12 his Due, will have (by changing the Scales)  $12\frac{1}{11}$  Pounds. For whereas in the first Position of the Balance  $F (11) \times A C (12)$  was equal to  $G (12) \times B C (11)$ , when G or 12 Pounds is plac'd in the Scale D, then  $12 \times 12$  will be equal to no less than  $C B (11) \times 13\frac{1}{11} G$ ; or,

As

Lect. III.

As the *Brachium* C B, 11 Inches long:Is to the *Brachium* C A, 12 Inches long ::

So will be F, or the Weight 12, plac'd in the Scale D :

To  $G = 13\frac{1}{11}$ , or the Weight of the Commodity keeping the Weights in *Æquilibrio*.

AND therefore as this Analogy gives a reciprocal Proportion between the Weights and their Velocities, the *Momenta* will be equal, which with contrary Directions destroy one another.

N. B. *In all these Cases we suppose the Weight to hang freely from those Ends of the Balance to which they are fasten'd.* See the

\* Ann. 5. other Cases in the Notes \*.

### Of the L E A V E R.

28. THE *Leaver* (a known Instrument commonly call'd a *Hand-spike*, when of Wood, and a *Crow*, when of Iron) is, in Theory, to be look'd upon as an inflexible Line like the Beam of a Balance, and subject to the same Proportions, only that the Power applied to it is commonly an *animate* Power; and from the different ways of applying it, it is call'd a Leaver of the *first*, of the *second*, or of the *third Kind*.

### EXPERIMENT VI. Pl. 9. Fig. 11.

29. LET the Steel-yard P W be taken off of its Hook K, and let its Center of Gravity C be plac'd upon a *Fulcrum* or triangular Prism D E; and instead of the Weight 1 hanging at P to keep in *Æquilibrio* the Weight W of 4 Pounds, let an animate Power, as the Hand, be applied at P: The *Statera* or *Steel-yard* will then be turn'd into a *Leaver of the first Kind*, so call'd because the *Fulcrum* or fix'd Point is between the Ends as at C, in which Case the Power may be four times less in Intensity than the Weight; but equal to it, if C or the *Fulcrum* was remov'd to M (the middle Point of P W) and four times greater, if C was remov'd to 3. N. B. *In all these Cases, the Leaver is still said*

\* Annot. 6. *to be of the first Kind* \*.

30. WHEN the *Fulcrum* is at one End, the Power at the other, and the Weight between them, the Leaver is of the *second Kind* \*.

\* Plate 9.  
Fig. 12.

31. BUT

31. BUT it is of the *third Kind*, when the fix'd Point is at one End, the Weight at the other, and the Power between them \*.

N. B. *The Power and Weight are always supposed to act at right Angles with the Leaver, except it be otherwise express'd; for then the Cases will vary, as may be seen in the Notes \* upon the Balance, which are equally applicable to the Leaver.*

THE Proportions, which the Powers and Weights bear reciprocally to their Distances, are set down at the 11th, 12th and 13th Figures.

EXPERIMENT VII. *Pl. 9. Fig. 14.*

32. PLACE the three Leavers A, B, D, in such manner upon the *Fulcra* F, F, F, that A may have the Proportion of its *Brachia* as 5 to 1, B as 4 to 1, and D as 6 to 1; and let these Leavers act upon each other; then making use of a Power equal to one Pound at the End M of the Leaver D, it will keep in *Æquilibrio* W, or 120 Pounds at the End I of the Leaver A.

IN this compound Leaver, the *Proportion* or *Ratio* of the Weight W to the Power M is compounded of the several *Ratios* of the long *Brachium* of each Leaver to its short one; for  $5 \times 4 \times 6 = 120$ . And accordingly you will find by measuring the Ascent and Descent of the Ends of the first and last Leaver, that whilst the Weight W descends  $\frac{1}{120}$  of an Inch, the Power or little Weight M will ascend  $\frac{120}{1}$  or 12 Inches, the Force to be gain'd by such a compound Leaver being shewn by the reciprocal Proportion between the Masses and Velocities of W and M.

WHEN two Powers applied to the Ends of a Leaver, support a Weight resting upon the Leaver, they are to one another reciprocally as their Distances from the Weight. The Proportion is mark'd in the Figure \*.

\* Plate 9.  
Fig. 15.

EXPERIMENT VIII. *Pl. 9. Fig. 16.*

33. IN the Frame A B C D hang the Leaver E F at the Points I, K, loaded with the Weight 7 at C; and the Weights whose Ropes go over the Pullies G H, being to one another reciprocally as the Distances I C, and C K, will keep in *Æquilibrio* the

O

seven

Lect. III. seven Pound Weight. N. B. *The Weight of the Leaver is not consider'd here; for unless the whole Brachia EC and CF be to one another as 3 to 4 the Center of Gravity (or Point C) must be remov'd a little farther towards I, as was said before concerning the Balance* \*.

\* 25.

It is upon this Principle that Horses of unequal Strength may draw equally in a Coach; for if the Spring-tree Bar be unequally divided, that Horse must employ more Strength which is applied to the short End of the said Bar. Two Men also who carry a Barrel hanging from a Staff are unequally press'd upon their Shoulders, if the Barrel does not hang in the middle, the Man carrying most who is the nearest. This is farther illustrated by

#### EXPERIMENT IX. Pl. 6. Fig. 17.

34. THE Foot F supports an horizontal Board AB, on which must be laid the Leaver  $12$  divided in the Proportion of  $2$  to  $1$  (supposing it a Spring-tree Bar) then placing the Pulley  $n$  over-against  $2$ , and the Pulley  $m$  over-against  $1$ , let the Weights O ( $1$  lb) and M ( $2$  lb) hang by Strings over the said Pullies, and if they will keep in *Equilibrio* the Weight N =  $3$  lb drawing the Leaver in a contrary Direction over the Pulley  $o$  against C. N. B. *The Pullies slide in a Groove in the Edge of the Board, and remain in any Place where they are set.*

#### EXPERIMENT X. Plate 9. Fig. 18.

35. IF there be a Leaver whose *Brachia* make an Angle, as is represented in the Figure, with its fix'd Point at the Angle as C; the Weight W, pressing perpendicularly upon the End W, will be kept in *Equilibrio* by a Weight of one Pound drawing the other End of the Leaver P perpendicularly (by means of a Pulley over which the Line of Direction of the Power is carried) by two Pounds at  $p$  and three Pounds at  $\pi$ . It is in this manner that a Hammer is made use of to draw a Nail. Some call this a Leaver of the fourth Kind; but it is evidently a Leaver of the first Kind, because the Weight W is at one End, the Power P at the other, and the Center of Motion C between: And if the Arm CW be set streight in a Line with PC, so as to bring W to  $w$ , and  $c$  be made the

*Fulcrum*

*Fulcrum*, the Instrument will plainly appear to be a Leaver Lect. III. of the first Kind \*.

\* Ann. 7.

*Of the P U L L E Y.*

26. WHEN a little Wheel, commonly a *Sheave* or *Sheever* is so fix'd in a Box or Block as to be moveable round a Center-Pin passing thro' it, such an Instrument is call'd a Pulley \*: And sometimes, tho' improperly, a Box or Block with several Sheevers in it is also call'd a Pulley, as in the second Figure \*. The first of these is by Workmen a *Snatch-block*.

\* Plate 10. Fig. 1.

\* Pl. 10. Fig. 2.

A Rope going round one or more Pullies to raise a Weight, is call'd the *Running-rope*.

WHEN a Block with its Sheevers is so fix'd, that whilst it remains immoveable, another Block and Sheevers rises with the Weight hanging at it; such a Machine is call'd a *Pair of Blocks*.


37. AN upper or fix'd Pulley adds no Force to the Power, but only prevents the Friction by making the Rope run easily; and so much the more as the Sheever is bigger than the Center-Pin about which it turns \*.

\* Ann. 8.

EXPERIMENT XI. *Pl. 10. Fig. 3.*

HAVING fasten'd to the Ends of a pliable Rope the Weights 1 and 3, the first of one, and the other of three Pounds, if the Rope be thrown over the Square but smooth Beam A B, the Friction of the Rope on the Beam will be so great as to hinder the three Pound Weight from raising the one Pound, tho' its *Momentum* (without that Hindrance) is three times greater, because its Velocity is the same and its Matter triple \*. But if only one Pound be fasten'd at the End of the said Rope, and the Rope thrown over the Pulley E D, the single Pounds will keep each other so exactly in *Æquilibrio* that the least Weight added to either of them will make it over-power. That these Weights ought to be in *Æquilibrio*, appears from a sight of the Figure, since the Weight on the right Hand cannot descend to *d* without causing the Weight on the left to ascend up to E precisely with the same Velocity, 1 *d* being equal to 1 E. N. B. This Pulley is also call'd a Roller.

\* L. 2. 3.

Leçt. III. 38. A lower Pulley, that is such a one as is moveable with  the Weight, takes off half the Weight; so that a Power of half its Intensity will sustain it.

EXPERIMENT XII. *Pl. 10. Fig. 4.*

To an Hook coming from the Center of the Pulley *g e* hang a Weight of 2 lb; then having made fast the running Rope to the Hook *f* of the Arm *A*, bring it under the said Pulley *g e* and over the Pulley *d*, and 1 lb at the End of the Rope will sustain the 2 lb hanging from *c* the Center of the Pulley *g e*.

\* 37. THAT the Power pulling down at *i* acts in the same manner as if it pull'd upwards at *d* is evident, because we have already shewn \*, that an upper Pulley neither encreases nor diminishes the Action of the Power.

WE may at one View see in Pullies how the Force of a Weight is diminish'd, by considering how many Ropes (or Parts of the Rope) are employ'd to raise it, and which divide the Weight as they are applied to the lower Pullies to which the Weight hangs, whilst the Power only draws by one Rope. For Example, in this Case the Ropes *f e* and *d g* sustain the Weight; but *f e* is supported by the Hook *f*, whilst the Power only draws up the Rope *d g*.

39. FROM this we may deduce this general Rule to know the Advantage to be gain'd by a Pair of Blocks, let their Number of Pullies or Sheevers be what it will, (*viz.*) *As One is to the Number of the Ropes (or of the Parts of the Rope) applied to the lower Pullies; so is the Power to the Weight.*

\* Plate 10. FIG. 4, 5, 6, 7, 8. THUS it is evident, only by a sight of the Figures \*, that one Pound will sustain 4 lb as in *Fig. 5*. Six Pounds as in *Fig. 6*. Five Pounds, as in *Fig. 7*: And six Pounds as in *Fig. 8*. N. B. *The Pullies and their Ropes represented by the 5th, 7th, and 8th Figures are call'd Tackles of Four, Tackles of Five, and Tackles of six.*

40. THE Machine represented in *Fig. 6*. is the most inconvenient for raising the Weight 6; but the most convenient for bringing together the Ends of two Beams without danger of bend-

bending them, as if the Ends A and B were to be brought together gradually. N.B. *This Way of using Pullies is call'd lacing.* Le&t. III.


41. WE must observe that the Rule abovemention'd is only applicable to the Cafes in which the lower Pullies rise all together in one Block with the Weight; but when they act upon one another, and the Weight is only fasten'd to the lowest of them, the Force of the Power is very much encreased, each Pulley doubling it. As for Example \*, a Power whose Intensity is equal to 8 lb (applied at *a*) will, by means of the lower Pulley A, sustain 16 lb \*: A Power equal to 4 lb (at *b*) will, by means of a lower Pulley B, sustain the Power of 8 lb acting at *a*: A third Power equal to 2 lb (at *c*) will, by means of the Pulley C, sustain the Power of 4 lb at *b*: A fourth Power of 1 lb (at *d*) will, by means of the Pulley D, sustain the Power of 2 lb (at *c*); and this is not alter'd by having its Rope carried over the upper Pulley or Roller E \*. N. B. *What Weight each Pulley and each Rope bears in this System of Pullies is set down in the Figure.* Pl. 10. Fig. 9. 38. 37.

## EXPERIMENT XII. Pl. 10. Fig. 9.

A Weight of 16 lb hanging at the Pulley A, (of the Machine Fig. 9. made up of four singly moveable Pullies, a Roller at E and four Hooks upon the Arm E F) let the little Weight 1, which keeps in *Æquilibrio* the Weight 16, be raised up to *k* 16 Inches high, and the Weight 16 will only descend in the Line *g h* an Inch long. This shews the reciprocal Proportion between the Weights and their Velocities to be applicable to this, as well as to all other Cafes of Pullies, as may be known by moving the Weight or Power in any Combination of Pullies, and measuring the Spaces which they go thro'. Thus in Fig. 4 \*, while 2 goes down to *a*, 1 goes up to B, just twice as far, Pl. 10. Fig. 4.  
 Sc. \* *This Proportion therefore will always produce an Æquilibrium in this as well as in all other Mechanical Engines.* Ann. 9.

N. B. *The Ropes, ascending and descending, are always to be suppos'd parallel, except where it is otherwise express'd \*, and in every Figure representing Pullies, the Power and Weight are mark'd with the Letters P and W.* Ann. 10.

*Of the A X L E in the Wheel.*

Lect. III. 42. WHEN a Power by help of a Rope, or any other Means,  is so applied to the Circumference of a Wheel as to cause the said Wheel together with its Axle to turn round and raise a Weight applied to the Axle in any Manner, such a Machine is call'd an *Axle in the Wheel*, or *Axis in Peritrochio* \*.

\* Pl. 10.

Fig. 10, 11.

SINCE in this Instrument the Wheel and its Axis move together; it is evident, that in one Turn of the Wheel, when the Power \* P descends a Length equal to the Circumference of the Wheel, the Weight W rises an Height equal to the Circumference of the Axis A by the winding of the Rope which carries the Weight upon the said Axis. And since, when there is an *Æquilibrium* between two Weights as W and P, there must be a reciprocal Proportion between their Masses and Velocities; W will be to P, as the Circumference of the Wheel to that of the Axis (supposing the Rope of no Thickness) or as the Semidiameter of the Wheel to the Semidiameter of the Axis (that is as \* D K to K A) because the Semidiameters of different Circles are in the same Proportion to one another as the Circumferences.

\* Plate 10.

Fig. 11.

HENCE it follows, that the less the Axle is in Proportion to the Wheel, the greater may the Weight be, which is sustain'd or rais'd by the Power.

EXPERIMENT XIII. *Pl. 10. Fig. 10.*

43. THE Machine represented by the Figure is a Model (made by a Scale of an Inch to a Foot) of such an Axle and Wheel as is often made use of to draw Water out of a Well, by means of a Power drawing by a Rope applied to the Circumference of one of the Wheels of the Machine, (or by pressing down successively the Handles E, F, G, H, I, K, whilst another Rope or Chain is wound up upon the Axis A or B, a Bucket hanging at it instead of the Weight W. Here in the Experiment one Pound hanging at the Circumference of the biggest Wheel C D will keep in *Æquilibrio* 12 Pounds hanging at the smallest Axis A, or 6 Pounds at the Axis B, and only 3 Pounds upon the Circumference T V. In the same manner, when the Weight hanging at the Axis continues in the same Place, and to be of the

the same Quantity, *viz.* 12 Pounds, then the Power which, at Lect. III. the Circumference of the Wheel C D, is equal to 1 Pound, must be equal to  $1\frac{1}{2}$  Pound, if it be applied at S R; but if it be applied at any of the Handles at the Distance of  $\frac{1}{4}$  of an Inch from the Circumference of the Wheel C D (which is the same as if a new Wheel was added of  $\frac{1}{2}$  an Inch more in Diameter) then a Power equal to no more than  $\frac{1}{4}$  of a Pound, will keep the Weight in *Æquilibrio*; and raise it, if its Intensity be encreas'd ever so little.

THIS is more clearly represented in Fig. 11. where the Weights\* Plate 10. are represented by the Letters W, w, w, and the Powers by the Letters P, p,  $\pi$ , and where you must observe, that unless the Power acts upon the Handles abovemention'd at right Angles, or in the Lines E  $\pi$ , F f, or G g, &c. the Effect cannot be the same as making use of a new Wheel E, F, G, H, to which the Lines E  $\pi$ , F f, and G g are Tangents. Fig. 11.

44. FOR if a Power, as for Example, P, acts obliquely upon any of the Handles, as in the acute Angle P F K, or its correspondently obtuse Angle P I K, the Line of Direction of the said Power \* becomes the Tangent of the Circumference D C; and consequently the Power acts as if it drew by a Rope going round the Wheel C D: And if the Obliquity was greater as when the Power p draws the Handle G in the Angle p G K, the Effect will be no greater than if the said Power should draw by a Rope going round the Circumference S R. The Powers in these Cases must be encreas'd in the same Proportion as the Lines D K and S K are shorter than E K. See the Numbers in the Figure, which express the Intensities of the Powers. \* Ann. §.

N. B. We have taken no Notice here of the Thickness of the Rope, to which a Regard is to be had in Practice, always adding half the Thickness of the Rope to the Semidiameter of the Axis: And if the Rope is wound up upon it self, for every such Turn we must still add half its Thickness, which is the Reason why more Power must be applied when the Axis is thus thickened, as often happens in drawing Water from a deep and narrow Well, over which we cannot have a long Axis.

45. IF the Rope to which the Power is fasten'd be successively applied to different Wheels, whose Diameters are bigger and bigger, the Axis will continually be turn'd with more ease, unless

Lect. III. unless the Intensity of the Power be diminish'd in the same Proportion ; and if so, the Axis will always be drawn with the same Strength by a Power continually diminishing. This is practis'd in Spring-Clocks and Watches, where the Spiral Spring S, which is strongest in its Action when first wound up, draws the Fuzee F, or continued Axis *in Peritrochio* by the smallest Wheels near B ; and as it unbends and grows weak, draws at the larger Wheels near A, in such manner that the Watch-work is always carried round with the same Force \*.

\* Plate 10.  
Fig. 12.

46. As Leavers and Pullies acting upon one another are sometimes joyn'd together to encrease the Action of the Power, whereby it may sustain or lift a greater Weight ; so it is usual to make a compound Axis *in Peritrochio* by combining together two or more of these Machines. Because tho' by means of a long Axle capable of receiving a great deal of Rope, one may draw Weights from a great Depth ; yet, as a very small Axis would be too weak for very great Weights, or a large Wheel would be very costly, if sufficiently strong, or make a cumbersome Engine which would take up too much room ; it is more advisable to combine Wheels and Axles, by means of Pinions or small Wheels upon the Axles, whose Leaves (or Teeth) take hold of Teeth made in the large Wheels, as we see in Clocks which have several Axles and Wheels, or some sort of Cranes which have only two of them combin'd together.

\* 82, 83.

#### EXPERIMENT XIII. Pl. 10. Fig. 13.

47. THIS Machine consists of two Wheels with their Axles, the first of which A B C (on whose Circumference A B is coil'd the Rope which carries the Power P, which is a Weight of one Pound) has a Pinion of eight Teeth on its Axis at C, which takes the Teeth of the Wheel F G of the other Axis *in Peritrochio*. The Wheel F G has forty Teeth, and its Axis H K is in Diameter the eighth Part of the Wheel A B. Hang the Weight W of 40 lb upon the Axis H K, and it will be kept *in Equilibrio* by the Power P, which is equal but to 1 lb.

IF we suppose the Axis C I of the same Diameter as the Axis K H, it is evident that the Power P would have sustain'd but 8 lb hanging on the said Axis ; nor would it have sustain'd more

more on the Axis K H if this last Axis had gone round as often as the first Wheel A B (which would have happen'd if the Wheel F G had had no more Teeth than the Pinion C) but since its Wheel has five times more Teeth than the Pinion on the Axis C I, it must go five times slower than the said Axis; and consequently the Weight W goes up five times slower than it would have done on the Axis C I; therefore it has forty times less Velocity than the Power P, rising only one Inch whilst P descends forty.

Lect. III.  
Plate 10.  
Fig 13.

HENCE it follows that the Ratio of the Power to the Weight is compounded of the Ratio of the Diameter of the Axis of the last Wheel (where the Weight hangs) to the Diameter of the first Wheel (where the Power is applied) and the Ratio of the Number of Revolutions of the last Wheel, to the Number of Revolutions of the first Wheel in the same Time. As for example, here the first Ratio is of 1 to 8, and the last Ratio is of 1 to 5; therefore the Ratio of the Power to the Weight is as 1 to 40, viz. the Composition or Multiplication of those two Ratios, because  $5 \times 8 = 40$ . *And this holds good, let the Machine consist of ever so many Wheels.*

### *Of the Inclined PLANE.*

48. FOR the better understanding the Use of the inclin'd Plane in Mechanics, we must remember what has been said before concerning the Velocity of a Weight \*, viz. that *whatever* \* Ann. 8. *Line the Weight describes in its Ascent by the Action of the Power,* Lect. 2. *we are to call its Velocity only that Line which represents its perpendicular Ascent or Descent.*

IF it should be requir'd to lift up a very heavy Body as W or w \*, the Height C B, it would be impracticable to raise it up in the Line C B without a Power whose Intensity is equal to that of the Weight; and even in that Case very inconvenient to do it, especially in building. But if an inclin'd Plane A B, be laid raising from the horizontal Line A C, from whence the Weight is to be raised, a less Power than the Weight will serve for that Purpose, unless it pushes the Body directly against the Plane, (as in the Direction W T) or draws the Body away from the Plane (as from W towards e, t, or L, or) in any Direction on that side of the Line E e \*.

\* Ann. II.

Lect. III.

Plate II.

Fig. 14.

49. THE Direction in which the Body can most easily be drawn or push'd up the Plane (as in driving a Wheel-barrow) is in the Line  $W w M$ , parallel to the Plane and passing thro' the Center of the Weight; for whether the Power drives a Plane  $k K$  (in a Direction perpendicular to it) along the Line  $W M$ , or the Power  $P$  (by its Descent to  $p$ ) draws it in the same Line, the Velocity of the Power will be equal to the Line  $W w$ , the Space describ'd by the Center of Gravity of the Weight, whilst the said Weight rises only the perpendicular Height  $Z B (=n W)$  or has the said Line properly to express its Velocity. If the Body was a Cylinder, as a rolling Stone, and the Plane  $T t$  were to pass thro' the Gudgeons or Axis of the said Stone; it is evident that the Case would be the same; and as the Weight  $P$  has its Rope running over the Roller (or upper Pulley)  $M$ , the Line  $P p$  will be the Velocity of the Power. Therefore in this Case the Weight (if kept in *Æquilibrium*) will be to the Power as  $W w (=T B)$  to  $w Y (=B Z)$  or as the Hypotenuse  $A B$  is to the Perpendicular  $B C$ , which (by *Eucl.* 4. 6.) are in the same Proportion; and consequently if the Power be ever so little encreas'd, it will draw the Weight up the Plane.

N. B. *In Practice the Power must be pretty much encreas'd if the Body is not smooth and spherical or cylindrick, and the Plane very smooth. But as all sorts of Bodies are to be drawn up, to reduce them as near as we can to a Sphere or Cylinder, we must fasten Wheels to them, or (which is the same Thing) lay them on a Wheel-Carriage.*

50. THAT the Power acts with the greatest Advantage, whilst it draws in the Line of Direction  $W w$  (parallel to the Plane) is evident; because if one End of the said Line of Direction remaining fix'd at  $W$ , the other should move towards  $B$ , or beyond it, then the Body would be partly drawn against the Plane, and therefore the Power must be encreased in Proportion to the greater Difficulty of Traction: And if the End  $w$  of the Line abovemention'd should be carried to  $D$ , or beyond it, the Power must be also encreased in as much as it endeavours to lift the Body off from the Plane. *How much the Power must be*

*be encreased in Proportion to the Angle, which its Line of Direction makes with the Plane, will be shewn in the Notes \*.*

Lect. III.  
\* Ann. 12.


51. IF the Power draws in a Line of Direction  $W B$ , parallel to the Base of the Plane; then in order to keep the Weight  $W$  in *Æquilibrio* by the Power  $\Pi$ ; the said Power must be to the Weight; as  $Z B$  to  $Z T$ , or as the Perpendicular  $B C$  to the Base  $A C$  of the Triangle  $A C B$ . For if we suppose the Pulley  $R$  at so great a Distance from  $W$ , that the Line of Direction  $W R$  may not sensibly alter its horizontal Position, whilst the Body  $W$  rises the Height  $B Z$ , then will the Power  $\Pi$  descend to  $\varpi$  whilst  $W$  rises the Height  $B Z$ , in such manner that  $\Pi \varpi$  ( $= W Y$ , and not  $W \varpi$ ) will be the Velocity of the Power. So that the Velocity of the Power to that of the Weight will not be as the Hypotenuse to the Perpendicular, as in the former Case, but as the Base to the Perpendicular in the Triangle  $A C B$ .

IF the Power be encreased just enough to overcome the Friction of the Plane and draw up the Body  $W$ , let the Pulley  $R$  be lifted up gradually to  $r$ , so as to keep the Line  $W R$  parallel to itself till it comes to  $\varpi r$ , and the Power will be descended to  $\pi$  when the Weight is come to  $\varpi B$ . But  $\Pi \pi$  together with the Distance  $R r$ , is equal to  $\Pi \varpi$ , or  $W Y$ , &c. And this Traction being constantly made in the Angle  $W B T$ , is the Case

*Of the W E D G E.*

52. THE Wedge is a short triangular Prism, whose two opposite and parallel Planes are rectangular Triangles, such as the Section  $A B C$ \*, the rest being rectangular Parallelograms; the Edge or entring Part of the Wedge is made by the meeting of two Planes in whose Section is the Point  $A$ , and the Back of it is the Plane opposite to the Edge, on which the Hammer or Mallet strikes to drive the Wedge forwards. The Representation of it is seen in the Figure  $A B C D E$  \*.

IF upon the horizontal Line  $A C$ \* produc'd towards  $p$ , be laid the Weight  $w$  at the Point  $d$ , and a Plane as  $G g$  to stop it from going towards  $A$ , whilst the Wedge  $A B C$  is push'd under it from  $d$  towards  $A$ ; as the Wedge is driven from  $C$  to  $A$ , the whole Length of its Base  $A C$ , the Weight  $w$  rises just the

Lect. III. Height  $CB$ , or the Thickness of the Wedge: Therefore the  
 Power will be to the Weight, as  $BC$  to  $AC$ .

53. THE Proportion of Force us'd would be just the same if a Plane as  $Ff$  were to move parallel to it self, and perpendicular to  $AC$ , and push up the Weight, as  $W$ , from  $A$  to  $B$  along the Wedge suppos'd then immoveable. Nay, it would be the same if the Weight was only push'd from one part of the Wedge to another, the Plane moving but from  $Ff$  to  $Gg$ , or (which is the same) from  $Ee$  to  $Dd$ ; for then  $wY$  would express the Rise of the Center of Gravity (or the Velocity) of the Weight, and  $WY$  the Velocity of the Power, which are still in the Ratio of  $BC$  to  $CA$ .

THIS is the second Case we mention'd concerning the inclin'd Plane, and may be confirm'd by the following

#### EXPERIMENT XIV. *Pl. II. Fig. 2.*

\* Plate II. Fig. 2. 54. TAKE the Machine describ'd in the Figure, in which the smooth Plane or Board  $BAHI$ , moveable on Hinges at  $B$  and  $I$ , may be rais'd so as to make any Angle with the horizontal Board  $NLI$ , by help of the Quadrant  $D$ , and fix'd in its Position by means of the Screw at  $T$ ; and having made fast the Head  $DE$  in the Groove  $Ss$  by a Nut under the horizontal Board, raise the Pulley  $C$  of the Arm  $DC$ , so that a Line going over the said Pulley in the Direction  $CM$ , may be parallel to the Plane  $AB$ . Then take a little wooden Cylinder  $M$ , the Ends of whose Axis or Pivots go thro' a brass Frame, as at  $Q$ , so that it may be drawn easily by a String fix'd at  $M$ ; and lay on the said Cylinder upon the inclin'd Plane, having fasten'd to  $M$  a String which goes over the Pulley  $c$  and is joyn'd to the Ball  $P$ , which being made use of as a Power will sustain the Cyndrick Weight  $M$ , when  $P$  is to  $M$  as  $AC$  the Height of the Plane, to  $AB$  its Length. *A little more Weight added to  $P$  makes it draw up the Cylinder.*

BUT if the Arm  $EC$  be lower'd, so as to come into the Position  $EG$ ,  $p$  must be to  $M$  (in order to keep it in *Æquilibrio*) as  $AC$  the Height of the Plane, to  $CB$  its Base. *N. B. Here it is to be observ'd that tho' the Addition of a little Weight to  $p$  will make it begin to draw the Cylinder  $M$  up the Plane; that the Cylinder will not be drawn quite up by the Descent of  $p$ , because*

cause the Angle which the Line of Direction of the Power makes with the Plane will be encreased as the Cylinder rises up; but if the Pulley G be gradually lifted up to K, whilst the Cylinder is drawn up to m, the Power, which then will be at  $\pi$ , will have the same Effect on the Weight at m, as it had before at M, &c. and the Weight (when kept in *Æquilibrio* in the Line of Direction MG or m K) will still be to the Power as M n to n m, or BC to CA.

THIS Machine also experimentally shews the Effect of the Power, whatever Angle its Line of Direction makes with the Plane.

55. THE Wedge, which we have hitherto consider'd, is a Wedge acting in the most simple manner, only with one of its Surfaces; for when it slides upon the Plane CA\* to lift the Weight W, it does only act with its Surface AB, the Surface AC only applying it self to the Line Ap, without removing it from its Place. Thus if a Moulding was to be separated from a Wainscot as Mm\* from Ww by Means of the Wedge ACB, it is plain that *a* the Velocity of the Wedge (so far as it is driven in, when from the Position *abc* it comes into the Position ABC) wou'd be to *am* the Velocity of the Moulding as AC to BC. Therefore, &c. So it would be if a Pillar standing upon a Floor was to be raised up without moving the Floor.

56. BUT in the common way of using the Wedge, both Sides of it act, as in cleaving Wood: Then the Proportion of the Power and Weight are different from the former; for then the Power will be to the Weight as half the Wedge's Thickness to its Length. But then, we may easily reduce this to what has been said before; because here we make use of a double Wedge. For let us suppose Ca\* an immoveable Plane, and on each Side of it a Wedge, as BCA and bCA, made use of to remove a Weight as *c* or *d* from the said Plane; the Power pushing the Wedge will be to the Weight but as *c* C (= AC = BC) or *d* D (= AD = Cb) to the Length of the double Wedge CBA b, which is now got to A c a d. Some Mechanical Writers have mistaken this Case by adding together the single Velocities of *c* and *d*, viz. Cc + dD, and calling that Sum the Velocity of the Weight, which they compar'd with Aa the Velocity of the Power; but we must consider that if the two Bodies

Lect. III. Bodies to be remov'd from each other (as the Parts of the Wood to be cloven) were laid upon one another on, either Side, as for example at  $d$ , their Velocity would only be  $dD$ , and they would be remov'd as easily then by one Wedge as  $bCA$ , from  $d$  to  $D$ , as by the two Wedges in the former Case. *For tho' in considering the Momentum of Bodies, the Sum of the Momenta of all the Parts is to be taken for the Momentum of the Whole; we are not to take the Sum of the Velocity of the Parts for the Velocity of the whole, but only the Velocity of the Center of Gravity of the Body, when we consider how far it is remov'd out of its Place. If a Leaver or Balance was made like a Fork\*, and the Part  $CA$  was just double of either of the Parts of the Fork as  $CB$  or  $CD$ , and the Center of Motion fix'd at  $C$ ; it is evident that one Pound at  $A$  will keep in Æquilibrium two Pounds at the forked End, tho' the Pounds were to hang single, a two Pound Weight being always equivalent to two single Pounds; but if the Sum of the Velocities of  $B$  and  $D$  were to be taken, there must then be two Pounds at  $A$  to sustain the Weights at  $B$  and  $D$ , the Radius  $CA$  giving no more Velocity to the Weight at  $A$ , than the Radius  $CB$  or  $CD$ , of half its Length, to the Weights at its Ends, which is absurd, &c.*

\* Plate 11.  
Fig. 5.

57. In both Cases of the Wedge, it is not only necessary to apply a Force something greater than either in Proportion of the half Thickness or of the whole Thickness of the Wedge to its Length, that the Power may overcome the Weight; but because the Surfaces even of the most polish'd Wedge are very rough (in Comparison to the Mathematical Smoothness which we have suppos'd) and the Bodies to be separated likewise very far from having their Surfaces truly plane; we must make use of an additional Force, to overcome the Stickage arising from that Roughness.

This Friction or Stickage, which is not great in other Engines, is very considerable in the Wedge, Experience shewing that a Wedge laden with a vast Weight, has hardly any Effect, especially in cleaving Wood; because not only the Surfaces of the Wedge, as was said before, but the Parts of the Wood to be cloven are always rough, and so close, that their Friction very much hinders the Motion; which Obstacle we endeavour to remove by Percussion, which here is of wonderful Use; for Experience shews, that a Blow upon the Head of a Wedge, makes it enter


enter easily into a hard Body; the Reason of which seems to be, *Leſt. III.* that a Blow, by putting all the Parts of the Wood in motion, makes them tremble and be diſunitied, ſo as to leſſen the Stickage, and facilitate the Motion of the Wedge. The Effect of Percuſſion will be greater in Proportion as the percutient Body is heavier and moves ſwifter.

*Plate II. Fig. 6.*

58. To prove by Experiments what has been ſaid of the Wedge, we muſt make uſe of the Machine repreſented in the 6th Figure of *Plate II.*

A B C D is a Braſs Frame, conſiſting of two horizontal Pieces A B and C D, and two vertical ones A D and B C, ſtanding upon, and fix'd to the former, each of theſe laſt Pieces has on the Inſide about the middle, two little Pullies N O, P Q not exactly in the ſame Plane, leſt the String that goes over the one ſhould fall foul upon the String which goes over the other. E F, G H are two Cylinders with Steel Axes, which are brought together (their Axes rolling on the Vertical Pieces) by the Deſcent of the Weights R S, each of which is divided into two Parts, by means of its Pulley T or V, ſo as to draw the Cylinders equally towards one another, by the Strings and Braſs Loops T N H, and t Y y, and two ſuch other Loops at the other End of the Axis of the Cylinders. That the Cylinders may come cloſe together, without touching the Pulleys N, O, P, Q, the Plates at their Baſis (which are a pretty deal larger than the Cylinders) are made convex towards the Ends of the Axis. The two Plates Z M, Z M, juſt wide enough to apply themſelves to the Cylinders, without rubbing again the Plates at their Ends, are join'd at M in the Manner of an Hinge, ſo as to make a Wedge, and be open'd to any Angle meaſur'd by the graduated Arc I K L, which paſſing thro' the Plate, holds them faſt, by means of the two little Screws Z, Z. X is one of two bended Wires, whoſe Ends being ſlipp'd into two Holes, keep the Cylinder E F from coming out of its Place, and only allow it to turn upon its Axis, when the Loop F (and its oppoſite under E) is carried over the Pulley Y (and the contrary Way) to the Axis of the other Cylinder at f, when only this laſt Cylinder H G is to be push'd away by the Deſcent of the Wedge, which by its own Weight, or the Addition of the Weight W, ſeparates one Cylinder from the other, which is fix'd (when the prick'd

Line

Lect. III. Line represents the String) or removes them from each other,  
 (when Y y represents one String, and H N the other.)

#### EXPERIMENT XV.

59. EVERY thing being in the Position represented in the Figure \*, open the Wedge to any Angle at Pleasure; as for Example, to an Angle of 20 Degrees, and hang on such a Weight W, as may, together with the Weight of the Wedge, draw down the Wedge, and, by separating the Cylinders, raise the Weights R and S. Open the Wedge to an Angle of 40 Degrees, and twice the Weight will be required to bring down the Wedge; but if you slip on the Wires, such as X, to fix the Cylinder E F, and hang the four Loops and Strings upon the Cylinder H G, the Wedge acting only with one Surface to remove the Cylinder H G, will require twice the Weight to bring it down, that it would if it acted with both Surfaces; that is, being open'd to an Angle of 20 Degrees, as much Force will be requir'd to bring it down, when only one Cylinder is moveable (tho' the other can freely turn about its Axis) as if it was open'd to an Angle of 40 Degrees, and both Cylinders moveable.

\* Plate II.  
Fig. 6.


#### *Of the Screw.*

60. A Screw is a Cylinder cut into several concave Surfaces, or rather a Channel or Groove made in a Cylinder, by carrying on two spiral Planes the whole Length of the Screw, in such manner that they may be always equally inclin'd to the Axis of the Cylinder in their whole Progress, and also always inclin'd to the Base of it in the same Angle.

THE Screw may be also consider'd as a Wedge carried round a Cylinder, which in that Case is call'd the Arbor of the Screw, the Wedge so carried on making what is call'd the Thread of the Screw, as may be seen in the 7th, 8th, 9th, 10th and 11th Figures \*: The Arbor of the Screw being A B in Fig. 7, and Fig. 7, 8, 9, a c b d in Fig. 8. as if the Cylinder A C B D was inscrib'd within the Screw.

\* Plate II.  
Fig. 7, 8, 9,  
10.

IN the 8th Figure we may see the manner how a Screw is made, if it be cut out of the Cylinder P H I Q, then H K L M N O P is a spiral Line going about the Cylinder, marking the prominent Part to be left of the said Cylinder, and b k l m n o, the Line

Line marking the Depth to which the Screw is to be cut Le<sup>st</sup>. III.  
 (supposing the same Line to go round the inner Cylinder or )  
 Arbor A B C D, tho' not express'd here to avoid Confusion) Plate II.  
 and then *b L l*, *l N n*, &c. will represent the prominent Part Fig. 8.  
 or Thread of the Screw. Now, if instead of cutting the Hol-  
 lows H *b L*, *L l N*, *N n P*, &c. into the Cylinder P H I Q a  
 continued Wedge be fix'd to a smaller Cylinder as A C B D,  
 or rather *a c b d*, the same kind of Screw will be made, and  
*a b c d* will be the Arbor of that Screw. Sometimes the most  
 prominent Part of the Thread, as L, N, &c. is not sharp but  
 flat, and then the Thread is call'd a square Thread, as in *Fig. 11.*  
 which represents the Section of such a Screw. This sort of  
 Thread is not us'd in Wood, but in Iron and other Metals it  
 is of good Service, being commonly more durable, and raising  
 the Weight with more ease than the sharp Thread, as will be  
 more fully shewn in the Notes \*.

\* Ann. 13.

### *Of the Force of the SCREW.*

To make an Estimate of the Force of the Screw (which may  
 be compar'd either to an inclin'd Plane (as we have consider'd  
 it among the mechanical Powers) or to a Wedge, according as  
 its Arbor does or does not advance in a progressive Motion  
 whilst it turns round its Axis to raise or stop a Weight or  
 to press Bodies together, which are the several Uses of a Screw)  
 let us take a flexible Wedge, as for Example one of Paper, and  
 coil it round a Cylinder \* as is represented in the Figure, where \* Plate II.  
 A B is the Arbor, C I D one Thread or Helix, D H E ano- Fig. 7.  
 ther, and E F G Part of the Wedge left to shew the Pro-  
 portion between the Power that turns the Screw and the  
 Weight W.

63. IF the Weight is push'd up the Wedge (or which is  
 the same, rais'd perpendicularly by the Wedge slipping under it)  
 from F to H in the Direction W w, then will H G be the Ve-  
 locity of the Weight, and G F the Velocity of the Power, which  
 is the Case of the inclin'd Plane becoming a Wedge; and this  
 will be the Analogy for the Screw \* thus acting.

\* 52.

Q

As

Lect. III.  


As a Circle whose Diameter is  $Hb$  :

To  $HI$  the Distance of two Threads :: (or as the Base  $FG$  : to the Perpendicular  $HG$  ::)

So is the Weight ::

To the Power applied to the Arbor at  $A$  to raise a Weight up the Thread  $HDI$ .

N. B. *We suppose the Diameter of the Arbor at  $A$ , and of the Screw at  $H$  nearly equal.*

Plate II.  
 Fig. 10.

THIS is the Case of the 10th Fig. where the moveable Plank  $DK$  is carried down by turning round the Heads  $G, G$ , of the Screws  $AB$  and  $CD$ , in order to press strongly the Bodies placed between the Planks  $DK$  and  $ML$ , whilst the Piece  $HI$  fix'd on the upper Plank is either guided thro' an Hole, or being only look'd at, serves to shew whether the Plank  $KD$  be brought down horizontally as the Screws are turn'd. When long Leavers are thrust into the square Holes at the Heads of the Screws, the Force of the Screw is much encreased, and then the Weight : will be to the Power :: as the Circumference of the Circle describ'd by that Part of the Leaver to which the Hand is applied : to the Distance between two Threads. So in Fig. 13. as the Circumference of the Circle whose Radius is  $AH$  to  $Cc$  the Distance of two Threads of the endless Screw  $CD$  :: so is the Resistance of the Teeth of the Wheel  $I$  : to the Power applied at  $H$ .

Plate II.  
 Fig. 13.

\* Pl. II.  
 Fig. 7.

64. BUT if the Weight  $W$  \* be driven along on the Wedge  $HFG$  in the Direction  $Ww$  parallel to the Surface of the Wedge, then will this Case be the same as that of the inclin'd Plane \*, and therefore the Analogy for the Screw in the Progression of it will be,

\* 49.

\* Plate II.  
 Fig. 7.

As the Spiral-Line  $HDI$  : \*

To  $HI$  the Distance of two Threads :: (or as the Hypotenuse  $FH$  : to the Perpendicular  $HG$ ) ::

So is the Weight :

To the Power apply'd to the Arbor at  $A$  in a Spiral Direction parallel to  $HDI$ .

THE 9th Figure represents the Practice of this, whether the Female Screw D E be brought down on the Male Screw towards the Plank B, or the Plank B and its Screw be brought up toward D E by screwing its Screw up higher into the Box or Nut D E. Therefore when Handles as D and E are us'd to encrease the Force, The Weight will be to the Power :: as the spiral Circumference describ'd in one Revolution by their Ends to which the Power is apply'd : to the Distance between two Threads of the Screw.

Lect. III.  
Plate 11.  
Fig. 9.

THIS Case is applicable in Practice to the Screw at the End of Gimblets to facilitate the boring of Wood; to the bringing down the Nuts in large cold Presses, and to the common Use of Screw-pins for holding together the Parts of Machines.


THE greatest Part of Mechanical Writers have only taken Notice of this last Proportion of the Force of the Screw.

65. THO' in the Theory, if the Product of the Intensity of the Power into its Velocity does ever so little exceed the Product of the Weight into its Velocity, the Power must raise the Weight; yet here the Power, or its Velocity, must be sensibly encreased to produce this Effect in Practice by Reason of the great Friction in the Screw, which is the same that we have taken Notice of in the Wedge; only we are to observe that there is more Friction in the sharp than the square Thread, which may appear by a Sight of the 14th Figure \*, where B A C D represents the Section of a square Thread, and *b a d* the Section of a sharp Thread. As the Part A C does not touch the Female Screw, which only bears with its Force and Weight upon the Flat A B, the Line A B in the square Thread compar'd with *a b* in the sharp Thread (on which Part the Female Screw bears) will shew the Friction to be less on the square Thread in Proportion to those Lines, besides the Obliquity to be consider'd elsewhere \*.

\* Pl. 11.  
Fig. 14.

\* Ann. 13.

66. THE Disadvantage to the Force of the Power occasion'd by the great Friction in the Screw is fully recompens'd in the Use of that Machine; because, on that Account, the Screw continues to sustain the Weight even after the Power is remov'd, or ceases to act, or to press the Bodies against which the Power had driven it; whereas in the Balance and Leaver and other mechanical Powers, the Weight ceases to be sustain'd and goes

Let III. back when the Power ceases to act. The Reason of this is, that  Weights when they are rais'd by Screws endeavour to descend perpendicularly, whereas the Screw has been push'd against them very obliquely, so that it can only be push'd back again in that oblique Direction, which cannot be given to them by the Gravity of the Weight tending downwards, if there be the least Friction against the Thread of the Screw: So likewise when a Body is press'd by a Screw, its Surface re-acts in Lines perpendicular to it, that is in the Direction of the Arbor of the Screw, whereas the Screw cannot be driven back unless it be mov'd in the Direction of its Thread, which makes a great Angle with the Axis of the Arbor. Thus the \* Plank K k *Fig. 10.* cannot fall down in the Direction of its Gravity towards M L, unless it could cause the Screws to move in the Direction B b or D d. Neither can the Teeth of the Wheel I \* *Fig. 13.* pressing against the Screw C c D in the Direction C B move the Screw or cause the Handle H to move in the Direction of the Circle whose Radius is A H, tho' the Power which went round in that Circle be remov'd, and the great Weight W does very strongly endeavour by the Axis E F to turn the Wheel I.

\* Plate II.  
Fig. 10.

\* Plate II.  
Fig. 13.

HENCE appears the great Use of the Machine of *Fig. 10.* which by Means of the strong Piece I may support the Side of an House, and several of them applied in different Parts can sustain a whole Building whilst the Foundation is mended or renew'd.

67. WHETHER the Screw be consider'd as a Wedge or an inclin'd Plane, it follows from what has been said, that the closer the Thread of the Screw is, the greater is its Force, and the more advantageous the Friction; the Power bringing forward the Screw with most ease, and the Difficulty to push back the Screw becoming the greater.

68. As Percussion is useful in the Wedge to lessen Friction; so in some Cases there is a kind of Percussion used to drive on a Screw, as in *Printing, Coining* and *Sealing* Instruments with large Seals where a great Pressure is requir'd; and that is done by a *Fly*, which is a Balance going thro' the Arbor of the Screw and loaded with Weights at its Ends, or sometimes only one *Brachium* of it, as in the *Printing-Press*.

HERE

HERE instead of the Hammer or Mallet, which comes on up-  
on the Back of the Wedge with an accelerated Motion in a  
circular Direction to drive it into the Wood; the Weight of  
the Fly comes down also with an accelerated Motion, but on  
an inclin'd spiral Plane, and thereby having overcome the Fric-  
tion of the Threads of the Screw as it descends, pushes the  
End of it with great Force against the Bodies to be press'd.  
*But this will be further explain'd hereafter in another Lecture.*

69. THE Friction of a Screw join'd by a Fly is often made  
use of to regulate Motion by retarding a Weight which would  
descend too fast, reducing the accelerated Motion of an heavy  
Body in its Descent to one that is uniform, by destroying just  
as much Force as Gravity would super-add to the descending  
Body in Motion. But we shall explain this more fully when we  
speak of the Fall of Bodies.

THO' Balances, Leavers, Pullies, Axes *in Peritrochio*, and even  
Wedges can be made to act upon one another to encrease the  
Force of the Power (as has been shewn in the Consideration  
of the Leaver, Pulley and Axis *in Peritrochio*.) Yet Screws can-  
not be applied to each other directly, without the Intervention  
of some other mechanical Power; but in Composition with o-  
ther mechanical Powers or simple Machines, the Screw will serve  
to make an Engine of vast Force. Which leads me to the Con-  
sideration of

### *Compound E N G I N E S.*

70. THE Combination of two or more of the simple Ma-  
chines, for the Uses of Life (whether they be of the same or  
of different Kinds) by means of a Frame of Wood or any Me-  
tal, makes what we call a *compound Engine*. As it would be  
endless to recount the several Sorts that are made use of, we  
shall only give an Account of some few, whereby we may  
judge of what may be performed by any Engine which has  
been made, or may be hereafter put in Execution, upon sight  
of an exact Draught of the intended compound Machine; that  
we may have a just Value for what is or may be useful, and  
not be deceived by Pretenders to Perpetual Motions, and those  
who promise greater Effects by Machinery than is conformable

to

Lect. III. to the reciprocal Proportion between the Intensities of the Powers and Weights of their Velocities \*.

\* Ann. 14.

Plate 6. 71. THE two Machines represented in *Fig. 1* and *2* of *Plate 6*.  
*Fig. 1, 2.* represent the Scorpions which the Ancients used in War for  
 \* L. 2. throwing Stones, as I have describ'd them already \*; I only  
 Ann. 7. take Notice here, that they are compounded of a forked Leaver, two Axes in *Peritrochio*, and two Pullies, which last in *Fig. 1.* only direct the Rope as they are Rollers, but double the Force of the Power in *Fig. 2.*

Plate 7. 72. THE Water Engine of *Plate 7. Fig. 14* and *15* consists  
*Fig. 14, 15.* of two Leavers E D and I K, and its Use is describ'd where it  
 \* L. 2. is first mention'd \*.


Ann. 9. at  
 the End.

73. THE 14th *Figure* of *Plate 9.* represents a Machine of great Force compounded of three Leavers acting upon one another as describ'd above \*.

\* Plate 9.  
*Fig. 14.*

74. THO' in the tenth *Plate, Fig. 4, 5, 6, 7* and *8* the Machines represented are made up of several single Pullies or Sheevers (Wheels moving round a Center-pin) like that of *Fig. 1.* yet as the same Rope runs over or under them all, they must be consider'd only as one Machine; for the Workmen call them by the single Name of a *Pulley* or a *Pair of Blocks*; and the Ancients consider'd a single or many Pulleys join'd in the manner represented in those Figures only as one Engine, which they call'd by various Names, according to its Number of Sheevers; as *Monopastum* when it had but one, *Dispastum* when it had two, *Trispastum* when three, *Tetrapastum* when four, *Pentapastum* when five, and usually *Polypastum* when it had many Sheevers.

\* Plate 10. 75. BUT the Machine of *Fig. 9* \*, is a compound Engine;  
*Fig. 9.* because in it the Pullies act upon each other and encreate the Force of the Power in a greater Proportion than according to the Number of Sheevers in the others abovemention'd, as has  
 \* 41. been shewn \*.

76. THE 13th *Fig.* represents an Engine compounded of two Axes *in Peritrochio*, whose Operation has been describ'd and Force calculated \*. 

\* 47.

77. THE 9th and 10th *Figures* of *Plate 11* \*, are compound Engines, because such great Screws cannot be turn'd without Leavers applied to them. N. B. *It is to be observ'd that when a Leaver or Leavers carry an Axis round, in which Case they are call'd Hand-spikes or Bars, they perform the Office of a Wheel turning an Axis, and may therefore together with the Axis they turn be consider'd as an Axis in Peritrochio.*

\* *Plate 11.*

*Fig. 10.*

THIS is more evident in observing the Motion of the Handle A H of *Fig. 13* \*, made use of to carry round the Screw and Axis A B.

\* *Plate 11.*

*Fig. 13.*


78. BECAUSE a Screw, which greatly adds to the Force of the Power, by reason of its Shortness, raises the Weight but to a small Height; and the Axis *in Peritrochio* (for the Reasons given in the Description of it \*) tho' it may raise the Weight to a very great Height, or from a great Depth, does not much encrease the Force of the Power; the Combination of these two Machines preserves the Advantages of each of them, and takes off their Defects: And this is done by causing the Threads of the Screw C D to take hold of the oblique or skew Teeth of the Wheel as c, and by continually turning the Wheel round to draw up a great Weight as W by means of the Rope which is wound on the Axis E F. If on the Axis E F another Screw be added, as at G H, to turn a second Wheel L, whose Axis M being of sufficient Length receives the Rope N instead of the Axis K O, a much greater Weight may be raised; or an Engine of vast Force may be made for boring hard Metals by means of cutting Instruments fix'd on the End of the Axis M, or performing Operations where much Strength is requir'd.

\* 46.

79. A Crane is an Instrument of such general Use, that we cannot avoid giving its Description here. It is of two Kinds: in the first, only the Gibbet moves upon its Axis, and in the second Kind call'd the *Rat-tail'd Crane*, the whole Crane, with its Load, turns upon a strong Axis.

80. THE

Lect. III. 80. THE first sort of Crane is represented by the 1st *Fig.* of  
 Plate 12. seen in Profil. L B E D is a Section of that Part of a  
 Wharfe on which it is fix'd, L B being the horizontal Line. A C is a strong horizontal Piece of Timber making the upper Part of the Crane, into which are fram'd the three upright Pieces X, Y, Z, (of which, the last call'd the main Piece, is stronger than the others) with its Cill I E, and its Braces H I and b E longer and stronger than the Braces and Cills M N and D S of the other two upright Pieces, and pinn'd with Iron where the others are only pinn'd with Wood. When the Wharfe is not of Stone-work where the Crane is fix'd (as is here represented) the three Cills must be all in one Piece reaching from D to E. Four Braces, such as K, join the upright to the horizontal Pieces. To the abovemention'd horizontal Piece is fasten'd with strong Iron Pins, a short Piece *p p* having a Bell-metal Collar to receive the Iron Pevet or Axis of the upright Shaft R F which is an Axis in *Peritrochio*, whose lower End of the Axis is also of Iron, turning in another Bell-metal Collar let into the firm Piece of Wood F. The Axis in *Peritrochio*, instead of a Wheel, has four Bars *e*; *f*, *d*, and another behind *d*, going thro' its thicker Part, which is eight square, the upper Part being round to receive the Rope. When this Piece is hoop'd with Iron above and below *d*, it is best to use but two Bars instead of four, pushing them quite thro' as *e b*, and Men at each end of them go round in the Direction *b O e* to wind up the Rope and raise the Weight at the End of it. This upright wooden Axle, with its Bars, is call'd the Capstane of the Crane (every such Axle being call'd a Capstane when it turns in a perpendicular Situation, like that of the Capstanes in Ships, and a Windless when it runs in an horizontal Position, tho' it should be for the same Purposes as the other) and the Rope R r r, which goes first over the Pulley or Roller T, then between the Pulleys P and Q; and lastly over the Pulley r has at its Ends a double Iron-hook, call'd a Ram's-head, to which the Goods to be cran'd up are fasten'd. The Gibbet G V B is moveable upon its Axis C B, by means of the Iron Center-pins or Pevets at its Ends B and C, so that when the Weight is rais'd up sufficiently high, by a small Rope fasten'd to it, or to the End of the Gibbet at g, it may be easily brought from over the Ship or Barge, suppos'd on the Water at W, to a Cart, or any other Carriage on the Wharfe towards

wards *w* on the right or left of the Piece *Z*. There is a Roof *Le*ct. III. or small wooden Shed *A a Q* to shelter the Rope from Rain,  when the Crane is not in use, the Gibbet being brought under it towards *Y*.

THE second Figure \* shews the Plane of the upper Part of the \* Plate 12. Crane, or as it would be seen from above, where we are to ob- Fig 2. serve the Position of the Pulleys *P* and *Q*, and of the Place of the Center of the Gibbet which must be at *C*, in a Line touching the Circumference of both the Pulleys; for if the said Center of Motion of the Gibbet were in a Line with the Center of the Pulleys, the Gibbet when loaded would require a Force to bring its End *g* over the Wharfe on either Side, and that Force ceasing to act, the Weight and Gibbet would run back and rest over *W*. See this more particularly explain'd in the Notes, where the 3d Figure, being Part of the Crane and various Situations of the Gibbet and Rope drawn by a larger Scale, is consider'd \*. Ann. 15.


THIS Crane is very expeditious with many Hands, it being always requisite that some should stand at the Bars to keep the Weight from running down again, which might be of dangerous Consequence. But if instead of the Capstane at *O*, there was an Iron endless Screw, and horizontal Wheel and Axle (commonly call'd a Worm and Wheel) fix'd to the Piece *X* made very strong for that Purpose, or a Machine in the manner of Fig. 13. of Plate 11. only with a Pinion instead of a Screw at \* Pl. 11. *G H*, making the Axis *E F* short and the Axis *M* long to re- Fig. 13. ceive the Rope; then two Men, nay sometimes one Man, may easily draw up Goods from a Barge; because by the Resistance of the Screw against the Teeth of the Wheel the Machine will hold the Weight at any Height, whilst the Man quits the Handle to bring the Weight by the Guide-rope fasten'd at *g* \* over a \* Plate 12. Cart on the Wharfe to receive it; and yet the Weight will go Fig. 1. down gently of it self, if a Man with a sudden Jerk sets the Handles a going the contrary way to that whereby he rais'd the Weight which is convenient for loading the Cart. N. B. *This sudden Impulse on the Handles does to the Screw what a Blow does to the Wedge to set it a going* \*. As this last sort of Crane \* No. 57. with a Worm and Wheel does very much encrease the Force of the Power, I need not put the Reader in mind that more Time must be spent in raising the Weight; because the Velocity of the Weight, compar'd with that of the Man's Hand who

R turns

Le<sup>ct</sup>. III. turns the Handle, must be diminish'd in a reciprocal Proportion  
 ~~~~~ of the Intensity of the Weight to the Intensity of the Man's  
 * L. 2. 9. Force *. This being true in all compound as well as simple
 12. 17. Engines.

* Plate 12. 81. THE fourth Figure * represents the *Rat's-tail Crain*, not
 Fig. 4. only useful on a Wharfe to crane up heavy Goods, but also of great Service in Building to raise great Stones and bring them round to any design'd Place. It consists of the following Parts. On the Cross-ground Cills L L L L L L is supported and fix'd by oblique Braces the strong upright Piece K call'd the Gudgeon of the Crane, on whose upper Part or Spindle S cover'd with Iron (and sometimes made wholly of Iron) the whole Machine turns, being easily mov'd from C to G, when it is charg'd with its Load H. C A is the Counter-wheel with its Axis D B bearing only on the Iron Ends of the said Axis in two hanging perpendicular Pieces at B and b; f F is the Brace and Ladder, whose Top F carries the Pulley above the Weight, the other Pulleys being in the Ends of the Pieces M, N, E; the remaining Parts are too plain in the Figure to need farther Explanation. The Power is sometimes applied by means of a Rope on the outer Circumference of the Wheel A, but most commonly Men, or an Horse, or an Ass, turn the Wheel round by walking in it. Sometimes also Strength is gain'd by having the Counter-Wheel made with Teeth, and giving Motion to its Circumference by Means of a Pinion.

* Plate 12. 82. THE 5th Figure * represents a Jack, which is an Instru-
 Fig. 5. ment of common Use, for raising heavy Timber or very great Weights of any kind; but as the Wheel-work of it is shut up in the strong Piece of Timber C B, I thought it not improper to represent the Inside of it in Fig. 6. * where you must only
 * Plate 12. suppose the Rack A B at least four times as long in Proportion
 Fig. 6. to the Wheel Q (the Figure of the Rack being here contracted for want of Room) and the Teeth, which will then be four times more in Number, to be contain'd about three in an Inch. Then if the Handle be seven Inches long, five Turns of it, that is five times 22 Inches, or 110 Inches will be the Velocity of the Power, whilst the Weight rais'd by the Claw A, or depress'd by the Claw B moves one Inch; for as the Pinion has but four Leaves, and the Wheel Q twenty Teeth, there must be five Re-
 volutions

volutions of the Handle fix'd to it to turn the said Wheel once Lect. III.
 round, whose three-leav'd Pinion R will in that Revolution just 
 move the Rack three Teeth, or one Inch. This might have
 been also known without seeing, or even knowing the Num-
 bers of the Teeth of the Wheel and Pinions, by measuring a Re-
 volution of the Handle in *Fig. 5.* and comparing the Space gone
 thro' by it with the Space gone thro' by the End A or B. So
 in any other compound Engine, we may judge of its Force by
 comparing the Velocity of the Part to which the Power is ap-
 plied with the Velocity that moves the Weight, as has been
 before shewn *. Sometimes this Machine is open behind from * 32, 47.
 the Bottom almost up to the Wheel Q, to let the lower Claw
 (which in that Case is turn'd up as at B) draw up any Weight.
 When the Weight is drawn or push'd sufficiently high, it is kept
 from going back by hanging the End of the Hook S fix'd to a Plate 12.
 Staple over the curv'd Part of the Handle at *b.* All the Work Fig. 5.
 must be made very strong in this Machine, but chiefly those
 Parts which immediately sustain the Weight; and this must also
 be observ'd in all other compound Engines.

83. ALL the Mechanical Powers are united together in the
 Machine represented by the first Figure of *Plate 13* *, which Plate 13.
 resembles our common Chimney Jacks for roasting Meat. In a Fig. 1.
 Frame A B C D fasten'd by the Nut *o* upon the Stand *o* O, and
 held together by the Pillars V W and B *q*, is adapted first the
 Piece E F, whose Fans or Flies may be put in Motion by the
 Wind, or drawn by an Hair fasten'd at F, which represents the
 Leaver and Balance : At right Angles to this Piece is joyn'd the
 perpendicular Spindle G H having upon it the endless Screw H,
 which may be also consider'd as a Wedge * : This endless Screw * 63.
 or Worm takes the skew Teeth of the Wheel K, which is the
Axis in Peritrochio, and, in turning round, winds up the String
 L M upon its Axis, which passing round the Pulleys at M and
 N (or drawing by a *Tackle of Five*) raises the Weight P. But as
 the Screw has no progressive Motion on its Axis, it cannot here
 be said to take in the inclin'd Plane; therefore to make this
 Engine take in all the mechanical Organs or Powers abovemention'd,
 we may add the inclin'd Plane *r q* Q R by making it
 rest on the Ground at Q R, and on the Pillar *q* B at *q r*, and
 thereby the Force of the Power drawing at F will be farther
 encreas'd in the Ratio of Q T to T S *. The whole Force gain'd * 49.

LECT III. by this Machine is found by comparing the Space gone thro' by the Point F, with the Height that the Weight is rais'd in any determinate Number of Revolutions of F *. An hundred Pounds Weight at P will be easily rais'd by the Hair of a Man's Head drawing at F.

* 32. 47.
82.

* 14.

* 68.

84. IN the Machines abovemention'd, both simple and compound, the Power according to its Intensity is so applied to one Part of the Machine as immediately to act upon the Weight whose Resistance destroys all the Force of the Power, when an *Æquilibrium* is made by giving the Body mov'd and the Mover a Velocity reciprocally proportionable to their Intensities; and when the Product of the Power into its Velocity exceeds that of the Weight into its Velocity, there is no more *Momentum* left to the Power, but so much as it has more than the Weight *. But there are other mechanical Organs, in which the Force of the Power is accumulated, and, as it were, *condens'd*, before the Weight is acted upon at all; such as an Hammer, the Cock of a Firelock which carries the Flint to strike upon the Steel, the Battering-ram (already mention'd by the by in another Lecture) the Rammer to drive Piles into the Ground, the Fly applied to Printing or making a Stamp on Medals or current Coin (cursorily taken notice of in this Lecture *) and the Pendulum; and in a word, whatever is made use to give a sudden Blow, or strong Impression instantaneously; which tho' all reducible to geometrical Calculation, whereby one may be certain of their Effects, yet cannot be explain'd from the Principles already given; and therefore the Consideration of them must be defer'd till we have explain'd those Laws of Motion on which they depend. In the mean time I think it not improper to give Sir *Isaac Newton's* Account of all the mechanical Powers by one Scheme, as he has explain'd them in the second Corollary to his Laws of Motion; for tho' that Explanation depends upon such Laws as are to be hereafter consider'd, so much of them as is contain'd in his first Corollary being taken for granted, will serve our present Purpose.


85. If upon a Body * at A two Forces act at once, whose Intensities are as the Length of the Lines AB and AC, and Directions in those Lines; the Body so acted upon will describe the Line AD the Diagonal of the Parallelogram completed by draw-

drawing the two Lines CD and DB respectively equal and parallel to the two former; and that Diagonal will be describ'd by the joint Forces in the same Time that either of the Forces singly would have caus'd the Body to go thro' the Line AB or AC . And as by Composition of Forces a Body will move in the Line AD ; so likewise a Body moving in the Line AD , tho' by the Action of one single Force, may be consider'd as if it had been acted upon by two Forces, namely, resolving the single Force into two, as AC and AB .

86. IF the unequal Radii OM^* and ON drawn from the Center O of any Wheel should sustain the Weights A and P by the Cords MA and NP ; and we wou'd know the Forces of those Weights to move the Wheel. Through the Center O draw the right Line KOL , meeting the Cords perpendicularly in K and L ; and from the Center O with OL , the greater of the Distances OK and OL , describe a Circle, meeting the Cord MA in D : and drawing OD , make AC parallel and DC perpendicular thereto. Now, it being indifferent whether the Points K, L, D , of the Cords be fix'd to the Plane of the Wheel or not, the Weight will have the same Effect whether they are suspended from the Points K and L , or from D and L . Let the whole Force of the Weight A be represented by the Line AD , and let it be resolv'd into the Forces AC and CD ; of which the Force AC , drawing the Radius OD directly from the Center, will have no Effect to move the Wheel: But the other Force DC , drawing the Radius DO perpendicularly, will have the same Effect as if it drew perpendicularly the Radius OL equal to OD ; that is, it will have the same Effect as the Weight P , if that Weight is to the Weight A as the Force DC is to the Force DA ; that is (because of the similar Triangles ADC, DOK) as OK to OD or OL . Therefore the Weights A and P , which are reciprocally as the Radii OK and OL that lie in the same right Line will be equipollent, and so remain in *Æquilibrio*, which is the well known Property of the Balance, the Leaver and the Wheel. If either Weight is greater than in this Ratio, its Force to move the Wheel will be so much the greater. If the Weight p equal to the Weight P , is partly suspended by the Cord Np partly sustained by the oblique Plane pG ; draw pH, NH , the former perpendicular to the Horizon, the latter to the Plane pG ; and if the Force of the

Lect. III.


* Plate 12.
Fig. 3.

Lect. III. the Weight p tending downwards is represented by the Line pH ,  it may be resolv'd into the Forces pN , HN . If there was any Plane perpendicular to the Cord pN , cutting the other Plane pG in a Line parallel to the Horizon; and the Weight p was supported only by those Planes pQ , pG ; it would press those Planes perpendicularly with the Forces pN , HN ; to wit, the Plane pQ with the Force pN , and the Plane pG with the Force HN . And therefore if the Plane pQ was taken away, so that the Weight might stretch the Cord, because the Cord now sustaining the Weight, supplies the Place of the Plane that was remov'd, the Force of that Weight will be obtain'd by considering it as the same Force pN which press'd upon the Plane before. Therefore the Tension of this oblique Cord pN will be to that of the other perpendicular Cord pH as pN to pH . And therefore if the Weight p is to the Weight A in a Ratio compounded of the reciprocal Ratio of the least Distances of the Cords pN , AM , from the Center of the Wheel, and of the direct Ratio of pH to pN ; the Weights will have the same Effect towards moving the Wheel, and will therefore sustain each other, as any one may find by Experiment.

BUT the Weight p pressing upon those two oblique Planes, may be consider'd as a Wedge between the two internal Surfaces of a Body split by it; and hence the Forces of the Wedge and the Mallet may be determined; for because the Force with which the Weight p presses the Plane pQ , is to the Force with which the same, whether by its own Gravity, or by the Blow of a Mallet, is impelled in the Direction of the Line pH towards both the Planes, as pN to pH ; and to the Force with which it presses the other Plane pG , as pN to NH . And thus the Force of the Screw may be deduced from the like Resolution of Forces; it being no other than a Wedge impelled by the Force of a Leaver. Therefore the Use of this Corollary spreads far and wide, and by that diffusive Extent the Truth is farther confirm'd. For on what has been said depends the whole Doctrine of Mechanics, variously demonstrated by Authors. For from hence are easily deduced the Forces of Machines, which are compounded of Wheels, Pulleys, Leavers, Cords and Weights ascending directly or obliquely, and other mechanical Powers; as also the Force of the Tendons to move the Bones of Animals.

End of the THIRD LECTURE.

ANNO-

ANNOTATIONS upon the Third LECTURE.

1. [1. *Simple Machines, or Organs, call'd by some Mechanical Faculties, or Mechanical Powers.*]

THE Word *Power* here is to be taken in a very different Sense Annotat.
 from what it bears in the Second Lecture, where the Word Lect. III.
Power * (signifying whatsoever is made use of to raise a Weight) L. 2. No. 19.
 is defin'd at large; for here it only signifies the Organ or Instrument
 whereby a Power of known Intensity is made to act upon a Weight;
 and therefore we must take care not to attribute any real Force to any
 simple or compound Machines, as a great many People are apt to do mere-
 ly because the Name *Power* has been given to Mechanical Organs, not
 from their Effect, but from the Effect which the Power produces by their
 Means; for how much soever the Force of a Power is thereby encreased
 in order to sustain or raise a Weight far superior to it in Intensity; yet
 this cannot be done without losing in Space and Time what is gain'd in
 Force, contrary to what some have vainly imagined, because the Vulgar
 commonly speak of a Machine as they do of an Animal, and attribute
 that Effect to the Machine, which is the Effect of the Power by means
 of the Machine, as it is usual to say, *Such a Machine raises such a Quan-
 tity of Water, or performs such and such Work*; when we should say, if
 we would speak in the proper and Philosophical Sense, *Such a running Stream,
 such a Fall of Water, the Wind, or so many Men, Horses, or Oxen, &c. raise
 so much Water in such a Time, &c. by such or such a Machine*; as we have
 observ'd in the Notes on the last Lecture*. It were therefore to be * Ann. 6.
 wish'd, that the Word *Power* were confin'd to its proper Sense, and not L. 2.
 us'd to signify a Mechanical Organ; but as it has been customary to use it
 in that Sense, I thought proper also to make use of it; but withal to give
 this Caution.

2. [3. — *reducing all the Mechanical Powers to the Lever, or explain-
 ing all their Operations by that of, &c.*] Tho' one may easily shew (as we
 shall at the End of this Note) that every other Mechanical Organ does
 potentially contain a Lever; that is, that if so much of every other sim-
 ple Engine was cut away as to leave only a Lever, the same reciprocal
 Propor-

Annotat. Proportion between the Velocities of the Power and Weight and their
 Lect. III. Intensities; would as plainly appear on such a Leaver as on the Engine,
 before it is reduced to it: Yet we are not from thence to conclude, that the
 Leaver alone might be made to serve for the Purposes of all other Mechanical Organs; for they are contriv'd of different Forms to answer to different Ways of working, requir'd in Mechanical Operations for the Uses of Life, it being sometimes impossible to use one Mechanical Organ instead of another; and always more convenient to apply one than another, the Choice of which depends upon the Skill of the Artist.

Thus in the Balance (whether a Pair of Scales or a Steel-yard) the Commodity to be bought, or heavy Body whose Quantity of Matter is to be estimated, is not to be rais'd with any Velocity, but only supposed to make an *Æquilibrium* with a known Weight, which in this Machine serves for a Power: But a Weight is seldom or never made use of as a Power in any other Mechanical Organs, except in some few, and those Engines which the Ancients made use of in War, which are now out of Use.

The Force of one or more Men is the Power applied to the Leaver; there the Power must always overcome the Weight, by adding a little more Velocity or a little more Intensity to the Power, over and above the reciprocal Proportion requir'd in the Balance. With this Instrument heavy Bodies are removed a little way at a time, as great Stones in Building, large wooden, leaden, or iron Pipes in Water-works, and large Pieces of Timber; but Leavers will only serve to raise those Bodies high enough to lay them on a Carriage, &c.

But if we wou'd raise up a Stone to a considerable Height to lay it in its proper Place in a Building, or any other Body to any Height above three or four Foot, the Leaver becomes ineffectual, and then we must use
 * 37, 38, 39. Pulleys in some of the Ways mention'd in this Lecture *. Tackles of Pulleys are very convenient where there is no room for a Capstane, and where Bodies are to be rais'd in different Places, because they are easily moveable; but the Weight must not be too great, by reason that many Men cannot pull at once, and equally, by one running Rope; and if the Power was to be very much encreas'd by the Number of Sheevers or Wheels in the Pulleys or Tackles, the Rope must be of a prodigious, and therefore inconvenient, Length.

An Axis in *Peritrochio*, Capstane or Windless, which are all the same Organ differently posited, is of use where Pulleys are deficient. For Example, if Water is to be drawn out of a deep Well, a Wheel with Arms or Spokes serves to turn the Axle on which the Rope winds to bring up the Bucket or Buckets. In Building, a Capstane whose Make gives the Power no more Advantage than a Tackle or Pulley of many Sheevers, is yet more useful, because it will admit of eight, ten, or twelve Men to work at it and push its Bars, when only three or four could pull at the running Rope of the Pulley. If the four Bars of the Capstane are so long that three Men applying their Strength to each of them, the Man in the middle of the Three is at the Distance of three Foot from the Axis of Motion,

Motion, and the Axle on which the Rope winds is of six Inches Diameter, those twelve Men will do the Work of 72, but in the sixth Part of the Time. The same would be done, and in the same Time, by two Men walking within a vertical Wheel of 24 Foot in Diameter (such as is call'd by some a Counter Wheel) whose horizontal Axle might be of eight Inches Diameter; and about the Weight of one Ton and an half, or two Tons, might thus be rais'd by either of these Machines. But if at a Wharfe, where only two Men should be employ'd to work at the Crane, it was requir'd to draw up heavy Blocks of Marble of three or four times the Weight abovemention'd (for a Block of Marble six Foot long, four Foot wide and four Foot deep will weigh between 7 and 8 Ton) the Counter-wheel must be 72 Foot in Diameter for an Axle of eight Inches, which is impracticable, on account of the Bulk and Cost; or the Axle must be made three times less in Diameter, and then it cannot be strong enough to bear the Weight. In such a Case therefore there must be a compound Axis in *Peritrochio*, such as is describ'd in this Lecture*, but with one more Pinion; for example, if an Iron Wheel with Teeth of four Foot in Diameter be fix'd to the abovemention'd Axis of eight Inches Diameter, and that Wheel be lead by an Iron Pinion of six Inches Diameter, whose Wheel is of three Foot Diameter and lead by a Pinion of eight Inches Diameter, whose Handles are a Foot long, the same Operation will be perform'd by two Men, but they must employ three times more Time. N. B. *If the Teeth of the Wheel be of Brass, and the Teeth or Leaves of the Pinions of Iron, the Machine will move more smoothly, and the Teeth wear equally.*

Annotat.
Lect. III.
~

No. 47.

The Numbers of Teeth in the Wheels and Pinions may be as follows,

The first Pinion, to which the Handles are fix'd, 28.


The first Wheel, led by the said Pinion, 112.

The second Pinion, 19.

The second Wheel, 171.

And the Axis on which the Rope winds must be of about eight Inches, because it immediately bears the Weight by the Rope, which runs over some upper Pulleys or Rollers that add no Force to the Power.

If the Capstane with Bars abovemention'd, be fix'd to do the Work for which it is proper, and unexpectedly there should be occasion to lift very heavy Weights, there is no Necessity for taking it down to set up such a Combination of Wheel-work as we have just describ'd, because a Pair of Blocks to help at such a Time may be fix'd in any Part of the Building over the Place where the Weight is to come up and receive the Rope from the Capstane, so as to encrease the Force of the Power, according as the Blocks are a *Tackle of Two, of Three, of Four, or Five, &c.* and then the Blocks may be taken off again, and the Capstane work'd as before. In the Use of the Counter-wheel or Rat's-Tail Crain * the Power may be al- * No. 82. so thus occasionally encreas'd.

Annotat. Where there is no Room for the compound Axis in *Peritrochio* of two
 Lect. III. large Wheels and Pinions, the same may be affected by making a Worm-
 screw of one Thread turn'd by two Handles of a Foot long each, to lead
 a Wheel of two Foot Diameter, having 72 Skew-Teeth, and a Wooden
 Barrel or Axle of 8 Inches Diameter, the Advantage of which Machine is
 * No. 79, 81. shewn in this Lecture *.

If Hogsheds or Pipes of Wine, or other Liquors, are to be let down
 into a Cellar, or brought up out of it, a Plank is laid along the Stairs,
 which in that Case is an *inclin'd Plane*, the only mechanical Organ fit for
 that Purpose: So likewise in making Reservoirs for Water, in gardening,
 and in building Fortifications, where Carts can't come, inclin'd Planes made
 of Wood serve effectually for Wheel-barrows to run on in removing the
 Earth from a lower to an higher Place.

In cleaving Wood, the Wedge only is of Use; for an Hatchet, which
 may cleave small Wood, is only a Wedge with an Handle. Another con-
 siderable Use of a Wedge, is the raising up a Beam to underprop it when a
 Floor begins to give Way, by reason of too great a Burthen laid upon it,
 as in a Ware-house; and so much Force may be applied this Way, that
 some thousand Tons may be rais'd up together with the Floor, and all
 secur'd by means of this small Machine. For tho' Screws turn'd by long
 Leavers might serve in a great measure for the same Purpose, there must
 be Space to go round with the Hand-Spikes, which cannot be had when
 the lower Part of the Ware-house is full of Goods, without removing
 them with great Trouble and Cost. See the Manner of performing this
 in the 4th Figure of Plate 13.

Plate 13.
 Fig. 4.

BADEC is a Beam, which in the horizontal Situation, mark'd by
 the prick'd Lines Bb and CD supports a Floor. Now when very great
 Weights being laid upon the said Floor, make it give way and bend into the
 Curve BAb, or, which is the same, CED, an upright Post Pp is fram'd
 into a stout Plank FF, and an horizontal Piece E (here seen End-wise) is
 slip'd under the Beam, and on the Top of the Post; then the two Wooden
 Wedges W, w, as broad as the Post, are driven in with heavy Hammers,
 in striking at once in contrary Directions, so as to set the Beam streight,
 and reduce the Floor to its Place, without removing any of the Goods that
 lie on the said Floor, or moving any of those below, but just to make
 Room for the Post and Plank.


The Uses of the Screw for raising, sinking, drawing, pushing, pressing,
 or joining together Bodies, are commonly known; and it is evident by In-
 spection, that no other of the mechanical Organs will answer the same Pur-
 poses.

Now we will shew how all the mechanical Organs or Instruments may be re-
 duc'd to the Leaver.

This has been already shewn concerning the Roman Balance (and is ap-
 plicable to the common Pair of Scales) by the 6th Experiment of this

* No. 29.
 Pl. 9. F. 11. Lecture *

Secondly,

Secondly, Pulleys are thus reduc'd to the Leaver *. An upper Pulley as Annotat. E D (Plate 10. Fig. 3.) appears plainly to be a Leaver of the first Kind, Lect. III. by cutting away all the Sheever, except the prick'd Line E D, which about the Center C keeps in *Æquilibrio* two equal Weights. In the 4th Figure  * Plate 10. Fig. 3. the lower Pulley *ge*, by leaving only the prick'd Line *ge*, and Center Pin *c*, appears to be a Leaver of the second Kind, whereby a Power applied at *g*, does in the Direction *gd* raise a Weight *W* suspended at the Center Pin *c*. It is here evident, that *e* is the fix'd Point or *Fulcrum* of the Leaver, *ge* the Distance of the Power, and *ce* the Distance of the Weight; and accordingly, in the Experiment, the Power *P* : is to the Weight *W* :: as *ce* : to *ge* :: 1 : 2. And by comparing the compounded Leavers of Plate 9. Fig. 14. with the System of Pulleys of Plate 10. Fig. 9. one may see that the four Pulleys reduc'd to their horizontal Diameters, act upon one another as four Leavers of the second Kind, in every one of which the Distance of the Power is Two, and that of the Weight is One, and therefore the Ratio compounded of them all *, is that of the Weight to the Power, or 16 to 1; for $2 \times 2 \times 2 \times 2 = 16$. Or comparing this System of lower Pulleys with Fig. 5. of Plate 13, which is just such a System or Assemblage of Leavers of the second Kind, where the Leavers are mark'd with the same Letters, as well as the Forces pulling down each Leaver *. * No. 32.

One may, by the 11th Figure of Plate 10, easily reduce an Axis in *Petrochio* to a Leaver of the first Kind, represented by the prick'd Line E T, the fix'd Point being at K, the Power being applied at E, and different Weights successively at the Points A, B, and T, cutting away the rest of the Machine. But as the Rope sustaining the Weight does not move in the same Plane as the Rope drawn by the Power, it is better to consider it as a Leaver of the first Kind twice bended, and an Axis of Motion going thro' one of the bended Parts, as in the 6th Figure *, where the bended Leaver A C c B moves on the Axis II fix'd in the Frame I K L I. *Bc* represents the Radius of the Axle, and AC the Radius of the Wheel, supposing *Bc* and AC in the same Plane, and at right Angles to the Axis; otherwise, if oblique, they must be reduc'd to right ones, by calling their Lengths only the perpendicular Distances of B and A from the Axis II: Then *P* being the Power, and *W* the Weight, the reciprocal Proportion will be thus, A C : B c :: W : P. * Pl. 13. F. 5.

To reduce the inclin'd Plane to a Leaver, we must look for a bended Leaver in the Weight rolling up the Plane, whose Arms shall be as the Length of the Plane to its Height.

Since the Triangle ABC* is similar to *w Y B* (by 4. 6. *Euc.*) and *w Y B* similar to *w B N* (by 8. 6.) in the bended Leaver *w B N*, *w B* : *BN* :: *AB* : *BC*. Now, since *w N* is the Line of Direction of the Weight *w*, that Weight may be consider'd as pressing on NB the short Arm of the Leaver at the Point N, the Center of Motion being at B, where the spherical Weight touches the Plane, and the Power applied at right Angles at the End *w* of *w B* the longer Arm of the Leaver; therefore, calling *P* the Power, and *w* the Weight, *P* : *w* :: *N B* : *B w* :: *AB* : *BC*.

Annotat.
Leſt. III.



N. B. Here the Power by a Rope over the Pulley M, draws in a Direction parallel to the Plane; but if it drew in any other Direction, one might calculate the oblique Force of the Power by means of a bended Leaver. But we must refer this to our particular Observation of the inclin'd Plane, and ſtatical Conſiderations, in other Notes; except the Direction parallel to the Baſe, which reduces the Wedge to a Leaver.

Now the Triangle ABC muſt represent the Wedge, which being driven under the Weight, makes it riſe up the perpendicular Height CB, while the Power drives the Wedge the Length of its Baſe AC, or, which is the ſame Thing, the Power Π draws in the Line wr parallel to the Baſe AC. Here the bended Leaver is OBN, with the ſhort Arm NB ſupporting the Weight at N, while the Power faſten'd to O draws the Arm OB at right Angles.

Authors have ſhewn other Ways of reducing the Wedge to a Leaver:

* Plate 13.
Fig. 7.

For example, the Wedge BFC* of Plate 13. Fig. 7. is conſider'd as a Leaver of the ſecond Kind, whoſe Fulcrum is at F, and the Weight at W; which moving round the Center of Motion F, by a Power carried on at the End of the Leaver from A to L, raiſes up the Weight W. Or elſe, the Leaver being kept by the Prop B, ſo as always to make the ſame Angle with the Horizon, is carried on from the Poſition CFB into the Poſition cfb , the Fulcrum advancing forwards along with it to raiſe the Weight to w . In the 8th Figure*, where two Bodies are ſeparated to

* Plate 13.
Fig. 8.

represent the cleaving of Wood, the Wedge is reduc'd to two Leavers of the firſt Kind, having the Fulcra at F, f , the Weights at W, w , and

* Plate 13.
Fig. 9.

the Powers at L, l . Or in the 9th Figure*, two Leavers of the ſecond Kind are ſuppoſed to be very thin, and thruſt in between the two Weights, ſo as to preſs againſt each other's Ends, and make a common Fulcrum at F, the Powers moving from L, l , to a and b , while the Weights W, w , are ſeparated: Or elſe (what comes to the ſame Thing) the ſaid two Leavers being join'd in a fix'd Angle by the Prop L l (Fig. 10.*) are conſider'd

* Plate 13.
Fig. 10.

as thruſt in between the Weights, while the common Fulcrum advances in the Line FG. But becauſe in all theſe Ways of explaining, the Diſtance of the Power (and conſequently its Force) is continually changing, which is not true in the Wedge; I would rather propoſe another Method, which will agree with the Caſe of the ſingle Wedge or double Wedge (diſtinguiſh'd thus, as the Inſtrument acts with one or both Surfaces*) always making the Diſtance of the Power and Weight to keep the Proportion agreeable to the Angle of the Wedge. As for example, in the Wedge

* No. 55, 56.

* Plate 13.
Fig. 11.

LFW Fig. 11.* LFW is a bended Leaver, whoſe ſhort Brachium always remaining the ſame, lifts up the Weight x , and brings it to w , whilst the Leaver turns round the Center F, the Power at L deſcribing the Arc L l : So in the double Wedge Fig. 12. two bended Leavers, moving round the Center F, by their ſhort Arms FW, F w , ſeparate the Weights W, w ; and when they have brought them round to X and x , the long Arms LF, lF turning round F in the Arcs LM, lM , one may ſee that the

the same Thing is perform'd as if the whole Wedge FM had been driven between them in the Direction NM. Annotat.
Lect. III.

Having shewn that the Screw acts either as an inclin'd Plane, or as a Wedge, it is evident that what has just been said reduces it to a Leaver.

3. [—*The Quantity of Friction in Engines will be consider'd, &c.*] That there is a Loss of Force in the working of Engines on account of the Rubbing or Friction of their Parts, has been observ'd by most Writers of Mechanics; but that Friction has not been enough consider'd by them: Upon which Account, several Persons (who having applied themselves to the Study of Mechanics are not yet much acquainted with the Practice) imagine by comparing the Effects which are perform'd by means of such Engines as they examine with the Powers apply'd to them, that the Machine must be very ill contriv'd, because the Effect does so much differ from the Calculation which they make exclusive of Friction; supposing indeed that something is to be allow'd on that Account, but nothing near what they find to be lost in Force. Possess'd with this Notion, Projectors contrive new Machines (new to them, tho' perhaps describ'd in old Books, formerly practis'd and then diffus'd and forgot) which they suppose will perform much more than they have seen done with the same Power; because they allow too little for Friction. Full of this they go to the Charge of 70 or 80 *l.* for a Patent for their new Invention; then divide it into Shares and draw in Persons more ignorant than themselves to contribute towards this (suppos'd advantageous) Undertaking; till, after a great deal of Time and Money wasted, they find their own Engine worse than others which they hoped by many degrees to excel. This has been very much the Practice for these last twenty Years: For tho' some Projectors have been altogether Knaves; yet the greatest part have first deceiv'd themselves; and those who are really deceiv'd, by their Eagerness and Earnestness most easily deceive and draw in others. For this reason, I thought it would be of Use to the Publick to give as full an Account of Friction, as I possibly could gather from the Experiments made by others (especially the Members of the Royal Academy at *Paris*) and my own Experiments and Observations.

We shall not here consider, that Friction or rather Resistance which arises from an ill Contrivance of the Parts of an Engine which are to act upon one another by Application; making those Parts act obliquely which ought to act at right Angles, or at least more obliquely than they should do; that being owing to an incomplete Theory or a bad Workman. But only that Friction, which is unavoidable from the Nature of the Materials, however polish'd, at the first making use of an Engine; and that which Time brings on, as the Parts wear unequally, or grow rusty or rotten for want of Oil, Grease, or by constant Use: So that the touching Surfaces, which were as smooth as the Hand of a skilful Workman could make them, become very rough and uneven by this means, and add much Friction to that which, on account of the Nature of the Materials, could not be avoided at

Annotat. at first. So an old Jack for roasting Meat requires more Weight to make
 Lect. III. it go when the Pivots and their Holes are pretty much worn ; and a well
 made Lock grown rusty for want of Oil to guard it from the Action of
 the acid Salts in the Air will not without difficulty be open'd with a rusty
 Key ; tho' the Figure both of the Lock and Key be as perfect as at first.
 Wood will grow rotten, and swell beyond, or shrink from, its first Dimensi-
 ons by the Weather ; and Cords run round Pulleys or wind about Rol-
 lers with more difficulty according as they stiffen by wet, or any way be-
 come more twisted.

To proceed methodically, we will consider the unavoidable Friction of the simple Machines or mechanical Organs severally.

The Leaver, in respect to the Work done with it, is subject to very little Friction, moving on a small Surface crossing the Instrument like a Line where it is applied to the *Prop* or *Axis of Motion* ; which in Theory is consider'd only as a Point, and call'd the *Center of Motion*.

When the Balance (whose Make is nearest to that of the Leaver) has no more Friction in proportion to its Length than the Leaver ; yet it has a great deal too much for a nice Balance, as may be found by Experience. Few of the Scale-makers know wherein the Nicety of a Balance consists ; but generally follow a Fashion, or, when they would excel, endeavour to out-do one another in Ornaments or a fine Polish, consulting Beauty more than Use ; and thinking the Business is done if they have brought the Balance to turn with a small part of a Grain.

For the Benefit of those who would make, and those who would use very exact Scales, I shall here mention the Faults I have found in Scales esteem'd extraordinary good ones ; and shew in what manner I think such Faults may be avoided ; as also the utmost that may be expected from a well made Balance.

Mr. George Graham and I last Summer, in order to try some Experiments with Brigadier Armstrong, Surveyor of his Majesty's Ordnance, examin'd a Pair of Scales made by a very nice Workman, and kept in a Glass Case that the Air might have no Power over them. These Scales were thought extremely nice, because they turn'd with the 256th part of a Grain ; but upon Examination they appear'd to differ from that Nicety sometimes 3 of the Parts above-mention'd, which made us imagine one *Brachium* longer than the other ; but at last we found that this Difference was owing to the Situation of the Axis of Motion ; for if in the Circle *A a B b** (on the lower part of which the Axis or sharp Edge of *C* ought to bear at *A*) the said Edge should not rest at *A*, but at *a* or *b*, the *Brachium* on the other Side of *A* will preponderate ; which it did in our Trials, so as to make a Difference three times greater than what turn'd the Balance in its true Situation. Now since the Friction encreases in proportion to the Weights bearing on the Axis of Motion (as we shall hereafter shew) this Error will encrease in the same manner and become considerable, as heavier Bodies are weigh'd ; so that when we think we have an *Æquilibrium*, by only lifting up the Beam *EL* with the Weight hanging at its
 Ends

* Plate 13.
 Fig. 13.

Ends in the Scale, we shall upon letting it down again find the *Æquilibrium* to be lost; so difficult it is to lay the Edge or Axis of Motion upon the same Place which it had before, or to give it a true Bearing at first over the lower Part of the Ring at A. There is also another Fault which Scale-makers are sometimes guilty of, which is not to make the *Brachia* nicely of a length; and then to hide that Error, they adjust the Balance by filing away some of the Thickness of the longest Part of the Beam, and sometimes by the Scales. Others again, by making Ornaments in a nice Balance weaken a slender Beam just under the Axis where it ought to be strongest.

Annotat.
Lect. III.

To make a very exact Pair of Scales the following Directions should be observ'd.

1. The Axis C must be made of good Steel harden'd and well polish'd, but the Edge of it must not be so sharp as to cut.

2. The two Rings on which the Axis is to bear, such as A a B b, must also be of hard Steel and well polish'd, but the lower Part of them should be the narrow End of an Oval, the Hole being of the Figure K. The Planes of these Rings with the Pieces that carry them (here represented by m n as broken, with the Examen o between them also broken) must be exactly parallel to one another, and one and the same Line must be their common Axis.

N. B. If these Rings were made of *Agate* or any harder Stone well polish'd, they would do better than Steel.

3. The Points of Suspension of the Scales, such as S, must be exactly equidistant from C the Middle of the Beam, and the lower Part of the Hole S must be a sharp Edge of hard and polish'd Steel.

4. Each Scale being fitted with its Hook and Strings must be weigh'd singly in another Pair of Scales, observing to weigh each of them in the same Scale against the same Counterpoise in the other Scale, without shaking the Beam to alter the Position of the Axis in the manner above mention'd.

5. When the Beam being suspended, does by its *Examen* (that is, the slender perpendicular Piece over the Axis) appear to have its two *Brachia* exactly in *Æquilibrium*; to try whether the Points of Suspension are exactly equidistant from the Axis at A, hang on the Scales (prepar'd as before directed) and if they are in *Æquilibrium*, and continue so upon changing them for one another, then you may be sure that both Scales and Beam are well adjusted.

6. But if by the preponderating of one Scale, the Beam appears to be unequally divided; then with a Pair of Pliers bend the Part L of the Hook, in order to bring S nearer to, or to remove it farther from the Point A, and if that alters the *Æquilibrium* of the Beam without the Scales, hang a Thread or some small Weight on that End of the Beam which has been made shorter, to restore that *Æquilibrium*; then try with the Scale, and if the *Æquilibrium* continues, file off so much of the heaviest *Brachium*, as the Weight of the Thread amounts to. If the Beam be made

Annotat. made of Steel, after the Hook *LS* has been harden'd, one must with a
 Lect. III. Blow-Pipe bring down the Temper of the Part *L*, that it may be soft
 enough to bend without breaking.

7. The Points of Suspension for the Scales, as *S*, must be in the same horizontal Line as the Axis *A*, the Scales must hang very freely on their Hooks, and the Center of Gravity of the Beam must be a very little under *A*.

8. When the Beam of a Pair of Scales weighs from three or four Ounces to about one or two Pounds, sometimes the Points of Suspension of the Scales are shut up within Boxes at the Ends of the Beam such as *B**, thro' which a square Piece as Steel *C* passes, whose upper Part in the middle has an Edge like the Axis of the Beam, but only with the Edge upwards, to sustain the Eye *ee*, and the Hook of the Scale. *Cc** represents one of those Pieces out of the Box, and its middle Section is mark'd *D*, whose upper Edge supports the *E**. In order to adjust such a Balance nicely (all other Things being fix'd as beforemention'd) there should be a long Hole like *Hb**, for the suspending Pieces as *Cc*, to be mov'd in, so as to be brought nearer to, or farther from the Axis of the Balance, by means of a Screw Pin *P*; that when they come to be exactly equidistant from the said Axis (which can only be known by hanging on Scales or Weights exactly equal, having due Regard to the Effect of Pins lengthened or shortened without the Box at *P* by screwing) the Pieces *C* may be fix'd, the Screw Pins filed off, and the rest of the Hole fill'd up on each side the supporting Piece *C*.

* Plate 13.
Fig. 14.

* Plate 13.
Fig. 15.

* Plate 13.
Fig. 16.

* Plate 13.
Fig. 17.

N. B. It is not to be expected that a large Pair of Scales should be so exact as a small Pair; because the Friction encreases according as it is heavier: So that if a Balance, whose Beam and Scales weigh six Ounces Troy, be turn'd with $\frac{1}{6}$ of a Grain, it may be said to be as nicely adjusted as Brigadier Armstrong's Balance abovemention'd, which weighing 16 times less, turns with $\frac{1}{256}$ of a Grain. So likewise when a Balance turns with a small Part of a Grain, we must not expect it to turn as easily when the Scales are loaded; for they will always become less nice, according as they encrease in Weight.

The Pulley is liable to a great deal of Friction on account of the Stiffness of the Ropes, the Smallness of the Wheels or Sheevers in Proportion to their Center-Pins, and their rubbing against the Sides of the Block or Frame in which they move.

Great Care should be taken that Pulleys, which are to be us'd in Building, or any where at Land, should receive no Wet; for by so doing, the Ropes twist and thicken, so as often to require a great Force to draw them thro' the Blocks in that Condition, even when no Weight is rais'd; but when one cannot avoid wetting them, tarr'd Ropes must be us'd as at Sea.

To prevent the Wheels rubbing too much against the Blocks that contain them, there ought to be thin Collars of Brass or Iron of a Diameter much less than the Wheels on the Pins on each side of the Wheel.

To

To lessen the Friction on account of large Pins when great Weights are to be rais'd, we should considerably increase the Diameter of the Wheels, tho' in some Cases the Machine would be too cumbersome, and therefore not so easy to be managed. But as People do not commonly imagine so much Difference as there is between small and large Pulleys, provided their Number and Combination to be the same; I shall explain that Case in considering the Way of finding what the Quantity of Friction is, and only mention here that according to *Monf. Amonion's Experiments and Calculations**, there is so much Friction in Pulleys on account of the Force requir'd to bend the Ropes and overcome the Friction of the Pins when the Wheels or Sheeves are small, That if over an upper Pulley of 3 Inches Diameter with an Inch Pin, there was a Rope of $\frac{2}{3}$ Diameter having 800 Pound Weight at each End, which Weights must keep each other in *Equilibrio* in this Case (because here the Pulley only does the Office of a Roller) in order to make one of the Weights preponderate and overcome all the Frictions, so as to bring the other Weight over, one must add $436\frac{2}{3}$ to that Weight, call'd the *Power* in such a Case: But if the Pulley had been of 24 Inches Diameter, the Diameters of the Rope and Pin being as in the small one, 45 Pound would have been a sufficient Addition to the Power to enable it to give Motion to the Weight by overcoming the Friction. It is therefore well worth considering the Quantity of Friction in order to direct us in Practice; since in this one Instance by only using a Pulley-Wheel of 24 Inches Diameter instead of one of 3, the Force to overcome the Friction is less by the Quantity of $391\frac{2}{3}$ Pound; or less than $\frac{1}{6}$ of the Power added to it will bring up the Weight, when in the other Case there must be more added to the Power than one half of it.

Annotat.
Lect. III.
* *Memoirs of the Royal Academy for the Year 1699.*

The Axis in *Peritrochio* has but little Friction if the Wheel be large and the Axle small, except what arises from the folding of the Rope round the Axle if a large Rope be made use of to raise a great Weight. But yet we shall shew how to find the Friction of the Axle, whatever it be, after we have taken notice of the Friction of the other Machines in general.

The inclin'd Plane is not liable to much Friction, if the Weight which is roll'd up be spherical or cylindrick; for then the whole Friction arises only from what the Plane wants of perfect Hardness, so as to suffer the Body rolling up to sink a little, which alters the Inclination of the Plane in that Place by making it steeper, and renders the Line of Direction a little inclin'd to the Plane: So likewise, if the Plane being hard, the Body yields and alters its Figure a little; then it must be lifted at every Pull, or go up by jumps. But if the Body to be drawn up be flat, or of any Figure not spherical or cylindrical, as a Piece of Timber, or at least a Sledge loaded, then the inclin'd Plane will have a great deal of Friction; which we shall shew how to estimate.

The Wedge has a great deal of Friction; for besides all this last mention'd Friction of the inclin'd Plane when flat Bodies slide against one another, there must be added the third Part of the Pressure which the same Body gives more to the Wedge than the inclin'd Plane, by reason of the

T

Obliquity

Annotat. Obliquity of the Draught, a Wedge being only an inclin'd Plane with the
 Lect. III. Line of Direction parallel to the Base, instead of being parallel to the Hypotenuse of a Triangle, whose Height is the Thickness of the Wedge.

N.B. *We do not here take notice of the Cohesion of the Wood or Bodies to be cloven, because the Resistance there to be surmounted is to be consider'd as a Weight, and therefore attributed to the Weight and not the Power.*

The Screw has a Friction of the same Nature as that of the Wedge, because it is compounded of a Wedge; but greater, because it touches in all its Parts at once, which a Wedge does not**.

Pl.II.F.II. Screw representing only a single Wedge rising in the Direction HK or
 †Pl.II.F.I4. LM, &c. has less Friction than the sharp-threaded Screw†; because in this last the Surface of the Thread of the Screw is inclin'd to the Base as
 Pl.II.F.I3. well as to the Axis or Arbor of the Screw. But the endless Screw has yet more Friction than the sharp-threaded common Screw, because it is oblig'd to take obliquely the Tooth of the Wheel which it drives.

For this reason in Clock or Watch-work those that would change Wheels and Pinions into endless Screws should be aware of the great Friction that is in them, and not make use of them unless the Nature of the Movement requires it, and there be more gain'd by the Alteration of the Direction of Motion than is lost by Friction.

To settle such a Theory of Friction as may serve to direct our Practice (that we may not only make a just Estimate of its Quantity in every simple Engine; but also find out the Friction of the several Parts which make up a compound Engine, so as to enable us to know what to allow for the Friction of the whole complex Machine) depends upon so many Experiments and Observations, that I rather choose to consider that Subject in a *Lecture* on purpose (*viz.* the 4th *Lecture*) than go on with it here, which would swell these Notes to too great a Bulk; so much remaining to be consider'd in relation to the several Particulars of the 3d *Lecture*, that they will far exceed the Length of the *Lecture*; unless I were to omit some Things very material, which yet are too difficult to be mention'd in the *Lectures* themselves, which I have made so easy as to require only the Reader's Attention, without any previous Knowledge of Mathematics.

4. [20. — *The Beam hanging freely on its Center of Motion, which is plac'd a little above its Center of Gravity.*] As no Position of the Beam of the Balance but the horizontal can make us judge of the Weight of the Bodies to be compar'd by that Instrument, we must be careful that the Center of Motion or Point of Suspension of the Beam be not in the Center of Gravity, because then not only the Beam would remain in any given Position* (as well inclin'd as horizontal) but also continue in that Po-

*L. 2. 26.

***Tho', strictly speaking, the Friction of Bodies of the same Weight does not encrease in Proportion to the Number of Parts that rub; yet it so happens here, because in this Case the rubbing Surfaces apply less exactly to one another than in the Wedge.*

friction

tion when equal Weights are suspended at its Ends. To illustrate this, let us examine the 1st Figure of Plate 14. [†] Annotat.
Lect. III.

$ACbD$ represents the Section of a Beam of some Thickness, which when suspended by its Center of Gravity K , will as well remain in the inclin'd Position $AEBF$ as in the horizontal Position: Now if the heavy Bodies P, W (equal, if the Body be equally divided in its Length by the Point K , or reciprocally proportionable to the *Brachia* if the Beam be unequally divided) be suspended at the Ends A, B , they will hang in *Equilibrio* in any Inclination of the Beam, let the Beam be of any Size whatever. First, let the Beam be suppos'd so slender as to have little or no Weight in comparison of the Bodies, as the Line AB with its Center of Gravity at K ; it is evident that the common Center of Gravity of P and W , which is at G , will not be sensibly remov'd from the Point G by the Addition of the Beam suppos'd of little or no Weight, nor by the Removal of the Bodies to p, w , when the Beam is inclin'd into the Position ab^* . Secondly, if we consider the Beam with all its Weight; when in the horizontal Situation CD , the heavy Part CB pressing upon AB being equal and equally distributed over the Beam, and the lower Part AD equal to CB hanging under in the same manner as CB presses above, the Center of the Gravity of the Beam will not on that account be removed from the Point K ; but the common Center of Gravity of the Bodies P, W , and the Beam will be remov'd from G to g under the Center of Motion K ; therefore the Balance and Weights will remain in that Position, because the Point K of the Line of Direction is supported \parallel . Neither will the Weights (which in a Balance always hang freely) be able to alter this horizontal Position by their Suspension, because their Distances AK, BK from their Lines of Direction to the Center of Motion K (upon which their Velocities depend) are equal or reciprocally proportionable to their Masses $*$. Now if the Beam be inclin'd in the Position $EAFB$, we may still consider in it the slender Beam AB loaded above and below with the two equal Prismatical Wedges AFB and AED , whose Centers of Gravity being at m and n , their Lines of Direction will go through the Points r and s equally distant from the Center of Motion K , therefore they will balance each other \dagger , and consequently not alter the inclin'd Position of the Beam. [†] Then if we consider the Weights p and w suspended from the Points a and b , the common Center of Gravity of the Beam and Weights will still be at g , and the Distances of the Lines of Direction of the Weights now becoming tK and uK decrease exactly in a reciprocal Proportion to the Weights; so that there can be no Motion occasion'd by this Position of the Beam and Weights, because there is no Alteration of Place in the common Center of Gravity of the whole loaded Balance, or the respective Velocity of the Weights. * L. II. 31;
32, 33.
L. II. 45.
* L. II. 137
and L. III.
18 and 22.
† 18.

But if we remove the Center of Motion or Point of Suspension of the Balance to k a little above the Center of Gravity of the Beam K , the Line of Direction, which in the horizontal Position of the Beam is kg , will in the inclin'd Position of the Beam be out of the Point of Suspension,

Annotat. fion, which will then be remov'd to c , and the Center of Gravity must move
 Leſt. III. from g to q , which it can do deſcribing a ſmall Arc round the Point c ,
 which will reduce the Beam to an horizontal Poſition, in which the Line
 of Direction will be cq going through the Point of Suſpenſion c , and con-
 ſequently *Weights that are in Æquilibrio upon a Balance whoſe Center of*
Motion is above the Center of Gravity of the Beam will reduce the ſaid Ba-
lance from an inclined to an horizontal Poſition.

Now in fixing the Center of Motion of the Beam above the Center of Gravity, Care muſt be taken not to fix it above the Points of Suſpenſion, as ſome Authors have taught, and as the Practice is in making the common Scale-Beams, which may be conſider'd as made up of two Leavers making an obtuſe Angle at k the Center of Motion, whiſt the Points of Suſpenſion A, B , are under its Level in the Line AB . Such a Balance is uſ'd for common Purpoſes, becauſe it comes ſooner to an Æquilibrio, than if AkB was one Line; but it is a *false Balance*, and a *ſkilful Perſon may cheat with it in proportion as the Angle AkB is more acute*,
 * Pl. 14. F. 2. *eſpecially when there is no Perpendicular Piece or Examen as Cr^* , to ſhew when the Balance is truly horizontal; for unequal Weights may make an Æquilibrio on ſuch a Balance, and not be diſcover'd by changing the Scales,* which preſently diſcovers the Cheat in a Balance whoſe Beam has its *Brachia* unequal †. For example, let the Balance $ACBc$ (whoſe Center of Gravity is at c and Center of Motion at C with the equal Weight P, P , hanging at its Extremities A, B) be plac'd in the inclin'd Poſition ab , I ſay that as the Line of Direction Dd of the Weight P is brought nearer to the Center of Motion (*viz.* to $q\pi$) the ſaid Weight may be increas'd in proportion as its Diſtance Cu is diminifh'd by being reduced to Cq , and act with the very ſame Force on its Point of Suſpenſion: Whiſt the Line of Direction of the oppoſite Weight P being remov'd from Dd to $b\omega$, its Diſtance Co becomes Cb , and therefore P may be diminifh'd in proportion as its Diſtance is increas'd; therefore in that Situation of the Balance the Weights P and P will keep each other in Æquilibrio, when they differ in the reciprocal Proportion of their Diſtances qC and bC , or bC and Ca if the Balance be inclin'd the other way in the Line $\alpha\beta$; the Diſcovery being only made by the viſible Inclination of the *Examen* towards s or t . Q. E. D.

The nearer the Center of Gravity of the Beam is to the Center of Motion, the nicer will be the Balance, becauſe the Beam will be the more
 * Pl. 14. F. 2. apt to vibrate quick from Side to Side. As for example, if $acbC^*$ be the Beam and C the Center or Axis of Motion, the Difference between the Effect of having the Center of Gravity at K , or c , will be the ſame as if we compar'd the Vibrations of two Pendulums of the Lengths CK and Cc , whoſe Velocities in their Vibrations reciprocally are in a ſubduplicate Ratio of their Lengths (as I ſhall further ſhew when I come to treat of Pendulums) for the Beam is really a Pendulum.

E X P E R I M E N T. Plate 14. Fig. 3.

To the Beam AB, whose Axis of Motion is C, fix a screw'd Wire K Pl. 14. F. 3. and a Ball W so contriv'd that it may be screw'd on towards, or screw'd fromwards the Axis C. When by bringing the Ball to W the Center of Gravity is at K, the Vibrations of the Beam will be quicker than when the Center of Gravity is brought to *k* by lowering the Ball to *w*. *This may be used in Practice for some nice Experiments, because by such a Contrivance the Center of Gravity may be brought as near as you please to the Center of Motion.*

N. B. That the Center of Gravity ought never to be above the Center L. 2. 26. of Motion appears from what has been already said in the second *Lecture*; No. 26.

5. [27. *In all these Cases we suppose the Weights to hang freely from the Ends of the Balance to which they are fasten'd.*] Though in the common Use of the Balance the counterpoising Weights, or the Scales, generally hang freely; yet there are some Cases where they do not; and in compound Engines where Balances are often a Part of a complex Machine; instead of Weights, Powers are applied to their Ends in all manner of Directions, and then they become Leavers of the first kind, such as the Regulators in several Water Engines, Beams to blow Bellows, &c. Therefore I shall consider the Effects of Powers applied obliquely to a *Mathematical Balance* or straight inflexible Line, which will also solve all Cases of the Leaver, and (with proper Allowances) may be applied to all the Mechanical Organs.

Let the Balance AB*, 12 Inches long and equally divided by its Center of Motion C, sustain at its Ends the two equal Weights W, P, which last we shall consider as the Power. Whilst the Power draws in the Line BP, it acts according to its whole Intensity†, its Distance being then $CB = CA$ the Distance of the Weight, both Distances being measured upon the Beam: But if the Power be removed to *P*, and (its String B*x*P running over the Pulley *x*) it draws obliquely in the Line B*x*, which makes with the Beam the acute Angle CBF, or (what comes to the same) the obtuse Angle CBE (because here we suppose it as much greater than a right Angle as CBF is less, CBF being = CB*φ*) the Force of the Power will be diminish'd in proportion as CF or C*φ* the Distance from the Line of Direction of the Power acting obliquely is less than CB the Distance of the Line of Direction of the Power acting directly or at right Angles to the *Brachium* of the Balance CB*. Knowing therefore the Intensity of the Power, which acting at right Angles at B (or which hanging freely from B, if an heavy Body, as represented here, be the Power) keeps the opposite Weight W in *Æquilibrio*, one may easily find how much the Power must be encreas'd to keep the said Weight in *Æquilibrio* when it draws obliquely.

Annotat. obliquely in any known Direction, as for example in the Direction represented in the Figure; or (which is all one) how much the Weight P , whose String fasten'd to B runs over the Pulley α , must be greater than the Weight P which hangs freely to have the same Effect. With the Distance CF or $C\phi$ draw the Arc $Ff\phi$ which cuts CB at f ; then you will have the Quantity of P by this Analogy.

*As the Length Cf upon the Beam, whose Length is 4, 8 Inches :
Is to CB the whole Brachium of the Beam, here 6 Inches long ::
So is the Intensity of the Power, or the Weight P here suppos'd 40 Pounds :
To the new Power, or Weight P , 50 Pounds.*

Hence it appears that the Weight P thus found, would keep the counterpoising Weight W in *Æquilibrio* by hanging freely at the Point f , as well as it does it by drawing obliquely over the Pulley α ; because the Momentum of W (or $W \times AC$) being divided by Cf , will give the Quantity of the Weight P ; or, in other Words, there will be a reciprocal Proportion between the Weights W and P , and their Distances CA and Cf *. Therefore $Cf = C\phi = CF$ the Distance of the Line of Direction from the Center of Motion (always found by the Length of a Perpendicular from that Center to the said Line of Direction) may in all such Cases be properly call'd the acting Distance of the Power. See No. 20.

* 13.
and L. II. 9,
12, 13, and
16.

Another Way.

Through the Point α taken in the Circumference of the Pulley, over which the String $B\alpha P$ runs, draw αE parallel to the Balance AB , which will cut the perpendicular Line of Direction of the Power (or freely hanging Weight) P , at right Angles at D , $D\alpha$ being $= DE$, and the Angle $\alpha BD = DBE$ by Supposition. In the rectangular Triangle $BD\alpha$, as much as the Hypotenuse $B\alpha$ is longer than the Perpendicular BD , so much must the Quantity or Intensity of the Weight P (which hanging freely keeps W in *Æquilibrio*) be increas'd when it draws obliquely in the Line $B\alpha$; that is, so much must P be greater than P to keep W in *Æquilibrio* drawing obliquely. The same would be true if α was remov'd to E ; and therefore this will always be the Rule to estimate the Force of Powers drawing obliquely.

As the Sine of the Angle of Traction, viz. the Angle which the Line of Direction of the Power makes with the Beam :

Is to the Radius ::

So is the Intensity of the Power drawing at right Angles with the Beam :

To the Intensity of the Power drawing obliquely.

N. B. The Angle of Traction here is $CB\alpha$ or CBE , whose common Sine is BD ; then in this Case $BD : B\alpha :: P : P$.

That

That this Method will always have the same Consequence as the former, Annotat. may be seen by comparing together the two Triangles $B\alpha D$ and CBF ; *Lect. III.* for since CFB is a right Angle by Supposition, and $CB\alpha = B\alpha D$ because of the Parallels CB and αD , BCF must be equal to αBD and consequently the Triangles be similar, which will shew that CF (48) : CB (60) :: BD (40) : $B\alpha$ (50) : P : P . *Q. E. D.*

This might also be explain'd a third Way, by resolving the Force which draws obliquely in the Line $D\alpha$, into the Forces, the one drawing along the Leaver in the Line BC , and the other in the Line BD , after Sir *Isaac Newton's* Manner*; but we shall consider it in the Case of Trusion or* 85, 86. pushing obliquely against any Point of the Beam, which is the same as drawing the contrary Way.

All these Methods will appear to agree in the following

EXPERIMENT. *Pl. 14. Fig. 5.*

The Balance or Leaver AB , 12 Inches long, is moveable upon the Center C of the Stand S , which has a long Piece so propp'd at S as to remain in an horizontal Position, so as to carry the Pulleys α and E , each 3 Inches distant from the Point D which is placed perpendicularly under B . When the Power P of four Ounces equal to W hangs freely, it keeps W in *Æquilibrio*; but if the String PXB be thrown over the Pulley α or E , then will the said Power P be overpois'd by W till it be changed for P or p a Five-ounce Weight, which drawing obliquely over α or E will keep W in *Æquilibrio*. Now one may observe, that when P descends one Inch, or from P to r , it brings down the End of the Beam B only to the Point b in the horizontal Line fe , which raises the opposite End A to a just as high above the Line AB ; but when the Power at P descends one Inch, namely to q , it brings down the End B to β in the horizontal Line bg , and consequently raises the opposite End A so much higher, so as to give more Velocity to the Weight W bringing it to w instead of w . Now since the Powers P and P with the same Velocity (or descending equally) give W different Degrees of Velocity, their Intensities must be different in that Proportion, because *Causes are always proportionable to their Effects*; therefore P must be greater than P as much as the Arcs Aa and $B\beta$ are greater than Aa and Bb , or rather as much as the Sine αo is greater than αn . This also appears by observing (since $BX = \beta x (= yx)$ and By) that the Strings BP , $B\alpha P$, and $\beta \alpha Pq$ are all equal.

N.B. This is strictly true only in the Beginning of the Motion of the Beam; but that is sufficient for our Purpose.

The very Sight of the Machine makes it plain that a Power acting at right Angles is the most effectual. For as in removing the Power from the Perpendicular it draws more weakly at E , and at m it only pulls the Center C in the Direction Bm so as to do nothing towards raising A the opposite

Annotat. opposite End of the Beam; and going the contrary Way, the Power weakn'd at x , becomes wholly ineffectual when brought to the Beam to draw in the Direction BC, because then it only acts against the Center C no Way moving the End A: Therefore a mean Situation of the Line of Direction between those two ineffectual Extremes, must be the most effectual; and that is in the Line BP perpendicular to the Beam.


It is upon this Principle that most of the Feats of the pretended strong Men, or modern Samplons are perform'd; the Machines on which they sit or stand being so contriv'd that the drawing Horses or hanging Weights pull those Limbs of the Man resisting (which perform the Office of a Leaver or Balance) in such a manner that they are drawn directly against the Center of Motion. But this I shall explain more particularly when I come to speak of those Feats of Strength.

If at one of the Ends of a Leaver or Balance be fix'd a Weight, which (moving with the said End of the Leaver) does not hang freely, whilst the Power acting at the other End is either a heavy Body hanging freely, or an animate Power pressing perpendicularly towards the Earth; I say, that such a fix'd Weight will vary in Force according to the Position of the Beam, and that Force will vary in a contrary manner according as the Center of Gravity of the Weight is above or below the Beam; namely, when the Center of

Pl. 14. F. 6. *Gravity of the fix'd Weight is below the Beam (as in Fig. 6.) the Weight will become heaviest (or act most strongly) when rais'd above the horizontal Line (as at G) in the inclin'd Situation of the Leaver DCG; and it will become lightest when depress'd below the horizontal Line (as at E) in the inclin'd Situation of the Leaver ACE; on the contrary, if the Center of*
 Pl. 4. F. 7. *Gravity of the fix'd Weight be above the Leaver (as in Fig. 7.) it will become heaviest when depress'd below the horizontal Line (as at K) in the inclin'd Situation of the Leaver ACK; and become lightest when rais'd above the horizontal Line (as at E) in the Situation of the Leaver DCE. But the said fix'd Weight will act in the same manner as if it hung freely when the Leaver is in the horizontal Situation as at BF (Fig. 6.) and BI (Fig. 7.)*

Pl. 14. F. 6. The first Case (Fig. 6.) may be thus explain'd. In the Position of the Leaver BF, the Line of Direction qO going thro' the Point of Suspension q , and being at right Angles with the Leaver, Cq is the acting Distance*, of the Weight as well as the Distance of the Point of Suspension; therefore as $BC : to Cq ::$ so is the Weight F : to the Power R ; in the same manner as if the Weight F hung freely from q . But when the Weight is rais'd up to G , as the Center of Gravity O cannot get under the Point K (the same as q the Point of Suspension in the horizontal Leaver) Or becomes the Line of Direction instead of KM which would have been the Line of Direction if the Body had hung freely from K ; therefore Cr is the acting Distance of the Weight instead of CM , when the acting Distance of the Power is become LC ; and consequently the Weight has more Force, and can only be balanced by a greater Power as P . For whereas $CL : CM :: F$ (or G) : R , now $CL : Cr :: F$ (or G) : P ,
 From

* See Page
142.

a stronger Power or heavier Counterpoise. But if the Weight be brought to E, its Line of Direction instead of nH becomes MO , and consequently its acting Distance is less than it ought to be in Proportion to CQ the acting Distance of the Power, which therefore may be diminish'd in Intensity and become S instead of R . 

From what has been said, and a View of *Fig. 7.* * one may see that the contrary must happen when the fix'd Weight has its Center of Gravity above the Leaver or Beam. For in the Situation of the Leaver ED , the Line of Direction of the Weight coming forward too fast (becoming OL instead of Eb) CL the acting Distance of the Weight at E bears a less Proportion to CN the acting Distance of the Power at S , than Cg to CB , the Proportion in the horizontal Situation; and consequently the Power S of less Intensity serves as a Counterpoise instead of R . But when the Weight is depress'd to K , we must make use of P a greater Counterpoise; because Cg is become the acting Distance instead of Ct . And in the horizontal Situation of the Beam the Body O weighs the same as if it hung freely, because Og is its Line of Direction, as it would be if the said Body hung down from g . This may be observ'd in Practice. Suppose the Man M (*Fig. 8.*) is lifting Hay, Sheaves of Corn, or a large Faggot A , by means of the Prong or Fork AB resting upon his Knee c as a *Fulcrum* and pressing down the End B of the Prong or Leaver; if the Fork A goes under the Faggot, it will be the Case of the inclin'd Leaver AK (*Fig. 7.*) and the Burthen will grow lighter as it rises. But if the Fork had been thrust through the Binding above the Faggot, it would be the Case of the inclin'd Leaver AE (*Fig. 6.*) at whose End the Burthen becomes heavier as it is rais'd up; and then if the Man was just able to begin to lift it, he must let it go back again, or find a new *Fulcrum* as C in order to raise that Weight. Suppose again, that the Beam of a large Pair of Scales, was inclin'd in the Position ACE ; a Man in the Scale suspended at H may, by thrusting up his Hand hard against the Beam, put himself in the Condition of the heavy Body HE , and consequently appear to weigh less than his true Weight, being counterpois'd by such a Weight as S . But if the Scale which he gets into be rais'd up, its Point of Suspension being at K in the Position of the Leaver DC ; the Man by thrusting hard against the Beam above his Head, may throw his Body into a Position perpendicular to the Leaver, and together with the Scale be in the Condition of the Body KG ; so that if there is more Weight in the opposite Scale, the Weight need not be lessen'd till its Intensity be equal to that of the Man's Weight, but it will begin to be lifted up whilst it is still greater; so that in such a Case, a Man's Weight will appear greater than it is, by as much as the Counterpoise P is greater than R . Just the Reverse would happen, if a Man was to sit upon the Beam; for then he would weigh least, when rais'd above the horizontal Line going through the Center, as at EM (*Fig. 7.*) and most, when below the said Line, as at FK .

Annotat. Now tho' the Action of heavy Bodies on one another may thus in the
 Lect. III. Balance and Leaver and some other of the Mechanical Organs be estimat-
 ed by the Distance of their Line of Direction from the Center of Motion; yet this is only so far true, as the perpendicular Ascent and Descent is agreeable to that Distance; for there are Cases especially in the working of compound Engines, where the Distance of the Line of Direction of a rising or falling Body from the Center of Motion is not proportionable to the perpendicular Ascent or Descent of the said Body. Therefore the Velocity of a Power, when it is an heavy Body, must be consider'd in the same manner as that of a Weight; as has been fully explain'd in the 8th
 * L. 2, Ann. 8 Note of the second Lecture*; and may be further prov'd by the following

E X P E R I M E N T.

* Pl. 14. F. 9. ACBEKD* is a Balance in the Form of a Parallelogram passing through a Slit in the upright Piece NO standing on the Pedestal M, so as to be moveable upon the Center-Pins C and K. To the upright Pieces AD and BE of this Balance are fix'd at right Angles the horizontal Pieces FG and HI. That the equal Weights P, W, must keep each other in *Equilibrio* is evident; but it does not at first appear so plainly, that if W be removed to V, being suspended at 6, yet it shall keep P in *Equilibrio*; tho' the Experiment shews it. Nay, if W be successively moved to any of the Points 1, 2, 3, E, 4, 5, or 6, the *Equilibrium* will be continu'd; or if, W hanging at any of those Points, P be successively mov'd to D or any of the Points of Suspension on the Cross-Piece FG, P will at any of those Places make an *Equilibrium* with W. Now, when the Weights are at P and V, if the least Weight that is capable to overcome the Friction at the Points of Suspension C and K be added to V as *u*, the Weight V will overpower, and that as much at V as if it was at W.

From what we have said above, the Reason of this Experiment will be very plain. As the Lines AC and KD, CB and KE always continue of the same Length in any Position of the Machine, the Pieces AD and BE will always continue parallel to one another and perpendicular to the Horizon, however the whole Machine turns upon the Points C and K; as appears by bringing the Balance to any other Position as *abed*: and therefore as the Weights applied to any Part of the Pieces FG and HI can only bring down the Pieces AD and BE perpendicularly in the same manner as if they were applied to the Hooks D and E or to X and Y the Centers of Gravity of AD and BE, the Force of the Weights (if their Quantity of Matter is equal) will be equal; because their Velocities will be their perpendicular Ascent or Descent, which will always be as the equal Lines $\propto x$ and $\propto y$, whatever Part of the Pieces FG and HI the Weights are applied to. But if to the Weight at V be added the
 little

little Weight u , those two Weights will overpower, because in this Case Annotat. the *Momentum* is made up of the Sum of V and u multiplied by the *Lect. III.* common Velocity y .

Hence follows, that it is not the Distance c multiplied into the Weight V , which makes its *Momentum*; but its perpendicular Velocity y multiplied into its Mass*. *Q. E. D.*

* 43.

This is still further evident by taking out the Pin at K ; for then the Weight P will overbalance the other Weight at V , because then their perpendicular Ascent or Descent will not be equal.

To conclude all that relates to Forces applied in different Directions to Leavers and Balances, I shall explain the Action of oblique Forces by the *Composition* and *Resolution* of Motion after Sir *Isaac Newton's* Manner; applying the Solution to a Proposition on the Balance, which has not been taken notice of by mechanical Writers tho' often talk'd of by handicraft Workmen.

THEOREM. *Fig. 10.*

Pl. 14. F. 10.

AB is a Balance, on which is suppos'd to hang at one End B the Scale E with a Man in it, who is counterpois'd by the Weight W hanging at A the other End of the Balance. I say, that if such a Man, with a Cane or any rigid freight Body, pushes upwards against the Beam any where between the Points C and B (provided he does not push directly against B) he will thereby make himself heavier or overpoise the Weight W , tho' the Stop GG hinders the Scale E from being thrust out fromwards C towards GG . I say likewise, that if the Scale and Man should hang from D , the Man by pushing upwards against B or any where between B and D (provided he does not push directly against D) will make himself lighter or be overpois'd by the Weight W , which did before only counterpoise the Weight of his Body and the Scale.

If the common Center of Gravity of the Scale E and the Man suppos'd to stand in it be at k , and the Man by thrusting against any part of the Beam cause the Scale to move outwards so as to carry the said common Center of Gravity to x ; then instead of BE , Ll will become the Line of Direction of the compound Weight, whose Action will be increas'd in the Ratio of LC to BC . This is what has been explain'd by several Writers of Mechanicks; but no one, that I know of, has consider'd the Case when the Scale is kept from flying out, as here by the Post GG , which keeps it in its Place, as if the Strings of the Scale were become inflexible. Now, to explain this Case, let us suppose the Length BD of half of the *Brachium* BC to be equal to three Feet, the Line BE to four Feet, and the Line ED (of five Feet) to be the Direction in which the Man pushes, DF and FE to be respectively equal and parallel to BE and BD , and the whole or absolute Force with which the Man pushes, equal to (or able to raise) 10 Stone. Let the oblique Force ED (= 10 Stone)

U 2

be

Annotat be resolv'd into the two EF and EB (or its equal FD) whose Directions
 Lect. III. are at right Angles to each other, and whose respective Quantities (or Inten-
 sities) are as 6 and 8, because EF and BE are in that Proportion to
 each other and to ED. Now since EF is parallel to BDCA the Beam,
 that Force does no way affect the Beam to move it upwards; and there-
 fore there is only the Force represented by FD, or 8 Stone to push the
 Beam upwards at D. For the same reason, and because Action and Re-
 action are equal, the Scale will be push'd down at E with the Force of
 8 Stone also. Now, since the Force at E pulls the Beam perpendicularly
 downwards from the Point B distant from C the whole Length of the Bra-
 chium BC, its Action downwards will not be diminish'd, but may be ex-
 press'd by $8 \times BC$: Whereas the Action upwards against D will be half
 lost, by reason of the diminish'd Distance from the Center, and is only to

be express'd by $8 \times \frac{BC}{2}$; and when the Action upwards to raise the Beam
 is subtracted from the Action downwards to depress it, there will still re-
 main 4 Stone to push down the Scale; because $8 \times BC - 8 \times \frac{BC}{2} =$
 $4 BC$. Consequently a Weight of 4 Stone must be added at the End A
 to restore the *Æquilibrium*. Therefore a Man, &c. pushing upwards under
 the Beam between B and D, becomes heavier. Q. E. D.

On the contrary, if the Scale should hang at F from the Point D only
 three Feet from the Center of Motion C, and a Post gg hinders the Scale
 from being push'd inwards towards C; then if a Man in this Scale F
 pushes obliquely against B with the absolute Force abovemention'd; the
 whole Force, for the Reasons before given (in resolving the oblique Force
 into two others acting in Lines perpendicular to each other) will be redu-
 ced to 8 Stone, which pushes the Beam directly upwards at B, while the
 same Force of 8 Stone draws it directly down at D towards F. But as
 CD is only equal to half of CB, the Force at D compar'd with that at
 B loses half its Action, and therefore can only take off the Force of 4
 Stone from the Push upwards at B; and consequently the Weight W at A
 will preponderate, unless an additional Weight of 4 Stone be hang'd at B.
 Therefore a Man, &c. pushing upwards under the Beam between B and D
 becomes lighter. Which was also to be demonstrated.

Scholium I.

Hence, knowing the absolute Force of the Man that pushes upwards
 (that is, the whole oblique Force) the Place of the Point of Trusion D,
 and the Angle made by the Direction of the Force with the Beam at the
 said Point, we may have a general Rule to know what Force is added to
 the End of the Beam B in any Inclination of the Direction of the Force
 or Place of the Point D.

Rule

Rule for the first Case.

First find the perpendicular Force by the following Analogy, whose Demonstration is known to all that understand the Application of oblique Forces.

As the Radius :

To the right Sine of the Angle of Inclination of the Force to the Beam ::

So is the oblique Force :

To the perpendicular Force.

Then the Perpendicular Force multiplied into the Length of the *Brachium* BC, minus the said Force multiplied into the Distance DC, will give the Value of the additional Force at B, or of the Weight requir'd to restore the *Æquilibrium* at A.

Or to express it in the Algebraical Way. Let *of* express the oblique Force, *pf* the Perpendicular Force, and *x* the Force requir'd, or Value of the additional Weight at A to restore the *Æquilibrium*.

$$\frac{DE : DF (= BE) :: of : pf.}{pf \times BC - pf \times DC = x.}$$

The same Rule will serve for the second Case, if the Quantity found be made negative, and the additional Weight suspended at B. Or having found the Value of the Perpendicular Force, the Equation will stand thus : $- pf \times BC + pf \times DC = -x$; and consequently the additional Weight must be hang'd at B; because $-x$ at A is the same as $+x$ at B.

Scholium 2.

Hence it follows also, that if, in the first Case, the Point of Trusion be taken at C, the Force at B (or Force whose Value is requir'd) will be the whole Perpendicular Force; because CD is equal to nothing: And if the Point D be taken beyond C towards A; the Perpendicular Force pushing upwards at that Point, multiplied into DC must be added to the same Force multiplied into BC, that is $pf \times BC + pf \times DC = x$.

The Machine I made use of to prove this experimentally, was as follows. (Fig 11.) The Brass Balance AB is 12 Inches long, moveable upon the Center C, with a Perpendicular Piece Bb hanging at the End B and moveable about a Pin at B, and stopp'd at its lower End b (by the upright Plate GG) from being thrust out of the Perpendicular by the pushing Pipe FE, whose lower Point being put into a little Hole at H, the upper Wire or Point (when put into another little Hole under the Beam at D) is

Annotat. D) is by means of the Worm-Spring EF pressing against the Plug E to drive forward the said Wire *bD*, made to push the Beam upwards with the Force of the Spring EF. TSS is a Stand to which is fix'd the Pillar TC which sustains the Balance; and it has also a Slit SS to receive a Shank of the moveable Plate GG, to be fix'd in any Part of the Slit by a Screw underneath.

EXPERIMENT. Pl. 14. Fig. 11.

Hang on *Bb* as in the Figure. Then let EF be so applied to the Hole H, that its upper Wire *bDk* may go through a little Loop at D, so as not to thrust the Beam upwards, but be in the same Position as if it did, that by hanging on the Weight W the *Brachium* BC with *Bb* and FE may be counterpois'd, that the Action against D and H may be estimated without the Weight of the pushing Pipe.

Then drawing down the End of the Wire *k* thrust it into the little Hole under D, and B will be so pull'd downwards as to require the additional Weight P of 4 Ounces to be hung on at A, to restore the *Equilibrium*: When BH is 4 Inches; BD 3 Inches, and the whole Force of the Spring equal to 10 Ounces.

I need not here say, that for explaining the second Case, *Bb* is to be suspended at D, with the Plate GG fix'd to stop it at the Place M to keep it from being push'd towards T, and that the upper End of GFED*k* must push into an Hole made under B, in which Case the Weight P must be hang'd at B to restore the *Equilibrium*.

N. B. To shew experimentally that the Force which the Spring exerts in this oblique Trusion is equal to 10 Ounces: Take the Beam AB, which weighs 4 Ounces, from its Pedestal CT, and having suspended at each End A and B 3 Ounces, support it under its Center of Gravity by the pushing Pipe EF set upright under it, and you will find that the Beam with the two Weights will thrust in the Wire K*k* as far as *b*, the Place which the oblique Trusion drives it to.

6. [29. — *In all these Cases the Leaver is still said to be of the first Kind.*] There is another way of distinguishing the Leavers according to Aristotle and the Mechanical Writers among the Ancients; and that is as the Weight does, or does not, rise in the same Direction as the Power. As for example, in the Leaver of the first Kind; as it has the *Fulcrum* (Center of Motion, or *Hypomochlion*) between the Power and the Weight, the Power must move downwards whilst the Weight moves upwards; and that Leaver is by those Authors call'd the *Heterodromous* Leaver; that is, *working or moving different Ways*. But the Leaver of the second, and likewise the Leaver of the third Kind, are both call'd *Homodromous* Leavers, because the Power and Weight being on the same side of the *Hypomochlion*

lion or propping Point, they both go the same Way, tho' in the one the Power always gains, and in the other always loses. Annotat. Lect. III.

If we examine such Instruments as we have in common Use, we may plainly see that they are Leavers of one of the three Kinds. As for example, a Pair of Pincers * is made up of two Leavers of the first Kind, * Pl. 14. F. 12. whose common Center of Motion is at the Rivet C, the Power being applied at the Handles B b to press them together, and thereby pinch the Body D as a Weight at the opposite Ends A a; in which Case the Power acts six times more strongly than if applied directly to the Body D at A, a; since in the two Leavers AB and a b the Distance of the Power BC and b C is triple the Distance of the Weight CA and C a. So is a Pair of Scissors † made up of two such Leavers, whose common Center of † Pl. 14. F. 13. Motion is C, the Power being applied at B b, and the Body to be cut as a Weight at D; where it is evident that the nearer D is to the Points A a, the greater will be the Difficulty in cutting it; and the less, the nearer it is brought to C. The Force of a Leaver in this Way is remarkable in the Brasier's or Tin-man's Sheers *, whereby one Man pressing up * Pl. 14. F. 14. on the Handle B raising the lower Sheer AC, moveable about the Center C, is able to cut a Plate of Brass or Copper D a quarter of an Inch thick, the other Sheer a CE being riveted to a couple of strong Standards fix'd into the Block F. Nippers, Forceps, Snuffers and other such Instruments may in the same manner be found to be Leavers of the first Kind. The little Cart or Carr BCA * very useful in Building, is also * Pl. 14. F. 15. a Leaver of the first Kind: It is made to lift great Stones and carry them to the Builders; upon the farther End of whose Plank or Bottom A the Stone D being laid, and with a little Pains mov'd towards C by wriggling the Plank; the Force of one Man by taking hold of the long Handle at B, and weighing it down as it turns about the Axle-tree of the Wheels EF as the *Fulcrum*, will be able to raise that Stone; and being rais'd, by Help of the Wheels, be able to convey it to the intended Place.

The cutting Knife * us'd by Druggists and Patten-makers, to cut Drugs, * Pl. 14. F. 16. or the Wood they use, is moveable on the Joynt or Center of Motion C, whereby it is fasten'd to the Plank CE; the Power is applied at the Handle B, and the Weight is D the Wood or Drug to be cut: This shews the Instrument to be a Leaver of the second Kind †. So is a Door, † 30. whose Hinges are the Center or Axis of Motion, the Hand or Power being applied at the Part towards the Lock, whilst the Body of the Door is the Weight. A Pair of Bellows are two Leavers of the second Kind, whose common Center of Motion is at the End of the Boards where the Nose begins, the Power being applied at the Handles, whilst the Air to be press'd out between the Boards, by its Resistance acts against the middle of the Boards as a Weight. Thus one may easily perceive that the Latch of a Door, when drawn up by a String, is a Leaver of the second Kind; as are also jointed-Nut-crackers, Horse-barnacles, &c. The Oars of a Boat or Galley and the Rudder are also Leavers of the second Kind, tho' *Aristotle* suppos'd them Leavers of the first; but the Error lies in this,

Annotat. this, that he consider'd the Water as the Weight to be mov'd; whereas
 Lect. III. the Boat or Galley is the Weight to be mov'd. For the Water makes
 Resistance at C* as a *Fulcrum*, the Man or Power acting at B, whilst the
 * Pl. 14. F. Vessel FD is mov'd by that Part of the Leaver which presses on it at
 17 and 18. the Point D, as the Oar in Fig. 17. and the Rudder in Fig. 18. But *Aristotle*, in his *Mechanics*, rightly reckons the Masts of Ships among Leavers of the second Kind, assigning the *Fulcrum* or *Hypomochlion* to be in the
 † Pl. 15. F. 1. bottom of the Ship B† considering the Ship it self as the Weight or Burthen, resting with its upper Deck C upon the Mast, as upon a Leaver, and so to be mov'd forwards. Lastly he affirms, the moving Force to be the Wind gather'd in the Sail, which, by the Help of the Sail-yard, is applied to A; and then he gives a Reason from the Principles hitherto deliver'd, why, the higher the Yard DAE is, so much the swifter the Ship will be carried with the same Wind and Sail; viz. because the farther the *Fulcrum* is from the moving Force (all other Things being at the same time consider'd) so much the easier will the same Power or Force move the Weight.

The same Instrument, according to its different Application may be a Leaver of the first or second Kind: As for example, when a Man who has
 * Pl. 14 F. 15. carried a Stone upon the Carr* goes to deliver it at the Place intended so as to lay it upon its Side E, he lets the End of the Plank rest upon the Ground at E, and lifting up the End B turns the Stone over; which Operation changes the Leaver from one of the first Kind into one of the second, where E is the *Fulcrum*, B the Power, and the Weight is at A.

† Pl. 15. F. 2. The Sheep-sheers† are two Leavers of the third Kind; the common Center of Motion being at the springing Bow at C, whilst the Power or Hand is applied at Pp, and the Wool to be cut is the Weight at W. Thus are the two Legs of a Pair of Tongs visibly Leavers of the third Kind. A Ladder or a Pole to be rear'd against a Wall are also Leavers of the third Kind. But the Use of Leavers of the third Kind is most beautifully shewn in the animal Body, where the All-wise Creator has supplied Animals with a Means to move the Limbs with great Velocity by applying the Power of the Muscles very near the Center of Motion; but at the same time giving the Muscles such very great Force as to perform their Office very suddenly, raising the Limbs even when great Weights hang at their Extremities; as for example, when we lift Weights with our Hands or Feet, or when we hold or break hard Bodies with our Teeth. There is scarce a Bone in the Animal Body but what is a Leaver of the third Kind. It is a delightful and curious Contemplation to consider what Proportion is observ'd in the Animal Body as an Engine, from which Art only copies faintly: There we may see various Applications of Powers, and how they help each other in moving the Limbs, sometimes acting jointly at the same time, sometimes succeeding one another to change Directions, and sometimes acting against each other to stop and check Motion; at other times drawing over Pulleys to alter the Angle of Traction as need requires. But this Subject is so copious that we can only give
 one

one Instance or two here ; referring the curious to consult *Alphonsus Bo-* Annotat.
relli who has written a whole Book upon that Subject (*viz. de Motu Ani-* Lect. III.
malium) and from whom we shall take only a few Examples. Here fol-
lows the 21st Proposition of his first Book.

“ P R O P O S I T I O N X X I .

“ *The absolute Power of every animal Muscle, must necessarily be greater*
“ *than the Weight of the Limb suspended by it, but never less.* Plate 15. Pl. 15. F. 3,
“ Fig 3, and 4. and 4.

“ All-wise Nature has so contriv'd the Shape of Animals made up of
“ several Organs join'd together, as to enable them to move from one
“ Place to another, and perform the several Operations requir'd for the
“ Preservation of Life. But this could not be done by giving an Animal
“ an orbicular Shape like a Ball, but it was proper that he should be
“ made up of several Articulations, such as Hands and Feet, to walk
“ about and handle Objects. But these Limbs could not move about
“ Joints, unless they were drawn by muscular Cords, and those Cords were
“ contracted by a moving Force. We are going to shew, that that moving
“ Force or Power must not be less, but necessarily greater than the Weight
“ and Resistance of the Limbs suspended. Let us consider any Limb,
“ for example the whole Arm ; it is plain that it was necessary for it to
“ move round every way about the Joynt of the Shoulder, that it might
“ be able to draw, suspend, and impell the Resistances as well of the
“ Weight of the Arm as of external Bodies to be handled : Such Opera-
“ tions require a proper Figure, Forces and apt Instruments, and all fit-
“ ted for that Purpose. The Shape without doubt must be long like a
“ Leaver moveable about a Center, or the fix'd Point or firm *Fulcrum* of
“ the Shoulder. Then in the Leaver must be consider'd the Positions in
“ which the moving Power and the Resistance are applied. The moving
“ Power acts by contracting the muscular Cords, which can only be fa-
“ sten'd near the Center of Motion of the Leaver, as has been said be-
“ fore*, whilst the Resistance is applied at its utmost Length or farther
“ End,

* Our Author in his 20th Proposition has in-
geniously shewn that the Tendons to move the
Bones could not conveniently be fasten'd near
the End of the Bone (at a great Distance from
the Center of Motion as we endeavour to do in
mechanical Engines) but near the Articulation
or Joynt. These are his Words.

*Pl. 15. F. 3. ‘ Let there be two Bones * AB
‘ and GF join'd and articula-
‘ ted at AF, namely in such manner that AB
‘ may be drawn round C the Center of the Ar-

‘ tication, as the two Bones of the Arm ; and
‘ let the Muscle DE be fasten'd to G the fix'd
‘ End of the Shoulder at G, and let its End be
‘ joined to E, the End of the lower Bone of the
‘ Arm or Cubit AB, which End must be
‘ drawn round the Center of the Articulation
‘ C, describing the Arc BH ; I say, that Na-
‘ ture neither could, nor ought to fasten the
‘ tendinous End E near the End B of the Bone
‘ AB. For if it could be, let us suppose the
‘ Connexion E to be made near B the Wrist,
X
and :

Annotat. " End ; therefore the Power : will always be to the Resistance :: as the
 Lect. III. " greater Distance of the Resistance : to the lesser Distance of the Power
 " from the same fix'd Point ; consequently the moving Power is greater
 " than the Resistance.

PROPOSITION XXII.

" The first Enquiry of the absolute apparent Force, which can be exerted by the two Muscles, the Biceps and Brachieus, bending the Cubit (or lower Bone of the Arm) when the whole Arm is in a supine and horizontal Situation ; which is greater than twenty times the Weight that is sustain'd by them, and exceeds the Force of 560 Pound Weight. Plate 15.

Pl. 15. F. 4. Fig. 4.

" Let the Humerus EA, and the Cubit and Hand AB, be almost in a right and horizontal Line, but supine (that is, with the Elbow downwards) and let the Cord GB fold about the Ends of the Fingers of the expanded Hand G, to which Cord at G hangs the Weight R, which must be encreas'd by degrees, till the Excess of the moving Power of the Muscles DC becomes wholly insensible, and they can sustain no more Weight than R, but be just able to hold it with a Force brought to be equal ; then we may judge that the Momenta of the Powers of the Muscle, and of the Weight, are wholly equal, neither of those

" and then the Tendon, and the Muscle DE is either loose and may be separated from the Limb and the Bones DAB ; or it is bound down to it by some Ligament or Fascia as R ; if the first, the Consequence will be as follows : Because the Bone AB cannot be turn'd up towards FG quite to the Situation AH, unless it be drawn by the Contraction of the muscular Cord DE, in which Case its Length DE, in order to be shorten'd to DM, must become less than the eighth Part of DE, which shortening in the Arm will be of above a Foot and an half ; this will not only be troublesome, but even impossible. It would be troublesome, because the Breadth and Thickness of the Arm would be vastly encreas'd to take in the Dimensions CM equal to CE ; so that the Arm would upon this account only become as big as the Belly of the Animal, which monstrous Thickness would hinder the rest of the Motions of the Arm and of the Animal ; then because the Structure of a Muscle is such that it can be contracted but a little, seldom above two or three Fingers Breadth ; such a fastening of the Muscle,

" which requires so prodigious a Contraction (namely of above a Foot and a half) would be altogether impossible. But the Absurdity of such a Position will most evidently appear, if we suppose the Bone AB to be the Humerus (or upper Bone) of the left Arm, which is to be mov'd every Way round the Joynt of the Shoulder suppos'd at D ; that it may be brought to the Breast, it is plain that it must be drawn by the Muscle ED fix'd to D the right Side of the Breast ; another Muscle to raise it up must be fix'd on the top of the Head, and the Muscle to bring it down must have its Origin in the lower Part of the Belly ; which Muscles, together with those of the right Arm, require a vast swell'd Space like a great Tun ; and the same would be requir'd for the Muscles of the Feet, which would make a Man so far from being well joynted and clean limb'd ; that he would be a ridiculous unweildy Mass, unfit for Motion and handling of Bodies : Therefore such a Shape was entirely to be rejected.

Forces

“ Forces overcoming the other. Now Experience shews us, that in a ro- Annotat.
 “ buſt young Man the Weight R does not exceed 26 Pounds, to which Lect. III.
 “ muſt be added the whole Weight of the Cubit and the Hand, which
 “ are nearly equal to 4 Pounds, and this Weight acts, not in the End of
 “ the Leaver as at B, but in the intermediate Place H, namely where
 “ its Center of Gravity is; therefore if another Weight of 2 Pounds be
 “ ſuſpended at B, which has the ſame Proportion to the Weight of the
 “ whole Cubit as the Diſtance OH to OB, we ſhall have for our Lea-
 “ ver an indiviſible Line and without Weight, at whoſe End B are ſuſ-
 “ pended two Weights, namely R and the Weight of the Cubit, that is
 “ in all 28 Pounds; and then becauſe the Direction CD of the Tendon
 “ of the Muſcle that draws makes a very acute Angle with the Line
 “ CO, becauſe the Tendon of the Muſcle exactly touches the Head of
 “ the Joynt A, we muſt from the fix'd Point or *Fulcrum* O draw the
 “ right Line OI, perpendicular to CI the Direction of the Tendon, and
 “ then from the Principles above demonſtrated it will appear, that the
 “ *Power drawing the Muſcle DC : has the ſame Proportion to the Reſiſ-*
 “ *tance of the Weight R together with the additional Weight above mentio-*
 “ *ned :: as the Diſtance OB : has to the Diſtance IO ;* but by a ſtriſt
 “ Examination it appears that OB the Length of the Cubit and Hand
 “ is more than twenty times greater than the Semidiameter of IO the
 “ Head of the Bone. Therefore the Strength and Power drawing the
 “ Muſcle DC is above twenty times greater than the Weight R and the
 “ additional Weight; and ſince the 2 Weights are equivalent to 28 Pounds,
 “ therefore the apparent Force with which the Muſcle draws the Cubit
 “ and endeavours to bend the Elbow, is greater than the Force of 560
 “ Pounds.

“ PROPOSITION XXIII.

“ *To find the Force which the ſaid Muſcles exert, when the Humerus or*
 “ *upper Part of the Arm is perpendicular to the Horizon, and the Cubit is*
 “ *parallel to the Horizon. Plate 15. Fig. 5.* Pl. 15. F. 5:


“ Secondly, Let EA be the *Humerus*, and AB the *Cubit* making
 “ a right Angle with each other, the *Humerus* being perpendicular and
 “ the *Cubit* ſtill horizontal: In that Poſition the Length of the Lea-
 “ ver OB ſtill remains the ſame, and now at its End B is ſuſtain'd the
 “ great Weight of 33 Pounds (as appears by making the Experiment) by
 “ the ſame Muſcles DC; but becauſe the Angle ICO made by the Ten-
 “ don with the Bone OC is leſs acute than in the foregoing horizontal
 “ Situation of the ſame two Bones, becauſe when the *Humerus* EA is bent
 “ towards the *Cubit* AB, the Tendon of the Muſcle DC adhering to
 “ the *Humerus* is alſo bent; yet the Angle ICO does not become a right
 “ one, becauſe the Tendon at I is firmly bound with membranous *Faſcie*
 “ and the outward Skin, which Ligaments ſerve as a Pulley to keep the
 “ Tendon

Annotat. " Tendon towards A the Angle of the Joynt ; but yet the Tendon IC
 Lect. III. " is not so closely bound down at I but that it rises a little, and there-
 fore the right Line OI perpendicular to the Direction of the Tendon
 " CI becomes sensibly longer than in the foregoing Case, as we may
 " find by feeling our Arm ; and therefore the Distance OB will bear a
 " less Proportion to IO, than it was found to do in the former Si-
 " tuation ; but whatever Proportion the said Distances have, the
 " same reciprocally will the Force contracting the Muscle DC, and
 " drawing the Bone, have to the Resistance of the Weight R and
 " the Weight of the *Cubit* together : Therefore that Force will bear a
 " less Proportion to this Resistance, than 20 to 1 ; and if it appear'd by
 " the foregoing Enquiry that the greatest Force of the *Biceps* and *Brachieus*
 " Muscles was equal to the Force of 560 Pounds ; it will appear
 " from the present Enquiry, in which the great Weight R is of 33
 " Pounds, and, taking in the Weight of the *Cubit*, equal to 35 Pounds,
 " that the Distance OI is only a sixteenth Part of the Distance OB, and
 " not, as before, a twentieth Part of it ; and therefore that the Distance
 " IO being sensibly increas'd, there must of consequence be rais'd a
 " greater Weight now, namely 35 Pounds, by those Muscles.
 " Yet here we must take notice, that tho' by reason of the bending of
 " the Limb EAB the Muscles are not stretched as before, but must be
 " in some measure relax'd ; yet the moving Force of each Muscle has
 " a less Power of contracting, because really the Muscles DC are not
 " both fix'd to the Top of the *Humerus*, but the *Biceps* is fasten'd to the
 " *Scapula* or Shoulder-bone HLE at L, but the *Brachieus* to the middle
 " of the *Humerus* ; and because the *Scapula* HLE is always in the same
 " and transverse Situation, the *Humerus* EA revolving about the Center
 " E of its Articulation, must make the Angle LEO with the *Scapula*
 " the less acute the more the *Humerus* is bent downwards, and then the
 " Origin of the *Biceps* Muscle D is more rais'd and recedes farther from
 " the top of E the Head of the Bone, because the Length of the Line
 " LDI subtending the Angle LEO is increas'd, and therefore the fore-
 " said Muscle is so much the more stretch'd as the *Humerus* is bent down-
 " wards : Therefore tho' by reason of the Angle EOB the *Brachieus*
 " Muscle be relaxed ; yet the *Biceps* may be so much the more stretch'd
 " by reason of the Elevation of the Point D above the Head of the
 " *Humerus*.

" PROPOSITION XXIV.

" Hence probably may be singly found the apparent absolute Forces of the Bi-
 " ceps Muscle, which is equivalent to 300 Pounds, and of the Brachieus, which
 Pl. 15. F. 6. " is equal to the Force of 260 Pounds. Plate 15. Fig. 6.

" Let the *Humerus* OE be bent backward in order to make the Angle
 " HEO as acute as may be, and likewise let the *Cubit* AB be inflected so
 as

“ as to become parallel to the upper Line of the *Scapula* HL, and then Annotat.
 “ the alternate acute Angles HDI and CID will be equal to one another, Lect. III.
 “ and then, as much as the *Biceps* Muscle DIC is relax’d on account of 
 “ the Acuteness of the concave Angle COE, just so much is it drawn and
 “ stretch’d on account of the convex Angle HDO ; therefore the natural
 “ Tension of the *Biceps* Muscle is in no wise alter’d, and remains exactly
 “ of the same Length as it had in the horizontal Situation of the whole
 “ Arm ; and as it suffers no Relaxation, it will have the same Force of
 “ contracting itself as it exerted the horizontal Position. But the *Brachieus*
 “ Muscle has not the same Advantage, its Origin being in the middle of the
 “ *Humerus* at F, and its End or Insertion at I near the Head of the *Cu-*
 “ *bit* ; and because the Angle EOC is acute, therefore the *Brachieus* Mus-
 “ cle must suffer the greatest Relaxation, and therefore exert little or no mo-
 “ tive Force : In this Case therefore one may find the moving Force of the
 “ *Biceps* alone ; (that is, if the Distance OI from the Tendon to the Center
 “ of the Head of the *Cubit* be not varied.) Let us suppose the Weight
 “ R sustain’d in that Situation together with the Weight of the *Cubit* to be
 “ equal to 25 Pounds, and because the Distance IO is almost a twelfth Part
 “ of the *Radius* and the Hand BO, therefore the absolute Force of the
 “ *Biceps* Muscle will be twelve times greater than the appended Weight R
 “ together with the Weight of the *Cubit*, that is, it will be equal to the
 “ Force of 300 Pounds, when the *Brachieus* exerts no Force by reason of
 “ its very great Relaxation ; then because the joint Forces of the two Mus-
 “ cles, the *Biceps* and the *Brachieus*, working together in the first Experi-
 “ ment, were equal to the Force of 560 Pounds ; if from that Force we
 “ subtract the moving Force of the *Biceps* alone, just found to be of 300
 “ Pounds, the remaining Force of 260 Pounds will be that which was ex-
 “ erted by the *Brachieus* Muscle, and that was the Thing to be enquir’d
 “ into.

“ PROPOSITION XXV.

“ *To find what Force the same Muscles exert when the Cubit hangs down-*
 “ *wards, whilst the Humerus is kept perpendicular to the Horizon.* Plate 15. Pl. 15. F. 7.
 “ Fig. 7.

“ Now let the *Humerus* EA, and the *Cubit* AB be in one direct Line
 “ and perpendicular to the Horizon ; then the greatest Weight to be
 “ suspended at B, might be almost immense, if the Strength and Tenacity
 “ of the Ligaments could always resist and was wholly insuperable.

“ If afterwards the *Cubit* be a little inflected, so as to make an obtuse
 “ Angle EAB with the *Humerus* which is now perpendicular to the Hori-
 “ zon, and an acute Angle BAK, with the horizontal Line OK, then
 “ indeed may the great Weight R be very much increas’d, because if
 “ from B the Line BK be drawn perpendicular to the horizontal Line
 “ AK, then the Weight R drawing the Leaver AB obliquely, acts in
 “ the


Annotat. " the same manner as if it had been suspended in the Point K of the
 Lect. III. " Leaver OK; and therefore we have now a new Leaver OK shorter
 " than OB; but the Force of the Muscle raising the Leaver draws from
 " the Point I having IO for the Distance of its Line of Direction; and
 " consequently the absolute Power that contracts the Muscle (which is al-
 " ways the same) has the same Proportion to the Resistance of the Weight
 " R, as KO has to IO; therefore if KO be only the double of OI,
 " the Weight R which is sustain'd in that Position will be half of the
 " whole absolute moving Force, and therefore equal to 250 Pounds; and
 " if the Distance OK be less than OI, then also will the Weight R be
 " greater than the moving Force of those Muscles.
 " Hence may be gather'd, that in the Flexion, or Elevation of the *Cu-*
 " *bit*, the Effect of the same Force which draws the Muscle is continu-
 " ally diminish'd; because the Length of the Leaver OK is successively
 " encreas'd, and therefore the Weight R must be decreas'd in the same
 " manner.

" PROPOSITION XXVI.

Pl. 15. F. 8. " *To find the Force of the same Muscles when the Arm is plac'd in a su-*
 " *pine horizontal Situation.* Plate 15. Fig. 8.

" The Force of the Muscles bending the *Cubit* may be exerted in an-
 " other manner, namely when the *Cubit* AB being in a supine horizontal
 " Situation must be inflected downwards towards G by the Muscles DC
 " which are now below the *Cubit*; for the Cord BLG being thrown
 " over the Pulley or Wheel ML moveable about a fix'd Axis M, it is
 " evident that whilst the Hand B descends, the Weight R is rais'd, AB
 " being then the Leaver whose *Fulcrum* is O, and the Weight R draws
 " the End B of the Leaver upwards towards L, and the Power of the
 " Muscles DC draws down the Leaver AB from I towards D; and
 " therefore those Things which have been said before will also be veri-
 " fied here, only with this Difference, that in the former Case the End
 " B was drawn downwards, not only by the Resistance of the Weight R,
 " but also the Weight of the whole *Cubit* and Hand; but on the con-
 " trary, here the Weight of the *Cubit* does not act counter to, but helps
 " the Power of the Muscles to draw; because, that as in this Situation
 " the Muscles draw down the *Cubit*, so the *Cubit* also acts downwards by
 " its Gravity, and these two Powers taken together have a *Momentum* e-
 " qual to that of the Weight R; and as in the first Case the Weight of
 " the *Cubit* was added to the Resistance of the heavy Body R, now it is
 " added to the Power of the Muscles; and because the greatest Power
 " of the Muscles DC was shewn to be equal to the Power of 560 Pounds*,
 " therefore if the Leaver AB was of no Weight, when the Distance OB
 " is found to be twenty times the Distance OI, the Weight R ought to
 " be of 28 Pounds; but because the 2 Pounds added to R are in *Æqui-*
 " *librio*

* By the 22d
 Prop. above
 cited.

“ *librio* with the Weight of the *Cubit* AB (that is, as they render it a Annotat.
 “ Leaver without Weight) therefore the Weight with its Adjunct making Le&t. III.
 “ up 30 Pounds will be the greatest Weight that can be sustain’d by the 
 “ Force of the Muscles in that Situation.

“ This may be shewn another Way, because the descending Weight of
 “ the *Cubit* being as 2 Pounds hanging at B, is made to act equally with
 “ a drawing Force of 40 Pounds applied at I (by reason of the reciprocal
 “ twentyfold Proportion) and the proper Force DCI will be of 560
 “ Pounds; therefore this Force, together with the *Momentum* of the *Cu-*
 “ *bit* acting with it, will produce an Effect equal to the whole Weight of
 “ 600 Pounds.

But what appears most wonderful is the Force of the Muscles that move the lower Jaw, which the said *Borelli* considers in the 87th and 88th Proposition of the first Part of his Book abovemention’d, where he shews that those small Muscles, all which taken together do not in a Man exceed the Weight of one Pound, do yet exert a Force equal to 534 Pounds; and in Mastif Dogs, Wolves, Bears and Lions, have a Force vastly superior, to enable them to break large Bones as they do daily in their Feeding.

7. [35. — And if the Arm CW be set streight in a Line with PC, &c. — the Instrument will plainly appear to be a Leaver of the first Kind. Tho’ the bended Leaver is not an Instrument of common Use, except in the Hammer and Tools of that Kind, yet the Consideration of it is very necessary in the Explanation of several Machines that virtually contain such a Leaver; especially in statical Propositions, of which we will here give some Examples. The Cases of the *Inclin’d Plane* and the Wedge * may ^{48, 49, 50,} be clearly solv’d this Way. As for example, when the Weight P ^{51, 52,} sustains the Weight W upon the inclin’d Plane AB drawing the Center of the Weight in the Line MW parallel to the Plane, one may consider in the said Weight the *bended Leaver* W T n whose longest *Brachium* is W T and the shortest T n. Now, as the Line of Direction of the Power is MW parallel to the Plane, TW acted upon at right Angles must be the Distance of the Power; and as W n is the Line of Direction of the Weight, n T at right Angles with it must be the Distance of the Weight; consequently as n T (the short Arm of the bended Leaver) : is to W T (its long Arm) :: so is the Power P : to the Weight W. Now, as the Triangles W T n and ABC are similar, the Power thus consider’d : is to the Weight :: as BC the Height of the Plane : to its Length AB, which was prov’d in its Place *. But if the Power had drawn the Weight in a Line parallel * ^{49.} to the Base of the Plane, which is the Case of the Wedge, its Effect might be thus explain’d. As the Power acts obliquely at the End of the long *Brachium* of the bended Leaver W T n, its acting Distance must be found by drawing from the Center of Motion T, T o perpendicular to the Line of Direction of the Power, which now must be consider’d as the long *Brachium*.

Annotat. *Brachium* of the Leaver, whilst nT still is the short Arm and acted upon
 Leſt. III. by the Weight at right Angles. Therefore the Quantity or Intenſity of
 the Power Π will now be found by this Analogy, viz.

As the long Arm of the Leaver, now To :

Is to the ſhort Arm Tn ::

So is W the Weight :

To Π the Power ; or the Baſe AC : to the Height CB .

* 48, 49, 50,
 51, 52.

When I conſider'd the *Inclin'd Plane* as a mechanical Organ*, I only took notice of two Applications of the Power, the one with its Line of Direction parallel to the Plane, and the other (which reduc'd it to the Wedge) with its Line of Direction parallel to the Baſe of the Triangle, that is, inclin'd to the Plane as much as the Plane is inclin'd to the Horizon, the Angle WBA (call'd the *Angle of Traction***) being equal to the Angle BAC , becauſe of the Parallels WB, AC ; but as in compound Engines, and in the Uſe of Carriages the Angle of Traction or Inclination of the Line of Direction of the Power to the Plane is very variable, we give this Way of conſidering a *bended Leaver* in the Body to be drawn up for the Solution of all Caſes, which will appear by the following Example.

* L. 2. 45.

Let DLB be the Angle of Traction, as when the Power Π draws over the Pulley p in the Line pL . WTn is a bended Leaver whole Center of Motion is T the Point where the Ball W touches the Plane, nT the ſhort Arm of the Leaver on whoſe Extremity n may be conſider'd the Weight as ſupported and preſſing at right Angles, becauſe its Line of Direction goes through n *; and WT the long Arm of the Leaver to which the Power is apply'd obliquely. But as Tz perpendicular to the Line of Direction of the Power is its *acting Diſtance*, we may conſider Tz as the long Arm of the Leaver. Then the whole Caſe will be ſolv'd by this Analogy, viz.

As the long Arm of the Leaver, now zT :

Is to the ſhort Arm Tn ::


So is W , the Weight :

To Π the Power.

Since one may find this laſt bended Leaver, in any Direction of the Power, that is, in any Angle of Traction, this general Rule for all Directions of the Power may be deduced from it.

** The Angle of Traction is the Angle the Power draws in a Line of Direction parallel to the Plane, there can be no Angle of Traction.

*As the Sine of the Angle of Inclination of the Plane :
To the Sine Complement of the Angle of Traction ::
So is the Power :
To the Weight.*

Annotat.
Lect. III.


But to prove how the Rule is deduced from the bended Leaver, we must shew that nT , the short Arm of the said Leaver, is always to its long Arm zT , as the Sine of Inclination, to the Sine Complement of the Angle of Traction; or that CB in the Triangle ABC : is to Lz in the Triangle TLz :: as nT : is to Tz ; and then we shall draw some useful Corollaries from our general Rule.

D E M O N S T R A T I O N.

Since WT is perpendicular to AB , and the Angles WqT and AqE equal (because they are vertical Angles, the Triangle AqE (rectangular at E) having two Angles equal to two Angles in the Triangle WqT , the third Angle A must be equal to the third Angle qWT , consequently the Triangles will be equiangular, and therefore similar (by 4. 6. *Euch.*) and as qE is the Sine of Inclination for the Radius Aq (because in its similar BC is the Sine of Inclination for the Radius AB) so also will qT be the Sine of Inclination for the Radius Wq . But as Tn which is parallel to the Horizon falls upon Wq , which being the Line of Direction of the Weight, is perpendicular to it, the Triangle WTn is similar to WqT (by 8. 6. *Euch.*) and thus Tn the short Arm of our bended Leaver becomes the Sine of Inclination for the Radius WT : Which was one of the Things to be demonstrated.


Further, as Tz is by Construction perpendicular to LW , the Triangle WzT is similar to LWT (by 8. 6. *Euch.*) therefore the Angle zTW is equal to WLT the Angle of Traction; so that as LT is the Sine Complement of the Angle of Traction for the Radius LW , TZ (the long Arm of our bended Leaver) must be the Sine Complement of the Angle of Traction for the Radius TW . Which was the other Thing to be demonstrated.

C O R O L L A R Y I.

Here follows, that when the Line of Direction is parallel to the Plane, the Power is the least that can be for that Inclination of the Plane, because then the Sine Complement of the Angle of Traction is changed into the Radius or Whole Sine, that is, TZ becomes TW , or the long Arm of the Leaver WT being acted upon at right Angles expresses the acting Distance of the Power; or, strictly speaking, the Angle of Traction vanishes. But if the Power should draw directly up in the Line We , it

Y

must

Annotat. must be equal to the Weight; because then WqT being the Angle of
 Lect. III. Traction, its *Sine Complement* (qW being *Radius*) would be nT , which is
 equal to the *Sine* of Inclination here express'd by the said Line. Hence
 it is plain, that if an Horse draws a Burthen up Hill by means of a Cart
 Pl. 10. F. 14. or any rolling Machine, he will draw it with so much the more Ease, as
 the Line of Direction by which he draws the Load comes nearer to a parallel
 to the Slope of the Mountain along which he draws.

COROLLARY II.

It follows also, that if the Line of Direction, as BW and wW parallel to AB , make the Angle wWB equal to the Angle wWD , the Power applied at B will be equal to the Power applied at D ; because in such a Case the Angles of Traction WLB , WBL will be equal, since by 29.1. *Eucl.* the Angle WLB is equal to its external opposite DWw , which is suppos'd equal to the Angle wWB , and consequently to its alternate WBL . Whence it follows, that if the Line of Direction of the Power, as Wa , and the Line WT perpendicular to AB , make the Angle TWa equal to the Angle of Inclination A (or qWT) the Power applied at a will be equal to the Weight by *Cor. 1.* because then the Angle of Traction Wal is equal to the Complement of the Angle of Inclination A ; that is, Wal is equal to $Wqa = ABC$.

COROLLARY III.

Lastly, it follows also, that if the Line of Direction of the Power be WT at right Angles to the inclin'd Plane AB , which makes the Angle of Traction a Right; the Power applied at T or in any part of the Line WT must be infinite. That is as much as to say, that a Power that should draw the Weight D directly from the Plane, or directly against it, would not keep it there, let its Intensity or Force be ever so great, because in that Case the *Sine* of the Complement of the Angle of Traction is reduced to nothing, or being *infinitely small*, the Power applied at T must be *infinitely great*; since, by what we have shewn, that Power : must be to the Weight :: as the *Sine* of the Angle of Inclination : is to the *Sine Complement* of the Angle of Traction.

* 28, 35, 45. Before I quit this Subject of Bodies supported or drawn on inclin'd Planes, I beg leave to apply what has been said in the second Lecture *, to shew generally why a Body will be sustain'd on an inclin'd Plane by another Body of less Weight (if the former be drawn in a Line of Direction parallel to the Plane, and the latter hangs perpendicularly) when the Weight of the great Body : is to the Weight of the little one :: as the Length of the Plane : is to its Height. The whole is deduced from this Principle laid down and explain'd in the second Lecture, *viz.* That if the Center of Gravity of a System of Bodies does not descend, the Bodies cannot descend.

Now

Now to apply this to our Purpose. ADB^* is an inclin'd Plane, whose Height is DB . If by means of a Pulley P the Weight w hanging perpendicularly does by a String hold the Weight W on the inclin'd Plane, and those Weights are to one another, as the Length of the Plane to its Height (which here is as 2 to 1) they will remain at Rest (that is, keep each other in *Æquilibrium*) whatever Part of the Plane W is laid on. First, let the Situation of the Bodies be W and w ; draw the Line mn which joyns their Centers of Gravity, and having found their common Center of Gravity at C^* ($Cn = 2 Cm$) draw Hh an horizontal Line through that common Center; and then whatever Position the Bodies are in, or whatever Part of the Plane W be plac'd on, their common Center of Gravity will still be in the horizontal Line Hh : If W be remov'd to V , w will fall to u , and the common Center of Gravity will be at k : If the Centers of the Bodies be at E and e , their common Center of Gravity will be at K , still in the same Line Hh , which may be easily prov'd, because the Triangles nhC , mHC , hpk , rok , &c. are all similar. Since therefore there is no Position of W on the Plane and of w in the Perpendicular wq which can alter the Height of the common Center of Gravity of the Bodies, they must be in *Æquilibrium*, because they can't fall unless their Center of Gravity descend. Q. E. D.

C O R O L L A R Y.

Hence also appears the Reason why two unequal Bodies will sustain each other upon unequal Planes of the same Height, whose Lengths are to one another reciprocally as the Bodies. For example, let the Weights F and G^* , join'd by the Cord FPG running over the Pulley P , be to one another in Weight as the Planes AB , BD (whose common Height is BE) on which they respectively rest. Find their common Center of Gravity C , and through C draw an horizontal Line; then you will find as before, that however you alter their Situations on their respective Planes, their common Center of Gravity will still be found in the same horizontal Line that goes through C , &c.

8. [37. — *An upper or fix'd Pulley adds no Force to the Power, but only prevents the Friction by making the Rope run easily; and so much the more as the Sheever is bigger than the Center-Pin upon which it turns.*] How much the Friction of a Roller or upper Pulley is diminish'd in proportion as the Center-Pin, or the Gudgeons are less in Diameter than the Wheel or Roller I shall exactly shew in the next Lecture. Now I shall take notice of the Diminution of Pressure upon the Center-Pin of a Pulley which always happens in a certain Proportion of the Weights which hang on each side, when they are in Motion, and that without Regard to the Bigness of the Pin, which in this Case may be consider'd as a Line. The Thing in general is comprehended in the following

Annotat.
Lect. III.

P R O P O S I T I O N .

When a String or Rope runs over a single Pulley or Roller by the Descent of the preponderating Weight (the other Weight rising at the same time) the Pressure on the Axis of the Pulley is always equal to the Quadruple of the Product of the Weights multiplied into one another) and divided by the Sum of the same Weights.

Pl. 15. F. 11.

Plate 15. Figure 11.

The Pulley is DE its Center C, the Weights p and q , and the running Rope $pD \div E q$. I say, that the Pressure on the Axis or Center C is =

$$\frac{4pq}{p+q}.$$

Pl. 15. F. 12.

Plate 15. Fig. 12.

The Pulley being represented in this Figure by the same Letters as before, draw at pleasure the Lines Dd and Ee in which the Weights rise or fall. Draw from the Center of the Pulley Ck parallel to the said Lines; then thro' the Point o taken any where in the Line Ck draw the horizontal Line pq and taking the Distance pP equal to qQ draw the oblique Line PoQ . From the Point c the common Center of Gravity of the Weights represented by the Letters p, q , and suppos'd to hang at those Points, let fall the Perpendicular cg till it meets the Line PQ . If we consider pq as a Balance of unequal *Brachia* whose Lengths are in the reciprocal *Ratio* of q to p , so that the Weights p and q shall upon it keep one another in *Æquilibrio*, we may call cq the Leaver of the greater Weight q , and pc the Leaver of the lesser Weight p ; and if we suppose the Weight q to descend as far as Q whilst the Weight p rises as far as P (through an equal Space) because Pg is to gQ as pc to cq , the Line cg will represent the Descent of the common Center of Gravity of the two Weights.

D E M O N S T R A T I O N .

The Sum of the Weights $p+q$: is to their Difference $q-p$:: as the sum of their Leavers $pc+cg$: to their Difference $pc-cg$: and consequently :: as half the Sum of their Leavers oc : to half their Difference oc ; and likewise by 2 and 4. 6. *Euch.*) :: as qQ the Velocity of the Weight descending : to cg the Velocity of the Center of Gravity descending. Now taking out of q a Weight equal to p , there remains only

$$q-p$$

$q - p$ to give Motion to the Weights by the natural Gravitation of it. Annotat. But the *Momentum* of $q - p$ by the natural Gravitation is $qv - pv$ (v being taken for the natural Velocity of the Weight falling) therefore the Velocity express'd by qQ will be $\frac{qv - pv}{q + p}$; but before, we had this Analogy $p + q : q - p :: qQ : cg$; therefore we have $p + q : q - p :: \frac{qv - pv}{q + p} : cg$ the Velocity of the Center of Gravity, which will be

$$\frac{q^2 - 2pq + p^2 \times v}{p^2 + 2pq + q^2}.$$

Therefore the *Momentum* of the two Bodies will be that Velocity multiplied by the two Bodies, and consequently the *Momentum* of the Fall

of the two Bodies will be $\frac{q^2 - 2pq + p^2 \times v}{q + p}$. Now if the two Bodies

$q + p$ fell wholly, their *Momentum* would be $qv + pv$; but as we must

subtract only that which falls of the Bodies, viz. $\frac{q^2 - 2pq + p^2 \times v}{q + p}$, there

is a Remainder of the Bodies which does not fall, and that Remainder of consequence must rest upon the Center and press the Axis. Now this Remainder, by making the Subtraction, appears to be $\frac{4pq}{p + q}$. Q. E. D.

Scholium.

Hence it follows, that if p be equal to q (in which Case there will be no Motion of the Weights, the Pressure will be $2p = p + q$; and if q be infinite, the Pressure is $4p$).

To try this practically, I made the following.

EXPERIMENT. Pl. 15. Fig. 13.

Pl. 15. F. 13.

I screw'd a very nice Pulley D to the bottom of the Scale b of the Balance AB moveable round the Center C, and having balanc'd the Pulley by Weights in the Scale a , I fasten'd to a String the two Weights p and q weighing 2 and 6 Ounces; then having put the String over the Pulley D, I plac'd 6 Ounces and one Penny Weight in the Scale a kept from rising by a Thread, which join'd in bottom to the fix'd Hook H, whilst the whole Balance hung by another fix'd Hook at M. Supporting the great Weight q by a Ruler held at e horizontally under it, I on the sudden remov'd e to give q leave to descend; which when it did, the Pressure on the Axis of the Pulley D was so much diminish'd (during the Rise

Annotat. Rise of p and Descent of q) that the Balance a sunk by the Action of the little Weight d , even when it was a great deal less than a Penny Weight, as plainly appear'd by the Thread Ha growing slack.

When q was = 12 Ounces and $p = 3$, the Counterpoise in the opposite Scale was 9 Ounces and 12 Penny Weight according to the Theory, and the Experiment agreed with it, there being a visible Descent of the Scale a , by the Addition of the little Weight d , even when it was less than the 300th Part of q , &c. Take q and p in any other Proportion, and still the Experiments will agree with the Theory.

This Way of considering the Pressure upon the Axis may also be applied to the Axle in the Wheel in the following manner.

Pl. 15. F. 14. Let ACB^ be an Axis in Peritrochio, whose Wheel is AB , C the Center or Axis of Motion and AX the Axle, q one Weight (commonly the greater Weight) and p the other, the Line qp , as a Leaver unequally divided at o , represents by its Part qo the Arm or Semidiameter of the Axle AC , and by its Part op the Arm or Semidiameter of the Wheel CB .

The Bodies p, q , being in the Situation represented in the Figure, and pb greater than qa , the Part of p which makes an *Equilibrium* with q will be $\frac{qa}{b}$, and consequently there will be left of p only $p - \frac{qa}{b}$ or

$\frac{pb - qa}{b} \times V$ (making V equal to the Velocity with which Bodies fall) to give Motion to the two Bodies; because the Remainder of the Weight of the two Bodies, $pV + qV$, namely $\frac{qbV + qaV}{b}$ of the two Bodies of

Equilibrium $q + \frac{qa}{b}$ will remain pressing on the Axis of the Machine. Let

u be the Velocity of p , the Velocity of q will be $\frac{au}{b}$, and its Momentum will be $\frac{qa u}{b}$, which added to pu the Momentum of p gives $\frac{pb u + qa u}{b}$ equal to the whole Momentum produced by the Force or Momentum $\frac{pbV - qaV}{b}$. Therefore since the Momentum produc'd is always equal

to the Momentum which produces it, we have $\frac{pb u + qa u}{b} = \frac{pbV - qaV}{b}$

which gives $u = \frac{pbV - qaV}{pb + qa}$ and $\frac{au}{b} = \frac{pbaV - qa aV}{pbb + qab}$ which

gives the Momentum of $q = \frac{qpba - qqa a}{pbb + qab} \times V$, which Momentum p

will give it as it bears on the Axis, and consequently the Axis must suffer the Pressure of it. And as q , by its Reaction upon p , will make it lose
as

as much (bearing also upon the Axis) or will as much retard its Descent, Annotat. the Axis must of necessity by this equal Action and Reaction bear the *Left. III.*

double Pressure $\frac{2qpba - 2qqa}{pbb + qab} \times V$; to which adding the Weight

$\frac{qb + qa}{b} \times V$ of the two Bodies of *Æquilibrium*, the whole Pressure on

the Axis will be equal to $\frac{qpbb + qqab + 3pbqa - qqa}{pbb + qab} \times V$ or (di-

viding by the Denominator as far as we can) it will be equal to $q +$

$\frac{3pbqa - qqa}{pbb + qab} \times V$. Therefore the Axis bears as much Weight as if

it sustain'd the Quantity of Matter $q + \frac{3pbqa - qqa}{pbb + qab}$.

C O R O L L A R Y I.

If $pb = qa$, the Pressure is $q + p$, or $q + \frac{qa}{b}$ (according as pb is set

down for qa , or the Reverse in the Form $\frac{qpbb + qqab + 3pbqa - qqa}{pbb + qab}$)

because in that Case $\frac{qa}{b} = p$.

C O R O L L A R Y II.

If p be infinite, the Pressure is $q + \frac{3qa}{b}$, or the Weight q together with three times the Weight that is able to keep it in *Æquilibrio* at the Distance of p , as in the Pulley above mention'd, where the Pressure is $\frac{4qp}{p + q}$. If q be infinite, the Pressure then is $4p$; and if b be infinite, the Pressure is only equal to q .

N. B. This may be given as a general Rule for the Pressure on the Axis either of a Pulley or of a Wheel and Axle, by the two Bodies acting against each other, viz.

As the Momentum of the two Bodies falling freely :

To the Momentum which is lost when they act on each other by means of the Machine ::

So is the whole Weight of the Bodies :

To the Weight pressing upon the Axis.

Annotat.

Lect. III.

9. [41. — While 2 goes down to a, 1 goes up to B just twice as far, &c.] There is another Case of raising a Weight by separated Pulleys, which I omitted here. It is mention'd by Dr. Pemberton in his *View of Sir ISAAC NEWTON's Philosophy*. I shall first give his Solution of it, and then shew how easily it may be reduced to our Rules, by proving, that however that Case is varied, there will be still a reciprocal Proportion between the

* Pl. 15. F. 15. Power and the Weight. The Weight W is sustain'd by the Power P by means of the three Pulleys C, D, E , of which D is fix'd and the others moveable, and a Rope goes from the Weight to each Pulley as represented in the Figure. "To explain the Effect of Pulleys thus applied,

* Pl. 15. F. 16. "it will be proper to consider different Weights hanging as in Fig. 16*. "Here if the Power and Weights balance each other, the Power P is "equal to the Weight w ; the Weight W is equal to twice the Power P or "twice the Weight w ; and for the same reason the Weight W is equal to "twice the Weight w , or equal to four times the Power P . It is evident therefore, that all the three Weights w, w, W together, are equal "to seven times the Power P . But if these three Weights were join'd "in one, they would produce the Case of Fig. 15 †. So that in that Figure the Weight W , where there are three Pulleys, is seven times the "Power P . If there had been but two Pulleys, the Weight would have "been three times the Power; and if there had been four Pulleys, the "Weight would have been fifteen times the Power.

† Pl. 15. F. 15. To explain this our Way, let us consider the Weight W to be rais'd one Inch, as from the horizontal Line AB to the horizontal Line ab , and from the Make of the Machine find what must be the Velocity of the Power. First then, the Point F of the Rope going over the Pulley C , must descend one Inch, viz. from the Line Ff to Gg (because W fasten'd to the said Rope rises one Inch by Supposition) and the Pulley D fasten'd to the said Rope DF must also descend one Inch. From the Descent of the Pulley D one Inch the Point H of its Rope must descend two Inches, as it is supplied from both Sides of the Pulley; but an Inch more of the said Rope must be supplied by the Rise of the Weight W ; therefore the Point H will descend three Inches, or from Hh to Ii . Lastly, as the Pulley E descends three Inches because it hangs by the Rope HI , the Point K of the Rope KP (being supplied from both Sides of the Pulley E) must descend 6 Inches on that Account, and one Inch more on account of the Rise of W : Therefore the Point K of the last Rope by which the Power P pulls, will descend seven Inches, viz. from the Line Kk to the Line Ll , whereby the Power also will descend the same Distance, namely from P to p . Consequently one Pound at P instead of the Hand will sustain the Weight W seven times as big, 7×1 being equal to 1×7 . Therefore in this Combination of Pulleys, as well as in all others, and indeed in all mechanical Engines (as we have often said) where there is an *Æquilibrium*, there will be a reciprocal Proportion between the Intensities of the Power and Weight and their Velocities.

10. [41. —

10. [41. — *The Ropes, &c. (applied to Pullies) are always suppos'd parallel, except where it is otherwise express'd.*] Tho' in a Combination of Pullies, where the last Pulley is a fix'd one as in the 4th, 5th, 6th, 7th, 8th and 9th Figures of Plate 10*; the Force exerted (supposing a Man* Pl. 10. F. 4. or Men to draw) is the same in whatever Direction the Power draws the running Rope; yet if the Ropes that are applied to the Block or Blocks which come up with the Weight, are not parallel, Force will be lost in Proportion to their Obliquity.

Suppose the Weight W^* together with the lower or moveable Pulley* Pl. 15. F. 17. C, from whose Center it hangs, to weigh 6 Pounds; if it was suspended at c , it would require a Force equal to 6 Pounds to support it; and therefore if you suppose two upper or fix'd Pullies as A, B, to have over them a Rope, at each End of which hangs a 3 Pound Weight, whilst the middle of the Rope comes under the Pulley C, it is evident that the two Weights (or rather Powers P and P, being both together equal to the Weight, must also support and keep it in *Æquilibrio*. Now, since P and P balance one another, if P be taken away and the Rope be made fast at a , P alone will support the Weight W, as we have said and explain'd already†. And this will appear more evident, if we reduce the Pulley C to † 37. a Leaver after the manner shewn in the second Note of this Lecture*. In * L. 3. Ann. this Case mn is a Leaver of the second Kind, in which the Center of Motion or *Fulcrum* is at n , the Weight W draws at right Angles at o with the Distance on , whilst the Power with the double Distance mn draws also at right Angles in the Direction mB . Now, if the Pulley B be removed to b , the Direction of the Power will be changed and become bm , consequently its Force will be diminish'd in proportion to the Obliquity of its Direction; that is, the Power able to sustain the Weight in the Direction bm : is to the Power which sustains it in the Direction mB :: as bm : is to Bm †.

From this Consideration may be deduced this general Rule for knowing the Intensity of the Power or Powers, which drawing obliquely over fix'd Pullies cause a Weight hanging from the Center of a moveable Pulley to rise directly up.


As twice the Tangent of the Angle of Inclination (that is, the Angle made by the Line of Direction of the Power, which is the oblique Rope) with the Horizon :

To the Secant of the said Angle ::

So is the Weight, when one End of the Rope is fix'd :

To the Power drawing obliquely.

But if two Powers (one at each End of the Rope) be made use of, then the Analogy will stand thus :

Annotat.
Lect. III.


*As twice the Tangent of the Angle of Inclination :
To twice the Secant of that Angle ::
So is the Weight :
To the two Powers taken together.*

P R E P A R A T I O N .

Pl. 15. F. 17. Let the Pullies A, B, be remov'd to a, b , and the Line bm and ao produc'd till they meet at C; join the Centers of the Pullies a, b , by the horizontal Line ab ; and from the Point E taken at pleasure in any Place of the oblique Rope draw ED parallel to ab , and making $Dc = DC$ draw EC: Draw *or* perpendicular to CB.

D E M O N S T R A T I O N .


* L. 2. 22. Since cC is the Line of Direction of the Weight W it must be perpendicular to ab * the horizontal Line, and consequently parallel to Bm ; therefore the Triangle cCb is similar to the Triangle Bmb (by 4. 6. Eucl.) and for the same reason DEC is similar to cbC , and cDE is also similar to them and equal to DCE, because of the right Angles at D, and the common Side DE and equal Sides $DC = Dc$. Besides, the Triangle aCc is similar to the others abovemention'd, because as the Weight W (or its Center of Gravity) descends as low as it can, aCb must be an *Isoceles* Triangle bisected by the Line of Direction cC . Now if Cc represents the Intensity of the Weight hanging on the Center of the lower Pulley C, its Half DC will represent the Intensity of the Power drawing directly or at right Angles to the Leaver mn , and CE its Intensity when drawing obliquely; and since the Angle $DEC = cbC$, EC the Secant of DEC will be the Secant of the Angle of Inclination; and DC the Tangent of DEC will be the Tangent of the Angle of Inclination; and this Double is Cc representing the Intensity of the Weight. Q. E. D.

If the two Powers are us'd, P will draw with the same Obliquity as P because the Angle $aCc = cCb$; therefore $P \dashv P : p \dashv p :: DC \dashv Dc : CE \dashv Ec$. Q. E. D.

The first Analogy may also be demonstrated by the Doctrine of the Leaver; for if we consider that bm is the Line of Direction of the Power drawing the Leaver mn obliquely, *or* perpendicular to bC (that is bm produc'd) will be its acting Distance*; and the Intensity of the Power acting at right Angles : will be to its Intensity acting obliquely :: as nr : to nm ; but because of the right Angle nrn and the common Angle nmr , the Triangle nmr is similar to oCm and cCb ; and therefore nr will be the Tangent and nm the Secant of the Angle of Inclination. Q. E. D.

* L. 3. Ann.
5. p. 142.

This may be experimentally tried by taking the Weight W together with its Pulley equal to 6 Pounds, and the Weights P and P equal to 5 Pounds each; for then if the Pullies A and B be fix'd at the Distance

of 8 Inches from each other at their Circumference, as at a and b , the Annotat. three Weights will not rest till the Line Cc be just 3 Inches long, in which *Left. III.* Case the Triangles aCc , cCb , EDC , DEc , moC , and mnr will have  their three sides in the Proportion of 4, 3, and 5. But the best Method *Pl. 15. F. 17.* of trying all Cases of this Kind, is to make use of the Machine contriv'd by Dr. s'Gravesande for that Purpose. (See his Introduction to Sir Isaac Newton's Philosophy, Part I. No. 205.) On the Plank or horizontal Board $H\uparrow$ are fixed two Standards S, S , which have, each on its upper Part, a *Pl. 15. F. 18.* Sextant with several Lines drawn from a Center taken on the upper Part of a Pulley, along which Lines going over the Pulleys may be stretch'd. In the middle of the Lines are written the Numbers which express the Secants of the Angles which those Lines make with the Horizon, and at the Ends of the Lines are written the Numbers expressing the Tangents of those Angles. Now in making Experiments it will appear in every Case where there is an *Æquilibrium*, that the Weights Q and Q are as the Numbers in the middle of the Lines along which the Threads are stretch'd; and the Weight P as the Sum of the Numbers at the Ends of those two Lines.

II. [48. — *A less Power will serve &c. — unless it pushes the Body against the Plane, &c. or draws it away from the Plane, &c.*] We have in the 7th Annotation consider'd all that relates to a Body moving on an inclin'd Plane, and therefore refer to that; but it will not be improper here to take notice of the Difference between high Wheels and low ones, as they roll over uneven Grounds or Rubs: Because tho' this Motion cannot be consider'd in every Respect like the rolling of Bodies on inclin'd Planes; yet there are a great many Things alike in both Cases.

Let the Line ab^* represent the horizontal Plane or Way on which a **Pl. 15. F. 19.* Wheel, represented by the Circle $ICbg$, is to roll from a towards b . m, n, o , represent three immoveable Rubs, whose Eminences or Tops reach as high as the Points dgb ; while the Power draws the Wheel in the Line of Direction cG . To know what the Intensity of the Power must be in proportion to the Intensity of the Weight (that is, in proportion to the Weight of the Wheel) we may suppose a bended Lever in the said Wheel, and are to consider its Effect in the Operation, which being well examin'd will give us this general Rule for all Cases of a Wheel going over a Rub on an horizontal Plane, the Line of Direction drawing along the Center of the Wheel being also suppos'd horizontal.

When the Circumference of a Wheel moving vertically on an horizontal Plane touches the Top of a Rub,

The Weight :

Is to the Power that can draw the Wheel over the Rub ::

As the Sine of the Angle, which a Line drawn from the Center of the Wheel to the Top of the Rub, makes with the horizontal Line :

To its Cosine.

Annotat.
Lect. III.



PREPARATION.

Pl. 15. F. 19. Through d the Top of the Rub md , and through g the Top of the Rub ng draw the horizontal Lines edt , rgs ; draw the Radii cd , cg and cb , the last of which is parallel to the Horizon and must be produc'd as far as p ; with the Radius cd and Center d draw the Arc ck ; then from the Points d and g draw dfk and gi perpendicular to cb , and round k for a Center draw the Circle C^*Dd equal to Clq , which will represent the Wheel rais'd up upon the Top of the Obstacle or Rub md .

DEMONSTRATION.

In respect to the Rub dm , cde is a bended Leaver whose *Fulcrum* is d and *Brachia* cd , de ; but as the Power draws obliquely to the *Brachium* dc in the Line pc , the *Brachium* dc must be reduced to fd perpendicular to the Line of Direction (being the acting Distance of the Power) but de preserves its whole Length, because cq the Line of Direction of the Weight goes thro' its Extremity e . Therefore here the Weight and the Power will be to one another reciprocally as the *Brachia* fd and de ; but fd is the Sine of the Angle fdc ($= cde$) which the Line cd makes with the horizontal Line de or its Parallel cf , and de its Cosine. Q. E. D.

Scholium.

If the Rub had been ng , twice as high, the Difficulty of drawing the Wheel over it would have been more than twice as great; because in considering the bended Leaver cgr acting in that Case, it must have been reduc'd to another bended Leaver igr , in which the Power : is to the Weight :: as rg : to gi , where the Disproportion of the acting Distance is more than doubly encreas'd to the Disadvantage of the Power.

COROLLARY I.

Hence follows, that the Difficulty of a Wheel to go over a Rub encreaseth in a greater Proportion than the Height of the Rub; the Rubs of different Heights compar'd together being always as the vers'd Sines of the Complement of the Angle of Inclination, when the Power : is to the Weight :: as the Sine Complement : to the Sine of the Angle of Inclination, which last *Ratio* encreases faster than the versed Sines.

COROL-

C O R O L L A R Y II.

Hence follows also, that a Wheel cannot by any Power, how great soever, be drawn over a Rub whose Top is as high as the Axis (as for example the Rub *ob*, because in that Case the Sine Complement is become the whole Sine *cb* and the right Sine is vanish'd; or, what is evident by the Scheme, the Power draws against the *Fulcrum* where it can have no Effect at all; let its Intensity be ever so great; unless the Direction of the Power be made to alter, and it should draw upwards: Therefore in Practice, especially where Carriages are to go upon rough Ways, it is usual to make the Horses or Oxen draw a little upwards from the Center of the Fore-Wheels.

C O R O L L A R Y III.

Hence likewise may be seen the Reason why high Wheels go over Rubs more advantageously than low ones, and that in proportion as they are higher; because the Lengths of versed Sines being (*cæteris paribus*) as the Diameters of the Circles to which they belong: That Rub, whose Height was the versed Sine of an Arc of a certain Number of Degrees in a lesser Circle, will be the Sine of an Arc of fewer Degrees in a greater, in proportion as it is greater; therefore the Sine Complement or horizontal *Brachium* of the bended Leaver bearing the Weight, will be less, and the Sine of Inclination or perpendicular *Brachium* of the Leaver, to which the Power is applied, will be greater. Besides, the high Wheel will not only go over Rubs impossible to the low Wheel, but several other Rubs still higher, provided their Height be not equal to the Semidiameter of the great Wheel. As for example in *Fig. 20**, the Intensity of the Power ^{*Pl. 15. F. 20.} *P* drawing the great Wheel *CD* over the Rub *D* along the horizontal Line *ab* is but half the Intensity of the Power *p* drawing the little Wheel *cg* over the same Rub at *g*: Because it is not the bended Leaver *igr*, which is transferr'd from the little Wheel to the great one at *FDE*, but the Leaver *fde*, whereby the little Wheel is drawn over a Rub only of half the Height. So in *Fig. 19*.†, the great Wheel, Part of whose Cir- ^{†Pl. 15. F. 19.} cumference is represented by the Semicircle *LHg*, goes over the Rub *MD* with the same Ease, that the little one goes over *md*; over the Rub *NG* (impossible to the little Wheel) with the same Ease that the little Wheel goes over *ng*, and has no Rub impossible to it (supposing the Intensity of the Power sufficiently great) till it becomes of the Height *OH* equal to its Semidiameter.

C O R O L L A R Y IV.

Lastly, we may observe from what has been said upon this Subject, that the greatest Difficulty to bring a Wheel over a Rub is in the first Effort, and

Annotat. and that the Action of the Power becomes continually easier as the Wheel
 Lect. III. rises, whether the Wheel was at Rest or in Motion just as it began to
 Pl. 15. F. 19. press upon the Rub; for the horizontal *Brachium* of the Leaver as *ed*
 (supposing *md* the Rub) rising continually round the *Fulcrum d*, when the
 Wheel is rising from *Cbgql* to *x Ddr*, diminishes its acting Distance as
 the Line of Direction *cq* of the Weight comes from *cq* to *kd*, whilst the
Brachium df by which the Power acts continues the same, as the Line of
 Direction of the Power goes from the Situation *cp* into the Situation *kx*;
 for the Distance of the Horses or Oxen, &c. is so great in proportion
 to the Height of the Rub, that we need not look upon the Point *x* as
 rais'd at all above the Horizon; nay, if the Horses draw a little upwards,
 every Advance of the Line of Direction of the Power towards the Point *x*
 will be a more advantageous Situation of it.

From this last Consideration it appears, that when a Wheel, as *C1q*,
 goes over a Rub as *md* or *ng*, we are not to consider it as if it roll'd up
 an inclin'd Plane as *qd* or *qg*, where the Power drawing horizontally (as in
 the Case of the Wedge) acts uniformly and that in the Ratio of the Base
 to the Height; the Power here being requir'd to exert much more Force
 at first, and less afterwards.

12. [50. — How much the Power must be increas'd in proportion to the
 Angle, which its Line of Direction makes with the Plane, will be shewn in
 the Notes.] All this has been fully consider'd in the seventh Annotation
 to this Lecture.

13. [62. — This sort of Thread is not us'd in Wood; but in Iron and o-
 ther Metals it is of good Service, being commonly more durable, and raising
 the Weight with more ease than the Sharp Thread, as will be more fully
 shewn in the Notes.] The Square Thread is seldom or never made use of
 in Wood, because then the prominent Parts of the Screw, such as *P, N,*
 Pl. 11. F. 11. *L, H, Q, O, M, K, I* (Plate 11. Fig. 11.) would have no more Strength
 than the lateral Cohesion of the Fibres of the Wood, for a Length very
 little greater than the Thickness of the Thread all the Way; so that in
 very great Strains the Arbor *abcd* might be stript of its Thread all the
 Way, and so let go what it was intended to hold. But to prevent this
 Inconveniency the hollow'd Part of the Screw is made sharp close to the
 imaginary Arbor or included Cylinder, which thickens the Thread of the
 Screw next the Arbor; but then of consequence it must come to a sharp
 Pl. 11. F. 14. Edge on the outside. This may be seen in the 14th Figure, where, instead
 of the hollow *BACD*, you have *bad*.

Tho' by this means the sharp-threaded Screw is strongest in Wood, it
 is weakest in Metals, where the Thread is generally fine, for if the Male
 and Female Screw do not exactly fit, but have a little too much Play, the
 Sharp Thread of the one will gull (that is, cut and wear away) the Sharp
 Thread of the other. Whereas in a Square Thread, tho' there should be
 some Play, there is no unusual wearing out, a Flat bearing upon a Flat.
 Besides,

Besides, on the Sharp Thread the Weight endeavours to descend (and therefore resists) with more Force than on the Square Thread; because, besides the Endeavour of the Weight to slide back on that inclin'd Plane which makes the Ascent of the Screw (which has been already explain'd in describing the inclin'd Plane and Wedge) it has also an Endeavour to descend along another Plane (as *ad Fig. 14.* *) which makes the Sharpness of the Thread, and consequently the Power must be encreas'd on that Score. Annotat. Lect. III.

To estimate what is the Force which the Sharp Thread has to bear the Weight on account of its Declivity, let us examine the Screw *CD F E M* * supposing a Female Screw to be rais'd with all the Weight, which consequently must press upon the Thread of the Male Screw here represented at *CIKPGH*, &c. which we will all reduce to the Pressure on the Point *P*, *WABP* being suppos'd the Weight pressing, and *WP* its Line of Direction. Now, if we resolve the Force *WP* representing the Pressure of the Weight downwards into the two Forces *WA* and *WB*, of which the first represents the Force exerted against the Declivity *AD*, and the last that whereby the Weight endeavours to go along *AD* without pressing against it; it will appear (because Action and Reaction are equal, that the inclin'd Sharp Thread only lifts the Weight with the Force represented by *AW*, whilst the Force left, *viz. WB*, carries down the Weight in the Direction *AD*, thereby drawing the opposite Side of the Thread of the Female Screw to press the harder against the Thread of the Male Screw at *C, M* and *E*, &c. So that if the Weight does not happen to rise perpendicularly (which it does not in many Cases) there will be a great Encrease of Friction on the upper Side of the Screw. * Pl. 11. F. 14.

The Friction of the Screw being owing to many Causes, we shall consider it in another Place.

14. [70. — Pretenders to Perpetual Motions, and those who promise greater Effects by Machinery than is conformable to the reciprocal Proportion between the Intensities of the Powers and Weights and their Velocities.] About the Years 1720 and 1721, the late Mr. John Rowley, Mathematical Instrument-maker, talk'd so much of the Wheel which he had seen at *Hesse-Cassel* (which he believ'd to be a perpetual Motion, as well as a great many Persons in that Country) that besides the common Herd of *Perpetual Motion Men*, which every Age affords, some very ingenious Men made an Attempt that Way, and were countenanc'd in it by some great Mathematicians, who, when the Scheme was laid before them, declar'd they knew no Reason why it should not do. But as I always declar'd against all Projects tending that way, I was desir'd at that time to publish my Reasons why the Thing seem'd impossible or impracticable; which I did in the *Philosophical Transactions* * in such a manner as might dissuade People at first from any such Attempts, in which so much Time and Money have been lost. I have here printed the whole Account again; but desire * No. 369.

Annotat. desire my Reader first to look over what I have said in these Notes, Page
Lect. III. 146 and 147.

The Wheel at Hesse-Cassel made by Monsieur Orfireus, and by him call'd a Perpetual Motion, has of late been so much talk'd of on account of its wonderful Phænomena, that a great many People have believ'd it to be actually a self-moving Engine; and accordingly have attempted to imitate it as such. Now, as a great deal of Time and Money is spent in those Endeavours, I was willing (for the sake of those that try Experiments with that View) to shew that the Principle which most of them go upon is false, and can by no means produce a perpetual Motion.

They take it for granted, that if a Weight descending in a Wheel at a determinate Distance from the Center, does in its Ascent approach nearer to it; such a Weight in its Descent will always preponderate, and cause a Weight equal to it to rise, provided that Weight comes nearer the Center in its Rise; and accordingly, as itself rises, will be overbalanc'd by another Weight equal to it; and therefore they endeavour by various Contrivances to produce that Effect, as if the Consequence of it would be a perpetual Motion.

But I shall shew, that they mistake one particular Case of a general Theorem, or rather a Corollary of it for the Theorem itself. The Theorem is as follows :

If one Weight in its Descent does (by means of any Contrivance) cause another Weight to ascend with a less Momentum or Quantity of Motion than itself, it will preponderate and raise the other Weight.

Cor. I. Therefore if the Weights be equal, the descending Weight must have more Velocity than the ascending Weight, because the *Momentum* is made up of the Weight multiplied into the Quantity of Matter.


C O R O L L A R Y II.

Therefore if a Leaver, or a Balance, has equal Weights fasten'd or hanging at its Ends, and the *Brachia* be ever so little unequal, that Weight will preponderate, which is farthest from the Center.

Scholium.

This second Corollary causes the Mistake; because those, who think the Velocity of the Weight is the Line it describes, expect that Weight shall be overpois'd which describes the shortest Line, and therefore contrive Machines to cause the ascending Weight to describe a shorter Line than the descending Weight. As for example, in the Circle ADB * Pl. 16. F. 2. (Fig. 2.)* the Weights A and B being suppos'd equal, they imagine, that if by any Contrivance whatever, whilst the Weight A describes the Arc Aa , the Weight B is carried in any Arc, as Bb , so as to come nearer the Center in its rising than if it went up the Arc BD ; the said Weight shall

shall be overpois'd, and consequently by a number of such Weights, a perpetual Motion will be produced.

Annotat.
Lect. III.


This is attempted by several Contrivances, which all depend upon this false Principle ; but I shall only mention one, which is represented by *Figure 4*, where a Wheel having two parallel Circumferences, has the Space between them divided into Cells, which being curv'd, will (when the Wheel goes round) cause Weights plac'd loose in the said Cells, to descend on the Side AAA, at the outer Circumference of the Wheel ; and on the Side D to ascend in the Line Bbbb, which comes nearer the Center, and touches the inner Circumference of the Wheel. In a Machine of this kind, the Weights will indeed move in such a manner, if the Wheel be turn'd round, but will never be the Cause of the Wheel's going round. Such a Machine is mention'd by the Marquis of Worcester, in his *Century of Inventions* in the following Words, N^o. 56.

“ To provide and make that all the Weights of the descending Side
“ of a Wheel shall be perpetually farther from the Center than those of
“ the mounting Side, and yet equal in Number, and Hest to the one Side
“ as the other. A most incredible Thing, if not seen ; but try'd before
“ the late King (of blessed Memory) in the Tower by my Directions,
“ two Extraordinary Ambassadors accompanying his Majesty, and the Duke
“ of Richmond, and Duke Hamilton, with most of the Court attending him.
“ The Wheel was fourteen Foot over, and had forty Weights of fifty
“ Pounds apiece. Sir William Balfore, then Lieutenant of the Tower,
“ can justify it, with several others. They all saw, that no sooner these
“ great Weights pass'd the Diameter Line of the lower Side, but they
“ hung a foot farther from the Center ; nor no sooner pass'd the Diameter
“ Line of the upper Side, but they hung a Foot nearer. Be pleas'd to
“ judge the Consequence.

Now the Consequence of this and such like Machines, is nothing like a perpetual Motion ; and the Fallacy is this. The Velocity of any Weight is not the Line which it describes in general, but the Height it rises up to, or falls from, with respect to its Distance from the Center of the Earth. So that when the Weight (*Fig. 2. **) describes the Arc Aa, its ^{* Pl. 16. F. 2.} Velocity is the Line AC, which shews the perpendicular Descent (or measures how much it is come nearer to the Center of the Earth) and likewise the Line BC denotes the Velocity of the Weight B, or the Height that it rises to, when it ascends in any of the Arcs Bb, instead of the Arc BD : So that in this Case, whether the Weight B in its Ascent be brought nearer the Center or not, it loses no Velocity, which it ought to do, in order to be rais'd up by the Weight A. Nay, the Weight in rising nearer the Center of a Wheel, may not only not lose of its Velocity, but be made to gain Velocity in proportion to the Velocity of its counterpoising Weights, that descend in the Circumference of the opposite Side of the Wheel ; for if we consider two Radii of the Wheel, one of which is horizontal, and the other (fasten'd to and moving with it) inclin'd under the Horizon in an Angle of 60 Degrees (*Fig. 3. †*) and by the Descent of ^{† Pl. 16. F. 3.} the

Annotat. the End B of the Radius BC, the Radius CD by its Motion causes the
 Left. III. Weight at D, to rise up the Line pP , which is in a Plane that stops the
 ~~~~~ said Weight from rising in the Curve DA, that Weight will gain Velocity,  
 and in the Beginning of its Rise it will have twice the Velocity of  
 the Weight at B; and consequently, instead of being rais'd, will overpoise,  
 if it be equal to the last mention'd Weight. And this Velocity will be  
 so much the greater, in Proportion as the Angle ACD is greater, or as  
 the Plane  $Pp$  (along which the Weight D must rise) is nearer to the  
 Pl. 10. F. 2. Center. Indeed if the Weight at B (Fig. 2.) could by any means be lifted  
 up to  $\beta$ , and move in the Arc  $\beta b$ , the End would be answer'd; be-  
 cause then the Velocity would be diminish'd, and become  $\beta C$ .

### EXPERIMENT. Fig. 3.

Take the Leaver BCD, whose *Brachia* are equal in Length, bent in  
 Pl. 16 F. 3. an Angle of  $120^\circ$  at C and moveable about that Point as its Center: In  
 this Case, a Weight of two Pounds hanging at the End B of the hori-  
 zontal Part of the Leaver, will keep in *Æquilibrio* a Weight of four Pounds  
 hanging at the End D. But if a Weight of one Pound be laid upon the  
 End D of the Leaver, so that in the Motion of D along the Arc  $pA$ ,  
 this Weight is made to rise up against the Plane  $Pp$  (which divides in  
 half the Line AC equal to CB) as having twice the Velocity of it, when  
 the Leaver begins to move. This will be evident if you let the Weight  
 4 hang at D, whilst the Weight 1 lies above it: For if then you move  
 the Leaver, the Weight 1 will rise four times as fast as the Weight 4.

Pl. 12. F. 3. 15. [80. — The Weight and Gibbet would run back and rest over W,  
 &c.] In the third Figure of Plate 12 CGrgs represents the Top of the  
 Gibbet with its Pulley at  $r$  its Extremity, and its Center (or the Top of  
 its Axis) at C. It is to be observ'd here, that the horizontal Part of the  
 Rope Cr is in the same Plane with the middle Line of the Gibbet, or  
 that the Rope Cr is parallel to the Line Cr (under it) in the Gibbet.  
 Now if it was possible for the Rope to continue parallel to the said Line  
 Cr in the middle of the Top of the Gibbet, whilst it is mov'd to the  
 left successively in the Situations C6, C7, C8, or to the right in the Si-  
 tuations C1, C2, C3, C4, C5, the Guider, or Person who pulls and  
 directs the Guide-rope fasten'd at the End near g, would bring the Weight  
 to any Place on the Wharf on either side the Crane without any more  
 Labour than what overcomes the Friction of the Axis of the Gibbet.  
 But when the Pulleys P and Q are plac'd at  $p$  and  $q$ , the middle line of  
 the Gibbet advances towards P and Q faster than the Rope does, which  
 being oblig'd to fold about the Pulleys makes an Angle with the Line  
 abovemention'd, as for example, the angle  $p8n$  on the left Side and  $q4t$   
 on the right, the angle being so much greater as the Gibbet is more drawn  
 back on the Wharf towards the Crane. The Consequence of this is that  
 the

the Weight (in this Motion of the Gibbet) must be rais'd in proportion Annotat. as the Rope is lengthen'd by folding about the Pulley, or in the propor- Lect. III. tion of  $Cp8$  to  $C8$  on the left, and  $Cq4$  to  $C4$  on the right. Now if the Excess of the lengthen'd Rope above the Length of the middle Line of the Top of the Gibbet be (for example) one tenth of the whole, then a Person holding the Gibbet by the Guide-rope, in a Position which makes that Difference, must sustain a tenth part of the whole Weight, viz. 224 lb in a Ton, too great a Weight for a Man thus employ'd to move with the Gibbet. Now as this Force gradually encreases by drawing round the Gibbet, it deceives the Man who pulls the Guide-rope, and as by a sudden Jerk he has brought on the Weight, he is often forc'd to let it go when it comes beyond his Strength, which sometimes proves of dangerous Consequence, as well to the Persons loading or unloading as to the Goods cran'd up.

To prevent these Inconveniencies, Workmen have plac'd the Pulleys in a different Situation, viz. in the Situation P, Q; so that in turning the Gibbet to the right, no Part of the Weight will be felt by the Guider till the Gibbet comes beyond  $C4$ ; and in moving to the left, till it comes beyond  $Cc$ . But then another Inconveniency arises from this Construction, which is the Reverse of the other Thing consider'd; namely, that the Weight descends, and consequently brings on the Gibbet with a Swing, which (if unexpected) may likewise do Mischief. As in large Weights these Inconveniencies are most sensible, Mr. Padmore of Bristol making a Crane for Mr. Allen (Postmaster of Bath) to raise Stone out of a Quarry, contriv'd an Application of the *Axis in Peritrochio*, which takes off this Danger and Inconveniency. In the first Figure  $x$  is a Wheel with Arms, Pl. 12. F. 1. whose Axis  $xu$  has on it a Pinion at  $u$ , which takes the Teeth of an horizontal contrate Wheel fasten'd to the Axis of the Gibbet; whereby one Man standing out of Harm's Way close to the Wheel has such Advantage of Power by means of the long Arms of the Wheel, as to move the Axis of the Gibbet with great Ease, notwithstanding the Inequalities above mention'd, and also to hold the Gibbet in any Position without difficulty.

The same ingenious Workman has made another considerable Improvement in another Crane fix'd by the River-side, whereby Mr. Allen lets down his Stone into such Vessels as come to fetch it away.

The Crane itself is not of an uncommon sort, but a rat-tail'd Crane with a double Axis in *Peritrochio* and two Handles, whereby four Men may raise very great Weights; and then turning the whole Crane about upon its upright Shaft, can fix it in any Position and let down the Weights speedily into the Boats or Barges which come near the Wharf to receive them. See the 5th Figure of Plate 16, where you may observe it to differ from that represented in Plate 12. Fig. 4. because the long Neck of Pl. 12. F. 4. the Crane is here of one Piece, and the Power differently applied. But this Construction is not new. Neither is it a new Invention to let down Goods after they have been rais'd by a Crane, by pressing an Arch of a

Annotat. Circle strongly upon a Wheel fix'd to the principal Axis, in order to retard and regulate the Descent by a Friction increas'd or diminish'd at pleasure, as is done in stopping Windmills. The Catch also, that hinders a Crane or Capstane from going back, is of common Use; but I don't know that any one has applied them both together in the same Crane, so as to depend upon one another, before Mr. *Padmore* did it, though many have done it since. Therefore I shall give a particular Description of this Contrivance, whose chief Intent is to prevent the great Mischiefs which often happen by the Carelessness of the Men employ'd to raise and let down heavy Burthens by the Use of the Crane. The sixth Figure represents an upright Section of so much of the Crane as the Contrivance abovemention'd is apply'd to.

PL. 16. F. 6.

AB is the great Wheel, whose large Axis A moving on two Iron Center Pins such as *a*, receives the Rope, or lets it run down according as it is turn'd, by means of the Handles fasten'd at C to the lesser Wheel or Pinion C, or as it is suffer'd to turn the other Way by the Gravity of the descending Weight when all Obstacles are remov'd. Upon the Axis of the Pinion is the Ratchet Wheel Dd, whose Teeth successively receive the Iron Catch fF (moveable on a Pin at F on the Iron Standard G, and to be rais'd up occasionally by the upright Iron Hh) to hinder the Weight from going back when the Handles are loos'd. Upon the same Axis behind the Wheel Dd is a wooden Wheel Ep, over which hangs the Half-ring of Iron OPo with a Groove or Hollow made in it to fit the Circumference of the said Wheel, so as to retard, or stop, or any way regulate the Motion of the Wheel (and consequently of the Axis and Pinion C, and the great Wheel and Axis AB which has the Rope VA) according as it is more or less strongly press'd down to make a Friction on the Wood, as it moves after the Catch is rais'd out of the Teeth of the Ratchet. The horizontal Leaver KL governs all these Motions in the following manner, viz. When the String QqK, fasten'd to the said Leaver at K, is pull'd, the Leaver moving on its Center M, does, by an horizontal Pin fix'd at right Angles to its Side at I, raise the Piece Hh, and consequently release the Ratchet by raising the Catch at f out of the Teeth: Then the Weight descends swiftly, moving the Wheel and Pinion round by its Force; but to prevent the too swift Descent, the Leaver is pull'd up a little more strongly by the Guider who holds the String QqK, which brings down the contrary End of the Leaver L, and consequently the Iron N, so low as to make the semicircular Ring OPo press hard upon the Wheel Ep, which it did not do when the Catch was rais'd but just out of the Ratchet. N.B. A strong Pull stops the whole Motion, and a more gentle one regulates the Descent. And if the Guider should be careless and let go the String; then immediately the Spring Ss, whose End s had been depress'd by the End L of the Leaver, will raise it up again (by its lateral Pin X) and restoring the whole Leaver to its first horizontal Position, the other lateral Pin I in the long Arm MK of the Leaver, will through the Notch H, press upon I the lower End of the upright Piece

Piece *Hh*, and so bring down the Catch *Ff* into the Ratchet Wheel at Annotat. *f*, the curved Piece *Opp* at the same time flying up and no longer pressing the wooden Wheel *Ep*. Lect. III.

Thus will Mischief never be the Consequence of Carelessness, because of the Catch; nor will the Weight go down by jerks, which would have been the Consequence of the Catch us'd without the Half-Ring, because the Catch is lifted quite out of the way when the Half-Ring is brought down and applied by pulling the String at *Q*. *N. B.* *T, t 1, t 2* is part of the upright Section of the Timber of the Frame. Pl. 16. F. 6.


To make this the plainer, let us examine the 7th Figure of Plate 16; Pl. 16. F. 7. where we have an horizontal Section of the Parts above mention'd. *TtT* is Part of the Timber of the Frame. *BB* is the great Wheel, whose Axel that holds the Rope is mark'd *AAA*, and its Iron Axis goes thro Bell-metal Boxes at *aa*. *CC* is the Small Wheel or Pinion, whose Axis is *cc*. *DD* is the Section of the Ratchet Wheel made of Iron. Between the prick'd Lines *Ap* and *Ei* is suppos'd the wooden Wheel upon the Axis of the Pinion (not drawn here to avoid Confusion, no more than the semicircular pressing Piece mark'd *OPo* in the last Figure) the Basis of whose upright fix'd Supporter is represented by *R*, and the End of the Piece which brings it down upon Occasion is shewn at *N*. *KL* is the horizontal Leaver, whose Center is at *M*, moving vertically by a Pull of the String fasten'd to it at *K*. *Ii* is the first lateral Pin of the Leaver, which at *I* goes through the bottom of the Piece *H*, Raifer of the Catch *Ff*, already describ'd with its Supporter *G*, on whose Top the Catch moves by a Center Pin. *ApA* is the second lateral Pin of the Leaver, whose Office is to press upon the End *s* of the crooked Spring *Ss* fasten'd to the bottom of the Frame at the farther End *S*. So that when the End *K* of the Leaver is pull'd up, the End *L* which is depress'd, must be lifted up again into its Place by the Force of the Spring restoring it self.

*N. B.* The crooked Figure of the Spring, and the Manner of its lifting the Pin is best shewn in the last Figure. Pl. 16. F. 6.

There is a fine Contrivance of an inclin'd Waggon-Way made of Timber to bring down the Stone from Mr. Allen's Quarry to the River's Side, the Distance of near a Mile, in Waggons which come down the Declivity on the artificial Way by their own Gravity, as the Coal-carriages do near Newcastle. But this Waggon-way differs quite from those in the North, every Part being very much alter'd for the better, and the Carriages themselves contriv'd to carry much more Weight without Danger. The Description of this Contrivance, wherein Mr. Allen and his Workmen have shewn great Skill and Ingenuity, I must now omit, and give it in another Part of this Book.

## LECTURE IV.

### *Concerning the Friction in Mechanical Engines.*

Lect. IV.  **W**HAT I have said hitherto in the Three first Lectures and their Notes, is sufficient for explaining the Principles of Mechanicks (strictly so call'd) enough to set People to work, who have a Genius for Practical Arts; but as they may be guilty of considerable Errors without making proper Allowances for Friction, and knowing how to find out (nearly) what it is in Engines already made, and such as they intend to make; I thought proper in this Place to give a Lecture on what I have been able to discover of the Nature of Friction, by reading all the Accounts I could meet with on that Subject, by repeating several Experiments already made the better to consider their Circumstances and whether the Accounts of them were true; and likewise by making a great many new Experiments my self.

*Tho' there are so many Circumstances in the Friction of Bodies, that the same Experiment does not always succeed with the same Bodies, so that a Mathematical Theory cannot be easily settled; yet we may deduce a Theory sufficient to direct us in our Practice from a great Number of Experiments, always taking a Medium between Extremes.*

FIRST then, it is observ'd that Wood, Iron, Brass, Copper and Lead, when greas'd or oil'd (as is done in Engines) have nearly the same Friction, and therefore the same Rule will serve for all those Substances. For tho' one may at first imagine that Metals must needs slip over one another more easily, because they may be made smoother and will take a better Polish; yet it is found by Experience that the flat Surfaces of Metals or other Bodies may be so far polish'd as to encrease Friction; and this is a  
mecha-

mechanical Paradox ; but the Reason will appear when we con-  
sider that the Attraction of Cohesion becomes sensible as we bring  
the Surfaces of Bodies nearer and nearer to Contact. This is ve-  
ry evident in drawing Glass Plates over one another, which re-  
quires more Force than if they were of Wood, and the same is  
true in Metals, for tho' the Pressure of the Air is sometimes a  
little concern'd in this Phænomenon, yet the Attraction of Cohesion  
does so far exceed it, that we need not take notice of the Effect  
of that Pressure in the Action of the Parts of an Engine on one  
another. It is true indeed that a couple of Marble Plates stick to  
each other by the Pressure of the external Air, when being first  
oil'd, to exclude the Air from between them they have been slipp'd  
upon one another ; and that when suspended in the Receiver of an  
Air-Pump they fall asunder as soon as the Air is drawn out of the  
Receiver. But if they be made very smooth, they will cohere  
strongly even after the Pressure of the Air is wholly taken off by  
the Air-Pump. I have applied together the flat Surfaces of two  
small Crystal Buttons, without wetting or oiling, which have  
coher'd so strongly as to hold 19 Ounces *Troy* before they were  
separated; when their Contact was but a Circle of one 12th of an  
Inch in Diameter ; in which Case the Pressure of Air could not be  
greater than the Weight of an Ounce ; because a Column of Air,  
whose circular Base is one 12th of an Inch in Diameter weighs no  
more. In Metals the same is evident ; but more especially in Lead ;  
for two Balls of Lead of about one or two Pound Weight each,  
if par'd clean with a Knife and applied close together, so as to  
touch in a Surface of about one 12th of an Inch in Diameter, will  
stick together so as not to be separated by a less Weight than of  
40 or 50 Pounds, tho' the Pressure of the Air in that Case could  
not amount to one 4th of a Pound. *See Phil. Transact. N. 389.*  
But to return to our Experiments of Friction.

EXPERIMENT I. *Pl. 13. Fig. 18.*

Pl. 13. F. 18.

*Cb* is a smooth Piece of Wood one Inch thick, four Inches  
wide and six long, with a little Hook in its Forepart *b*, and weigh-  
ing six Ounces. If it be drawn along an horizontal Plane repre-  
sented by the Line *AB* over the Pulley *P* by means of the String  
*HW*, the Weight *W* equal to two Ounces and as much more as  
will overcome the Friction of the Pulley, will draw it along whe-  
ther

Lect. IV. ther it slides on its greatest Surface or Flat as at *c*, or on its Edge as at *K*, tho' in the last Case there is only a fourth part of touching Surface. From hence it appears that *the Friction is equal to about one 3d of the Weight,\* and arises from the Weight that presses the Parts together, and not from the Number of Parts that touch*; for whether the Parts of the Surface be consider'd as Springs to be bent, or as small Eminences over which the Body drawn must be lifted, it is plain that the Difficulty will be the same, whether the Surface be greater or less, provided it be press'd in a reciprocal Proportion of its Bigness; for tho' the Base of *c* be four times broader than the Base of *K*, every Inch of it in Breadth is press'd but by one Inch of Wood in Thickness; whereas the Surface on which it rests in the Position *K*, which is four times narrower, is press'd by a Thickness of Wood of four Inches. The Friction therefore, or Impediment to the drawing the Body in the Line *AB*, arises not from the Number of Parts which touch, but from the Pressure on the touching Surface, as appears yet more plainly from the following

Pl. 13. F. 18.

## EXPERIMENT II. Pl. 13. Fig. 18.

LET the Pulley  $\pi$  (over which the Weight  $w (= W)$  drew along *K*, when it was rais'd up to the Level of *m* the Hook of *K* above *AB*) be brought down to the same Level as *P*, then  $w = W$  will not draw *K* along without the Help of the little additional Weight *q*; because the Traction made in the oblique Direction *mp* makes *K* press more against the horizontal Plane than when the Draught is in the Direction *m $\pi$* ; but if the Hook be fix'd at *n* so as to make the Line of Direction parallel to the Plane *AB*, then will the Weight  $w$ , without the Help of *q*, draw *K* along as *W* did *c*.

Pl. 13. F. 19.

## EXPERIMENT III. Pl. 13. Fig. 19.

UPON the Table or horizontal Plane *TMNO*, let a circular Plane or Plate of about two Foot Diameter move, bearing on a Pin round the Center *C* so as not to touch the first Plane; then let a Brass or Leaden Plate, such as *A*, *B* or *D*, be so laid betwixt the circular Plane and the Table, as to press upon the Table with all

all its Weight, and at the same time (by means of a Pin rising Lect. IV.  
from its middle) be carried round along with the circular Plate under the Point A, B or D. Then if the Weight of the Plate A be equal to 30 lb, a Force equal to 10 lb will draw the Plate round pulling in the Direction AT, when the said Plate is plac'd at A or D; but if the Power draws in the Line Bt or Et Tangent to the little Circle BE, it will require 20 lb to draw A round: Lastly, if the Power draws from A towards T in the Tangent of the great Circle AtD, whilst the Plate is fix'd at B, then a Force equal only to 5 lb will move round the Plate.

THIS shews that the Friction is increas'd when the rubbing Parts move faster, and decreas'd when they move slower than the Power. Therefore we may say that *the Resistances from Friction are to one another in a Ratio compounded of the Pressures of the rubbing Parts, and the Times or Velocities of their Motions.*

WHEN a smooth Body is laid on an inclin'd Plane, according to the Laws of Mechanicks it ought not to rest upon it, but must slide down. But the Friction being equal to one 3d of the Weight, as we have already shewn, the Body K\* laid on the inclin'd Plane\* Pl. 13. F. 29. AB will not slide down till the Perpendicular AC be the third part of the Base BC; because then the Body's Tendency downwards in a Line parallel to AB is equal to one 3d of the Weight of the Body.

HENCE it appears how necessary it is to consider the Friction in mechanical Performances; as for example, if a Beam AB† ly-† Pl. 17. F. 1. ing on the rising Ground Dc is to be drawn by a Power at P in the Direction AP parallel to Dc; when Aa the Rise of the Plane above the horizontal Line Hb is one 30th of ac, a Power equal to one 30th of the Weight would be able to draw it along according to the Laws of Mechanicks exclusive of Friction; and therefore 100 lb would draw along a Beam of 3000 lb along the Plane Ac: But as the Friction is equal to one 3d of the Weight, we must have a Power equal to 100 lb + 1000 lb before we can draw the Beam along, if the Plane was as smooth as a Marble Floor. But along the Ground, where even the smoothest Earth is rough, a much greater Force will be requir'd. *Monf. Amontons* of the Royal Academy at Paris found that a Force equal to 2250 lb was but sufficient to draw such a Beam up a Hill of the Ascent abovemention'd.

## Lect. IV.

Pl. 17. F. 2.

## EXPERIMENT IV. Pl. 14. Fig. 2, 3 and 4.

LET AB a wooden Cylinder of an Inch Diameter, smooth and well polish'd, turn freely on two Supporters Ss, likewise smooth, well polish'd and greas'd. Then at each End of a filken Thread or String thrown over the said Cylinder, or wound about it two or three times (for that makes no difference if the Parts of the String do not touch one another) suspend a Weight of 9 lb, which Weights will keep each other in *Æquilibrio*. Then adding several Weights to the 9 lb Weight P here made use of as a Power, to find exactly what Addition it must have to raise the Weight W, which is also only a 9 lb Weight, that Weight will not be rais'd up by the turning of the Cylinder till P and all the Weights added to it make all together 18 lb. This is agreeable to what was shewn before by the first Experiment of this Lecture. For as the whole Weight that presses the Cylinder on its Supporters is  $W + P$  or 18 lb, 6 lb must be added to P in order to overcome that Friction; but that additional Weight adds also a Friction of 2 lb the third part of its Pressure, for the overcoming of which Friction 2 lb must be added, whose Friction requires also two 3ds of a Pound, and so on, in a Series of Fractions as  $\frac{2}{9} + \frac{2}{27} + \frac{2}{81}$ , &c. Now as the Weight of the Cylinder itself was not consider'd, the proper Way to allow for its Friction, if it be not very heavy, is to double the Power at once, \* as appears in Fig. 4, where A represents the Section of the Cylinder above mention'd.

IF the Cylinder has Gudgeons or a small Axis, such as that represented in Fig. 2. † it will move as much more easily on its Supporters as the Diameter of the Axis at G and g is less than the Diameter of the Cylinder *ab*; as here the Weight added to *p* need be but  $1\frac{1}{2}$  lb when the Diameter of the Axis is but  $\frac{1}{2}$  of an Inch; because then the rubbing Parts move slower than the Power, in that Proportion, agreeable to what was shewn by the third Experiment of this Lecture. The fourth Figure || represents the Section of the Cylinder at *a*, and of its Axis at *g*.

HENCE we may know what difference there is between the Use of Sledges and Carts for drawing along heavy Goods. If the Ground was perfectly even like a marble Floor it would require  $\frac{1}{3}$  of the Weight of the Sledge, and all that is laid upon it, to draw it along; \* but as such a Plane is not us'd in Practice (and if it could be had, Horses

\* Pl. 17. F. 4

† Pl. 17. F. 3.

|| Pl. 17. F. 4.

\* Ann. 2.

Horses could not pull well on a slippery Pavement) we must al- Lect. IV.  
ways make use of a Power greater than one 3d of the Weight. ~~~~~  
But in a Cart, if the Wheel be six Foot high, and the Hole in  
the Knave for the End of the Axel four Inches, the Friction will  
be diminish'd 18 times; therefore a Power equal to  $\frac{1}{4}$  Part of the  
Weight will be able to draw it along upon an horizontal Plane;  
because the 18th Part of  $\frac{1}{3}$  is  $= \frac{1}{4}$ . Sledges therefore should on-  
ly be us'd where Streets are too narrow for Carts, and where  
Goods in small Parcels are to be laid on and taken off often, for  
the Conveniency of the Men; but the Horses must work the  
harder. This also shews that high Wheels in Carriages are pre-  
ferable to low ones on account of their having less Friction, be-  
sides the mechanical Advantage which we have consider'd in another  
Place; tho' it is not always expedient to make use of them, as for  
example, in Beer-Carts high Wheels would be inconvenient, because  
of the Necessity of often loading and unloading.

To reduce all this to Practice, take the following Rules for cal-  
culating the whole Friction of Engines.

*R U L E I.*

IN examining all the Frictions, begin with that which is near-  
est to the moving Power.

*R U L E II.*

To find the first Friction, consider the Spaces gone through in  
the same time by the Power and by the rubbing Part, and accord-  
ingly take a proportional Part of  $\frac{2}{3}$  of the Power. As for exam-  
ple, in equal Velocities of the Power and rubbing Part, the Fric-  
tion is  $\frac{1}{3}$  of the Pressure which is made not only by the Power,  
but by the Resistance equal to it, and therefore the third part of  
the whole Pressure must be equal to two 3ds of the Power.

*R U L E III.*

THE Power being unknown, must first be found by knowing  
the Velocity of the Weight, and thence deducing that of the Power  
by mechanical Principles: Then, by what has been shewn here,  
finding the Friction, there must be so much added to the Power  
to make the Engine work; always observing that what has been

Lect. IV. added for the Friction first found does still further encrease the Friction, and therefore there must still be allow'd something for the Encrease of Friction caus'd by every new Addition to the Power

N. B. We must remember, as has been said before, that if the Power be an heavy Body, its Velocity is its perpendicular Ascent or Descent ; but if of another kind, the Velocity is its Space gone through.

#### R U L E IV.

THE Friction of the several Parts of an Engine however remov'd from that Part to which the Power is applied may be found by comparing their Velocities with that of the Power, and thence deducing the Friction as has been shewn ; and when all the particular Frictions have been found and added together, their Value added to the moving Force will not enable it to overcome the Resistance in a compound Engine any more than a simple one ; for that Addition superadds a Friction to every rubbing Part, therefore we must still encrease the Power as shewn in the last Rule. An Example will illustrate both Cases.

- \* Pl. 17. F. 5. THE Diameter AB of an *Axis in Peritrochio* \* is equal to 3 Foot, and that of its Axle *ab* only to six Inches ; therefore if the Power P be = 108 lb, the Weight or Resistance W will be = 648 lb. Here the Friction will be 12 lb or  $\frac{1}{6}$  of 72 lb if the
- † Pl. 17. F. 2. Axle be supported on its Surface, as in Fig. 2.† Now as 12 is  $\frac{1}{9}$  of the Power 108, there must be added  $\frac{1}{9}$  of 12 to overcome the Encrease of the Friction caused by the 12 lb, viz.  $1\frac{1}{3}$  lb, and then  $\frac{1}{9}$  of this last and so on ; so that to overcome the Resistance = 648 lb the Power must be =  $108 + 12 + 1\frac{1}{3} + \frac{4}{27}$ , &c. lb or more than 121  $\frac{1}{3}$  lb. But if the Axle had Gudgeons of Iron CC of only one Inch Diameter, the Friction on them would be six times less, the Velocity of their Surface being diminish'd in that Proportion.
- ‡ Pl. 17. F. 6. Now if there be join'd to the Axle *ab* another Wheel DE|| likewise of three Foot Diameter with an Axle of six Inches FG, about which the Rope or Chain that supports the Weight winds instead of winding about *ab*, then by mechanical Rules the Weight X = 3888 lb will be sustain'd by the Power P of 108 lb.

N. B. We here suppose the last Axle FG to have also Iron Gudgeons as c of an Inch Diameter, and that (by means of a Strap of a Leather, or a Rope, or Chain going round the

the Axel *ab* and the Wheel *DE*) the second Axis in Peri-Lect. IV. trochio *DFcGE* is carried round in the Direction *aD*, so as to draw up the Weight *X*, six times more slowly than it would rise if at *W*. But the best way is to make use of a Pinion with Leaves (that is a small Wheel with Teeth) to carry round the great Wheel by its Teeth, as here the Pinion of 12 Leaves on the Axis *ab* will carry round the Wheel *DF* which has 72 Teeth (its Diameter being six times greater than that of *ab*.) Because a Rope, Strap, or Chain cannot perform its Office without being made so tight as not to slip, in which Case the Pressure caus'd by the Elasticity would occasion an additional Friction, besides the additional Resistance made by the Difficulty there is to bend the Rope, &c.

To find the Friction of this compound Engine, we must consider all the Parts that rub, viz. first *C* the Gudgeon with  $\frac{1}{3}$  of the Velocity of the Power; then the Teeth or Strap at *ab*, with  $\frac{1}{6}$  of the Velocity of the Power; and lastly, the Gudgeon *c* of the Wheel *DE*, with  $\frac{1}{2}$  of the Velocity of the Power; which gives for the several Frictions  $\frac{2}{3} = 2$  lb,  $\frac{2}{6} = 12$  lb and  $\frac{2}{12} = 1$  lb: The Sum of all the Frictions therefore is 14,333 Sc. lb; by which Number if you divide the Intensity of the Power or 108, you will have for your Quotient 7,5 Sc. which will give you a Divisor for every Weight to be added to the Power on account of the Friction. When therefore you have added 14,333 Sc. for the Sum of the Frictions, you must superadd  $\frac{1}{7,5} = 1,33$  for the Friction of the first additional Weight, and  $\frac{1}{7,5} = 0,25$  for the Friction of the second additional Weight, Sc. Therefore in order to raise *X* or 3888 lb by means of the compound Engine here mention'd, the Power must not be 108 lb + but 124,16 lb +.

N.B. All other Cases may be deduced from this.

Tho' I have, at a Mean, taken the Friction to be equal to a third part of the Weight; yet by making many Experiments since, I found it much to vary, being sometimes greater and sometimes less; so that it is scarce possible to come at an exact Theory: Yet for Practice, as I have already said, it is useful to know what Experiments have been made with any particular View to direct us in the like Cases by taking the Mean of the different Effects we find. As Carriages are Machines of the greatest consequence for the Uses of Life, I thought it would not be unwelcome here, to enlarge upon the Friction of such Instrument  
by


Leſt. IV. *by giving an Account of ſeveral Experiments made upon Sledges and Wheel Carriages by Mr. Camus a Gentleman of Lorraine, which I have ſince tried and found to ſucceed in the ſame manner in almoſt every Trial. The little Differences I obſerv'd being unavoidable in Things of this Nature, even when made by the ſame Hand: Therefore I chooſe to give the Account of them in the Author's own Words, translated from the French, in which Language he publiſh'd his Book entitled, Traité des Forces mouvantes, pour la Pratique des Arts & Métiers, &c. Par M. de Camus.*

As there will be a Difference in the Friction of the ſame Body drawn over different Bodies, it is neceſſary as much as poſſible to know what that Difference is, in order to know how to manage or apply Forces. For this reaſon the following Experiments have been made, in order to find out nearly what Forces or Powers are requir'd to draw Weights, or what Force is loſt by Friction, and what Metal is moſt proper to be us'd, or what Materials; that we may loſe the leaſt we can of the Force of the Power; and what Effect Water, Greafe or Oil have upon different Subſtances.

FOR this Purpoſe, we muſt take three Sledges, each of them an Inch and an half wide and three Inches long, each Side or bearing Part being two Lines wide. Different Weights muſt be laid upon each Sledge, and they muſt be drawn over flat Bodies of different Subſtances or Metals; and the following Effects will be produced.

TAKE three Plates 2 Inches wide and 4 or 5 Inches long, one of Iron, one of Braſs and one of Copper; let them be draw-ſil'd, that is, ſil'd longwiſe, without being poliſh'd, and rubb'd upon ſuch a Stone as the Streets are pav'd with, that they may be like the Sledges drawn along the Streets, and alſo their Grain may lie the ſame way as it does in the Holes for Pivots and Gudgeons in Machines in regard to the Direction of the Motion. Let theſe Plates be faſten'd upon a little Plank of Oak with a Nail whoſe Head is off, that they may be put on and taken off eaſily one after another by means of an Hole in the End of the Plate. At the End of the Board fix a little Pulley with very fine Pivots, and let the Copper and Braſs Plate be rubb'd ſmooth on one Side with Pumice-ſtone.


THEN

THEN take a light filken Purse with a strong Silk to run over the Pulley and draw an Ounce Weight (including the Sledge) then prepare twenty leaden Balls weighing altogether an Ounce, that you may put into the Purse such a Number of them that the Sledge and its Load may be drawn upon the different Bodies and Metals. Let there be also prepared 20 Balls weighing altogether one Pound, to draw such a Sledge as shall together with its Load weigh one Pound; as also a third, weighing three Pounds with 20 Balls equal in Weight to three Pounds. Let the Sledges be one of Iron, or arm'd with Iron, as most of the Sledges are which are drawn along a Pavement of Pebbles; one of Wood unarm'd; the third of Lead, or arm'd with Lead; and a fourth, if you will, of Brass or Copper. Lect. IV. 

IF you put the Balls gently into the filken Purse (or the great ones in a Bag of Linnen) stopping it so that it may not move, and then the Board be lifted up at the Pulley End, so as to rise an Inch in the Length of two Foot, that the Sledge being once in Motion may not run swiftly, but on the contrary may continue at rest when it has been stopp'd in the running softly at first; but it must not be kept long in one Place, because there it would stick or sink in, and more Force would be required to set it going at one time than at another. If the Board was set horizontal, or inclining towards the Pulley, the Experiments would not succeed; because the Sledge being once set a going would run all at once: And therefore we found the best Situation of the Board was to have it rise a little towards the Pulley, and to set the Sledge a going. Things being thus prepar'd, the Effect of the Experiments was as is express'd in the following Tables.

THE first Column shews the Number of Balls requir'd to draw the Sledge dry upon different Substances or Metals as set down. Thus *Iron upon Wood*, signifies the Sledge arm'd with Iron, sliding upon the wooden Board: *Iron upon Iron*, the same Sledge sliding upon the Iron Plate; and so of the rest: *Upon polish'd Brass*, signifies the same Sledge sliding upon the Brass rubb'd with the Pumice-stone: And *upon polish'd Copper*, the Copper Plate rubb'd with the Pumice-stone.

THE Columns over which are the Words — wet — greas'd — oil'd — signify that the Plates and Sledges being *wet*, *greas'd*, or *oil'd*, the Sledges were drawn along by the Number of Balls specified according to the Columns. Thus the first Column, where you


 Lect. IV. you find these Words *Iron upon Wood* 5, above which Number is written *Balls*, signifies that the Sledge which weighs three Pounds, slides upon a Board of Oak with 5 of those Balls, 20 of which weigh 3 Pounds; so that there is only required one quarter of its Weight to make it slide along a plan'd oaken Board: But if that Board be wet, 8 Balls will be required, which shews the Friction to be increas'd the Value of 3 Balls: If the Wood be greas'd, 4 Balls and a half will do, which is but little more than half of what is requir'd when the Wood is wet: If the Wood be oil'd, there must be 5 Balls; and so on for all the Columns upon different Metals.

WE have not set down the Number of Balls required for drawing the Weight of one Pound, because the Proportion was found the same as in the 3 Pound Sledges; at least the Difference was scarce perceivable. But for the Sledges weighing one Ounce, we give an Account of their Effects, because as they differ from those of 3 Pounds, we may the better from their Comparison deduce the Causes of Friction and Resistance.

---

## A TABLE of FRICTIONS.

*A Load of three Pounds is drawn upon a Sledge one Line\* and an half wide in the Under-Plates, and three Inches long, with*

|                      | Balls          |      | Balls          |        | Balls          |          | Balls          |
|----------------------|----------------|------|----------------|--------|----------------|----------|----------------|
| Iron upon Wood       | 5              | wet— | 8              | oil'd— | 5              | greas'd— | $4\frac{1}{2}$ |
| Iron upon Iron       | 3              | wet— | 3              | oil'd— | 3              | greas'd— | $3\frac{1}{2}$ |
| Iron upon Brafs      | $3\frac{1}{2}$ | wet— | 4              | oil'd— | 4              | greas'd— | $4\frac{1}{2}$ |
| up n polish'd Brafs  | 3              | wet— | 4              | oil'd— | $3\frac{1}{2}$ | greas'd— | 4              |
| upon polish'd Copper | $3\frac{1}{2}$ | wet— | $4\frac{1}{2}$ | oil'd— | $4\frac{1}{2}$ | greas'd— | $4\frac{1}{2}$ |
| <hr/>                |                |      |                |        |                |          |                |
| Wood upon Wood       | 7              | wet— | 14             | oil'd— | $3\frac{1}{2}$ | greas'd— | $3\frac{1}{2}$ |
| Wood upon Iron       | 5              | wet— | 11             | oil'd— | 4              | greas'd— | 4              |
| upon Brafs           | 4              | wet— | 6              | oil'd— | 3              | greas'd— | $4\frac{1}{2}$ |
| upon polish'd Brafs  | $4\frac{1}{2}$ | wet— | 7              | oil'd— | $3\frac{1}{2}$ | greas'd— | 4              |
| upon polish'd Copper | 5              | wet— | 8              | oil'd— | 4              | greas'd— | $4\frac{1}{2}$ |
| <hr/>                |                |      |                |        |                |          |                |
| Lead upon Wood       | 5              | wet— | 11             | oil'd— | 5              | greas'd— | 4              |
| Lead upon Iron       | 7              | wet— | 6              | oil'd— | 6              | greas'd— | 5              |
| Lead upon Brafs      | 5              | wet— | 7              | oil'd— | 5              | greas'd— | 5              |
| upon polish'd Brafs  | 6              | wet— | 8              | oil'd— | 6              | greas'd— | $4\frac{1}{2}$ |
| upon polish'd Copper | $6\frac{1}{2}$ | wet— | 9              | oil'd— | 7              | greas'd— | 5              |

*A Sledge weighing an Ounce with its Load is drawn along, with*

|                      | Balls          |      | Balls          |        | Balls |          | Balls |
|----------------------|----------------|------|----------------|--------|-------|----------|-------|
| Iron upon Wood       | 6              | wet— | 5              | oil'd— | 8     | greas'd— | 10    |
| Iron upon Iron       | 4              | wet— | 9              | oil'd— | 7     | greas'd— | 13    |
| upon Brafs           | 5              | wet— | 6              | oil'd— | 7     | greas'd— | 13    |
| upon polish'd Brafs  | $6\frac{1}{2}$ | wet— | 7              | oil'd— | 8     | greas'd— | 13    |
| upon polish'd Copper | 6              | wet— | $7\frac{1}{2}$ | oil'd— | 9     | greas'd— | 13    |

\* N.B. A Line is one 12th Part of an Inch.

## Lect. IV.

|                      | Balls | Balls   | Balls     | Balls       |
|----------------------|-------|---------|-----------|-------------|
| Wood upon Wood       | 7     | wet— 16 | oil'd— 6  | greas'd— 12 |
| Wood upon Iron       | 6     | wet— 15 | oil'd— 8  | greas'd— 11 |
| upon Brass           | 5½    | wet— 11 | oil'd— 8  | greas'd— 12 |
| upon polish'd Brass  | 6     | wet— 12 | oil'd— 8½ | greas'd— 11 |
| upon polish'd Copper | 7     | wet— 13 | oil'd— 9  | greas'd— 12 |
| Lead upon Wood       | 7     | wet— 10 | oil'd— 9  | greas'd— 11 |
| Lead upon Iron       | 7     | wet— 8  | oil'd— 9  | greas'd— 11 |
| upon Brass           | 6     | wet— 6  | oil'd— 8  | greas'd— 10 |
| upon polish'd Brass  | 7     | wet— 8  | oil'd— 9  | greas'd— 9  |
| upon polish'd Copper | 8     | wet— 8  | oil'd— 10 | greas'd— 11 |

*The same bigger Sledges with the Weight of Three Pounds were drawn upon a Pebble, such as the Streets are pav'd with, by*

|                                  | Balls | Balls   |
|----------------------------------|-------|---------|
| Wood upon Pebble or Paving-stone | 8     | wet— 13 |
| Iron upon Pebble                 | 10    | wet— 9  |
| Lead upon Pebbles                | 16    | wet— 15 |

*The little Sledges of an Ounce.*

|                  | Balls | Balls   |
|------------------|-------|---------|
| Wood upon Pebble | 13    | wet— 18 |
| Iron upon Pebble | 9     | wet— 12 |
| Lead upon Pebble | 15    | wet— 15 |

*A Load of Three Pounds upon a Sledge whose Sides are brought to a cutting Edge where they bear, instead of the common Way, was drawn along with*

|                      | Balls | Balls   | Balls     |             |
|----------------------|-------|---------|-----------|-------------|
| Iron upon Wood       | 4     | wet— 7  | oil'd— 4  | greas'd— 3  |
| upon Iron            | 3½    | wet— 3½ | oil'd— 3  | greas'd— 3  |
| upon Brass           | 3     | wet— 3½ | oil'd— 3½ | greas'd— 3½ |
| upon polish'd Brass  | 3     | wet— 3½ | oil'd— 3½ | greas'd— 3½ |
| upon polish'd Copper | 3½    | wet— 3½ | oil'd— 3½ | greas'd— 3½ |
| Wood upon Wood       | 10    | wet— 16 | oil'd— 5  | greas'd— 3  |
| upon Iron            | 3     | wet— 7  | oil'd— 3  | greas'd— 3  |
| upon Brass           | 3     | wet— 6  | oil'd— 3  | greas'd— 3  |
| upon polish'd Brass  | 4     | wet— 5  | oil'd— 3½ | greas'd— 3  |
| upon polish'd Copper | 5     | wet— 5  | oil'd— 4½ | greas'd— 3½ |

*An*

*An Ounce Load upon a Sledge brought also to an Edge,  
was drawn by*

|                      | Balls |      | Balls |        | Balls |          | Balls |
|----------------------|-------|------|-------|--------|-------|----------|-------|
| Iron upon Wood       | 5     | wet— | 8     | oil'd— | 6     | greas'd— | 7     |
| upon Iron            | 4     | wet— | 4½    | oil'd— | 5     | greas'd— | 7     |
| upon Brass           | 4     | wet— | 7     | oil'd— | 6     | greas'd— | 8     |
| upon polish'd Brass  | 5     | wet— | 7     | oil'd— | 6     | greas'd— | 8     |
| upon polish'd Copper | 6     | wet— | 8     | oil'd— | 6     | greas'd— | 8     |
| <hr/>                |       |      |       |        |       |          |       |
| Wood upon Wood       | 9     | wet— | 16    | oil'd— | 5     | greas'd— | 9     |
| upon Iron            | 4     | wet— | 10    | oil'd— | 5     | greas'd— | 9     |
| upon Brass           | 5     | wet— | 8     | oil'd— | 6     | greas'd— | 9     |
| upon polish'd Brass  | 6     | wet— | 8     | oil'd— | 7     | greas'd— | 9     |
| upon polish'd Copper | 7     | wet— | 9     | oil'd— | 8     | greas'd— | 9     |

*With the same cutting Sledges, loaded with three Pounds.*

|                  |    |      |    |
|------------------|----|------|----|
| Wood upon Pebble | 11 | wet— | 16 |
| Iron upon Pebble | 14 | wet— | 12 |
| Lead upon Pebble | 15 | wet— | 18 |

*The little Weight.*

|                  |    |      |    |
|------------------|----|------|----|
| Wood upon Pebble | 9  | wet— | 8  |
| Iron upon Pebble | 14 | wet— | 13 |
| Lead upon Pebble | 16 | wet— | 8  |

*Upon Clay.*

- For one Pound Weight with the cutting Sledge, 8 Balls.
- For the broad wooden Sledge, only 6 Balls.
- For the broad Iron Sledge, 4 Balls.
- For the cutting Iron Sledge, 5½ Balls.
- For an Ounce Weight with the broad Sledge, 13 Balls.
- For the same Weight with the cutting Sledge, 18 Balls.



## C O R O L L A R Y I.

THE different Effects of these Experiments will also admit of a different Way of explaining them. As for example, there is proportionably more Force requir'd to draw along the little Sledge than the great one; and that probably because it glues it self or sticks to the Parts to be overcome, and also the Oil or Grease sticking to it become an Obstacle, instead of helping it as they do the great Weight, by filling up the Holes in the Wood or Metals, their Parts perhaps as they are spherical may help as so many Rollers: As Oil is not so hard as Grease, it does not so much hinder the little Weight, which requires less Force to overcome its Resistance than that of the Grease.

## C O R O L L A R Y II.

FOR this reason the great Weight runs more easily over the greas'd than the oil'd Wood; because the Grease being firmer fills the Pores of the Wood better, and keeps together the little Parts that rise when it is not greas'd, and which must be broken or press'd down when there is neither Grease nor Oil. This obliges us to employ more Force, and much more when it is wet; because the Water penetrating into the Wood causes the rough Parts to rise, which are like small Mushrooms or Grains of Corn that one must either run over or press down flat.

## C O R O L L A R Y III.

AND therefore when Wood is wet, as in the wooden Sledge, there must be as much again Force applied as when the Wood is dry; and a Force more than double would be requir'd if the Wood had been long imbib'd in Water, or was of a Nature to suck in a great deal of Water and to swell upon that account; and though Water does not seem to be imbib'd by Metals, yet it is very reasonable to think from these Experiments that it does in some measure join and glue them together; which perhaps would not happen when heavy Loads are drawn by Horses, where  
this

this Adhesion would hinder but little in proportion to the great Weight.\* Le§t. IV.

#### C O R O L L A R Y IV.

As Iron sometimes appears to run more easily over Iron than Brass, and over Brass than Copper, the Reason seems to be, that Iron being harder than Brass it is less liable to yield and sink in; and as Copper is softer than Brass, it sinks more than Brass, and consequently has more Resistance and Friction than Brass; as this appears in several Cases and several Sledges; and may perhaps do the same in the large ones, of which ours are but as Models.

#### C O R O L L A R Y V.

WE must not conclude from thence that Grease or Oil should not be us'd in Engines, tho' here they have seem'd of little use, and sometimes have been an Hindrance, as here in the little Sledge; because we know them to be of use in great Machines, for two Reasons; because they fill up the Holes as we see, or else rolling under the Weight facilitate its Motion, and they hinder the Parts from wearing or flying off, and besides they hinder such Parts as are broken off; from sticking to the Work: For if large Machines, as large Presses and Screws for coining Money, were us'd without Grease, they would move much harder on account of the small Parts that would wear off and stick on in other Places, and by that means break off more, which would not only make the Engine harder to move, but wear it out much sooner.

#### C O R O L L A R Y VI.

BUT tho' Grease or Oil seems prejudicial to small Machines, their Motion is thereby render'd more equal; and tho' one may perceive the Motion to be more difficult, as in Pocket-watches when they are oil'd, because then they generally go slower, yet then they go more equally: And it is always an Advantage to have them

\* This is a plain Consequence of the Attraction of Cohesion; for since the Attraction of Cohesion is proportionable to the Surface or the Number of the touching Parts, and the Friction proportionable to the Weight, the Hindrance or Loss of Force on account of the said Attraction will always be less in proportion to the whole Friction, as the Weight encreases.

clean'd.

Leſt. IV. clean'd and oil'd, for then they go better, and the Holes do not ſo ſoon wear wider; the Balance plays better, and is not liable to vary. One might indeed uſe no Oil in ſome ſmall Machines which go eaſily and are but ſeldom to be put into Motion.

#### C O R O L L A R Y VII.


THE Uſe of Greafe is viſible, eſpecially when Wood acts againſt Wood; for then Greafe makes the Motion as eaſy again, nay two 3ds eaſier if the Parts be ſharp and cutting, as appear'd from the Experiments made with the flat and the ſharp Sledge of Wood drawn along on Wood not greas'd.

#### C O R O L L A R Y VIII.

FROM hence alſo it appears how uſeful Greafe is for Coach and Waggon Wheels; eſpecially in rainy Weather; for if the Naves were wet, or if there was no Greafe to hinder the Water from penetrating into the Wood, four times the Force would be requir'd to draw a Burthen, than when it is drawn with Greafe in dry Weather, as may be ſeen by the Sledge, in reſpect only of the Friction made upon the Axel-tree, which is but little when compar'd with the Reſiſtance made upon the Ground or the Pavement; but beſides that Obſtacle, the Hole of the Nave would grow conſiderably bigger; for by its ſwelling the Axel-tree would no longer be free, and muſt break off the Parts in its Way: Then as the Nave grows dry the Axel will no longer fill the Hole; the Wheel will vary and be more liable to break, in thoſe Jolts that happen as it goes over a ſudden Riſe or falls into a Hole. For theſe Reaſons therefore it is always well to greaſe Machines.

#### C O R O L L A R Y IX.

THO' theſe Experiments do not wholly determine every thing that relates to very great Weights to be drawn upon Pebbles, and the Motions of great Machines; yet they ſhew us which are the eaſieſt Metals for moving on one another: As alſo that there are Parts to be overcome or broken off in Frictions: That it is merely the Weight and the Motion that cauſe the Reſiſtance and Friction, and that the Quantity of Surface does not encreaſe it when the Parts that rub and bear do not move ſwifter; for tho' the ſharp  
Sledge

Sledge goes with more difficulty in several Cases, it is not to be considered as if that way of making it would serve to avoid or ease the Friction, but as an Edge or a Saw which penetrates into the Body of Metals or Bodies to be cut ; and if it appears to slide more easily upon Wood, it is on account of the Grain of the Wood, and the Way being once mark'd it follows the Grain as in a Path, and then it has fewer Paths to press down or overcome ; but on Clay which has no Grain, or upon Pebbles, it goes much harder, because the Obstacles to be remov'd are greater, as it sinks more when sharp than when flat. 

*C O R O L L A R Y X.*

By this one may see that it is possible to encrease Friction, by making the moveable Parts so small that they will penetrate and rub off Parts of what they bear upon, as it may happen to the Pivots in Watches, if they should be too small ; for then they would soon make the Holes too big ; as likewise in Sledges, if too narrow a Bar of Iron be put on, thinking thereby to lessen the Friction on the Pavement, and so in other Cases.

*C O R O L L A R Y XI.*

Hence may be easily conceiv'd that the Bottom of Sledges ought to be broad, and it is better to shoo them with two broad Plates of Iron under each Piece, than only one narrow one ; and when the Streets are dry, it would be more advantageous to have no Iron at all, the Experiment shewing that Wood will slide better) which still appears more probable, because the little Sledge of one Ounce (with its Load) slides more easily upon the paving Stones than the great one ; whereas it goes with more difficulty on other Occasions, by reason of its being in some measure glued by its close Cohesion.

*C O R O L L A R Y XII.*

Besides this Experiment, one may also infer that the Draught will be the easier in dry Weather with the wooden Sledge ; because when Stone-dust is strew'd before Sledges, the Draught is easier by one 4th part, whether it be upon Wood, upon Iron, or upon Brass : And as there are always small Parts rubb'd off from the

~~~~~  
 Lect. IV. the Stones of the Pavement by the Nails of the Wheels and the Horses Shoes, besides the Dust at that time; all which Things will facilitate Motion, as may be known by making a Comparison with the Experiment in which neither Stone-dust nor other Dust was us'd.


C O R O L L A R Y XIII.

As for the Iron Sledge, it is evident that it would go most easily in rainy Weather; but if it was dry Weather and the Pavement not wet, it would be worth while to try with the Sledge itself (not in Model) whether then it might not be as well not to wet the Pavement, as is commonly done by a little Barrel full of Water laid upon the Sledge with two little Holes for the Water to run out and wet the Pavement and the Sledge. For tho' it is certain that if the Sledge and Pavement were sufficiently wet, all would go easier; it is very probable that it is better not to wet them at all, than to wet them so imperfectly as only to make a kind of Dirt and Lumps that may prove an Hindrance. There would not be so many Parts for the Iron of the Sledge to remove in running its Length, as was observ'd by the Experiment on the small one, which running twice over the same Track, goes much more easily the second time than the first. This may also be observ'd in whetting a Knife, which slides easily over the Whetstone after having rubb'd it over once or twice when the Stone is dry, and so wears but a little; whereas it rubs hard and wears much when the Stone is wet.

C O R O L L A R Y XIV.

WE don't speak of the Leaden Sledge; which we only us'd for Curiosity, and to shew that Parts are broken off by Friction, and that Lead causes most Resistance in many Cases, where Pivots or Arbors are made to turn upon Lead or Pewter.

ALL the foregoing Experiments are of Use to direct us how to choose Matters or Substances to make use of, and how to calculate the Force according to the Weight and the Motion which the rubbing Parts will have. Those that would try these Experiments over again out of Curiosity, must take care to make them in the same Order; first, all those that are dry, then those that
 are

are wet, next those that are oil'd, and lastly those that are greas'd ; Lect IV.
 because if the Machines were greas'd before they were oil'd, the 
 Grease filling up the Pores, the Effects of the Oil would only be
 like those of the Grease, especially when we make use of Wood,
 which must, after having been wet, be suffer'd to be quite dry,
 and new plan'd again, before it be anointed with Oil or Grease.

THO' I have in the Notes to my last Lecture * mathematical * Lect. 3.
 ly consider'd the Effects of high and low Wheels, and compar'd Ann. 11.
 'em together; yet as Coaches, Waggon and Carts, and other
 Wheel-carriages are so necessary for the Uses of Life, that only
 the Disuse of them for one Month would be enough to put a
 whole Nation in Confusion, I shall here again give an account of
 some more of *M. de Camus's* Experiments and Reasonings about
 them, being well satisfied of the Truth he asserts from my own
 Experiments, and having found his Experiments to answer as near
 as can be expected in Machines that have so much Friction as
 Models of Carts and Waggon. I have indeed a Machine with
 Brass Wheels, whose Steel Axes have very small Pivots, so
 nicely made that any of the Wheels once set in motion will turn
 upon their Axes for the Space of more than half an Hour, making
 several hundred Revolutions before they stop; but the Use of my
 Machine being chiefly to shew how near those kinds of Experi-
 ments may be brought to agree with a mathematical Theory;
 we cannot expect that any Carriage to bear Weight can have so
 little Friction: † Therefore I choose rather to relate *M. de Camus's* † Ann. 3.
 Experiments made on Models of Carriages of an Inch to a Foot,
 every way representing Carts and Waggon, and liable to as much
 Friction in proportion to this Bigness; because it shews us direct-
 ly what is the real Friction in the Carriages at present in use. In
 the manner of remedying Friction I shall add some of my own
 Considerations and Observations to his.

I begin here with his 24th Proposition, Sect. 5. of his Book
 above mention'd.


PROPOSITION XXIV.

*The Wheels of Carriages must be exactly round, and the Fellies
 at right Angles to the Naves, according to the Inclination of
 the Spokes.*

IT is a general Rule in all Cases, that the Wheels be exactly
 round; for if they were not so but like EFGH, || and the Nave Pl. 17. F. 7.

D d

out

Le^{ct}. IV.  out of the Center ; it is certain that such a Wheel, in turning, would be affected in the same manner upon plane Ground, as other Wheels are when they rise and fall, and would not be *in Equilibrio* ; the Wheel turning towards H would move with as much difficulty, as if there was a Rise to ascend, and that Height being pass'd, it would fall on the sudden as if a square Stone was roll'd along, and the Jolts of the Wheel would precipitate and push the Horses at one time, and immediately encrease their Difficulty of drawing the next Moment ; and that in proportion to the Wheels being out of round : Suppose the Wheels to be without Angles, nay truly round ; yet if the Nave should not be in the middle, * Pl. 17. F. 7. the shortest Part as * F being upon the Ground, when such a Wheel begins to turn the Weight must be rais'd in the same manner as when another Carriage is going up an Hill, and from F to D or quite to G the Wheel would act like a Wedge, and at D or G it would fall and drive on the Horses as in a steep Descent ; so that Horses or Oxen would be as much tir'd with such Wheels upon even Ground, as in straining to climb up Hill, or to bear the Shock of a steep Descent ; and this would mostly affect the Tiller or Horse next to the Wheels : Therefore Wheels ought to be exactly round, and the Naves and Holes of the Naves exactly in the Center of the Wheels.

Secondly, THE Fellies must not cross wind, but be at right Angles with the Naves according to the Inclination of the Spokes, that is, the Plane of the Circle of the Wheel, which passes thro' the Fellies, must cut the Nave at right Angles, tho' it need not pass through the Place where the Spokes are inserted into the Nave ; for otherwise the Wheel, in turning, would find Inequalities, as it happens when the Hole of the Nave is too big, and the Wheel moves from Side to Side, which comes to the same as if the Wheel was out of round : and then the Inequality of the Spokes which would be too leaning or too strait upon the Nave descending into an Hole or rising upon an Eminence opposite to their Inclination, would cause them or the Fellies to break. Therefore the Wheels of Carriages must be exactly round, and the Curves of the Wheels, or Fellies, at right Angles to the Naves.

C O R O L L A R Y I.

HENCE follows, that where Wheels are not shod with Iron, great Care must be taken to fix Wood to the Fellies in order to keep

keep them round ; and that nothing is sav'd by using no Iron ; for Lect. IV. if in such Places Wheels wear but little, because they only go upon Earth, it would only be making the Iron Plates thin, which would then cost but little ; and that Expence once over, the Wheels would be preserv'd longer and more Work would be done, so as to make the Gain far exceed the Expence. Whereas in Countries where they do not shoe Wheels with Iron or secure them with Wood, the Wheels wear so as at last to be rather square than round, and thereby the Horses or Oxen being very much fatigu'd, hardly do half the Work that they would do if the Wheels were kept to their Roundness.

PROPOSITION XXV.

The Spokes must be inclin'd to the Naves, that the Wheels may be dishing, or concave.

IF Wheels always turn'd upon smooth and even Ground, it is certain that the Spokes ought to be strait upon the Naves, that is, at right Angles to their Axes ; because then they would bear perpendicularly, like the Spokes B of the Nave AC,* which is * Pl. 17. F. 8. the strongest Way for Wood : But because the Ground is unequal, and when the Wheels fall into the Ruts, that Wheel which is in the Rut bears a greater part of the Weight than the other, because it is lower (as we have demonstrated it) in such a Case, the Spokes of a dishing Wheel become perpendicular in the Rut, and therefore have the greatest Strength ; whilst the opposite Wheel being upon higher Ground bears a less part of the Weight, and consequently the Spokes need not be at their full Strength, and so will have a sufficient Force tho' that Force be less than what they have upon even Ground. The Spokes therefore ought to be inclin'd, to make the Wheel dishing as is the usual Practice.

PROPOSITION XXVI.

The Axel-trees must be streight in all respects, and at right Angles to the Shafts or to the Pole.

IN the Motion of all Bodies, there is one Way of moving, which is the easiest of all the rest ; and that happens here when the Axel-tree is every way streight. For if its Ends should bend back-

Lect IV. ward, so as to bring the Wheels nearer together behind as AE^* ,
 * Pl. 17. F. 9. and spread them much before as DC , it is certain that they could
 not go into the Ruts nor turn in going forward, or at least with
 great difficulty, dragging instead of rolling: There would be the
 same Inconveniencies in bending the Axel-tree forward, so as to
 † Pl. 17. F. 10. bring them nearer the Pole as IF ,† and make them spread behind
 as at BD . The less the Axel is bent, the less the Inconveniency;
 but there will always be some, when the Wheels are not parallel,
 therefore the Axel ought not to be bent at all.

|| Fig. 9. IF the Wheels spread outwards as DC ||, or inwards as IF ,
 there will still be three other Inconveniencies. If the Axel bends
 * Fig. 10. outwards so that DC^* bears upon the Ground, the Way will be
 too wide; it will be hard to turn; and the Weight being drawn
 † Fig. 9. forward will crush the Wheel, the Length of the Spoke CH † in
 that Case acting as the Arm of a Leaver to break the Axel or the
 Spokes, C being the long Arm, the Center of Motion at one End
 of the Nave, and the short Arm at the other. If the Axel be so
 * Fig. 10. bent as to bring the Wheels inwards as at I and F ,* the same three
 Inconveniencies would happen; the Way would be too narrow,
 and the Weight would tend to crush the Wheels, and there would
 be a Difficulty in turning: Besides, they would bear but on the
 Edge of the Iron and become cutting by their small bearing.
 Therefore as such Inconveniencies will happen more or less ac-
 cording to the bending of the Axel; it should not be bent at all.
 But there will be no Inconveniency when the Axel is streight and
 † Fig. 11. the Wheels are in the Situation OP , AD .† By this means the
 Wheels will have Liberty as they go along; but otherwise, tho'
 a Wheel when off of the Ground might turn upon the Axel, yet
 || Fig. 9. when on the Ground and drawn at H ||, it would only drag.

THE Axel must also be at right Angles to the Pole or Shaft;
 for if the Pole or Shafts were on one side as at B , the Coach or
 Carriage would be drawn on one side, and almost all the Weight
 would bear upon one Horse; but it must be at right Angles like
 * Fig. 11. the Pole G ,* as has been said before. Therefore the Axel-trees of
 all Carriages must be streight, and at right Angles to the Pole or
 Shafts.

C O R O L L A R Y I.

THIS shews the Inconveniency of those Coaches, whose Axels
 are bent so as to make the Wheels spread upwards that they may not
 touch

touch the Braces ; for this brings on all the Inconveniencies above mention'd, and the Coaches are more liable to turn over, because the Way is narrow'd, and they go into the Coach-houses with more difficulty spreading at top, the Tops of the Wheels being more afunder than if the Axel was streight, and when they strike against any thing at top, they are more liable to break ; and then likewise the rolling of the Wheels is hinder'd ; therefore it would be better to bring in the Braces nearer together, or set the upright Wheels further afunder.


C O R O L L A R Y III.

THIS Way of bending the Axel does also render uselefs the Advantage gain'd by dishing the Wheels, as has been explain'd : for in this Case the Spokes of both Wheels bear perpendicularly at the same time as if they had not been inclin'd to the Nave ; and when a Wheel comes into an Hole or deep Rut, the Spokes being no longer perpendicular, it is more liable to be crush'd by the Weight. This makes the Wheels and Axel-trees more subject to break, and the Spokes to be loose or break. Nay, in even Ground they cut more, and bear on the Edge of the Iron Plate, as one may observe in such Wheels that the shoeing is more worn outwardly than inwardly ; and this occasions their sliding the more upon paved Streets.

P R O P O S I T I O N XXVII.


The Hind-wheels do not drive or impell the Fore-wheels ; let the Hind-wheels be ever so high, and the Fore-wheels ever so low.

IN some of the Ancients Pictures one may see Chariots represented with four Wheels all high and equal. In some Countries where Fashions seldom change, they still retain those sorts of Wheels. People, in all probability, lower'd the Fore-wheels in order to turn the more easily because of the Shafts being in the Way. The Fore-wheels have also been lower'd still more in Coaches by reason of the Braces which in some measure hinder the turning short, and are liable to be cut by the Wheels : Afterwards Crane-necks have been invented for turning yet shorter, and by degrees the Wheels made low enough to go quite under the Bend
of

Lect. IV. of the Crane-neck. Then several Coachmen pretend that their  Horses tire because the Fore-wheels are too high, so as not to be driven on enough by the Hind-wheels: And this false Principle has been follow'd even to Childrens Carts and Toys which are made with very low Wheels before. It is very likely that they would be made still lower, if it was not evident that by making them a little lower they would not go at all over some Rubs, nor get out of deep Ruts. Coachmen generally encourage lowering the Wheels, not only because of the easy turning, and their Imagination that the Hind-wheels drive on the Fore-wheels if they be much smaller; but chiefly because of their getting up easily into the Box. Now if the Fore-wheels are upon smooth and even Ground, we shall find that the Fore-wheels will be at rest tho' ever so low, the Center of Gravity of each pair of Wheels being so plac'd that the Line of Direction falls between the Wheels both in the Fore and Hind-wheels. Therefore the Carriage cannot move of it self in that Situation, only the Fore-wheels are more loaded because they are lower than the Hind-wheels; but it does not follow from that, that they roll more easily; for if it was so, Carriages would be more apt to roll when most loaded, which is contrary to Experience. A Consequence also of this Principle of *Wheels driving* would make a Carriage with very high Hind-wheels and low Fore-wheels go of it self upon even Ground, which never was; therefore the Principle is false.

S C H O L I U M.

IF the Hind-wheels could, by their greater Height, drive the Fore ones, it would follow that a Coach or Carriage would go with most difficulty when the high Wheels go foremost by making the Horses draw behind. Now let a Carriage be made in Model with Hind-wheels of five Inches, and Fore-wheels of two Inches and three Lines, which is the common Size, if we take an Inch for a Foot (tho' some Hind as well as Fore-wheels are less in Proportion) and let this Carriage or Waggon be set upon a smooth Board, and loaded in the middle with five Pound Weight of Lead; let a small Pulley be fix'd at the End of the Board, over which runs a filken Thread fasten'd at one End to the Waggon and at the other to a Scale, or a little Linen Bag to receive Leaden Balls to draw the Loaded Waggon by their Weight. The same Weight that draws the Waggon with the small Wheels foremost, will also draw

draw it with the great Wheels foremost, provided the Line of Lect. IV. Direction of the Draught be in the same Situation in both Cases:  This shews that there is no driving whatever Difference of Height there be in the Wheels, even upon plane, horizontal and smooth Ground.

C O R O L L A R Y I.

IT follows that this Notion of *driving* must have been from some Workmen who thought the Case to be parallel between a Carriage upon an inclin'd Plane, and one with high Hind-wheels and low Fore-wheels, tho' on an horizontal Plane; but the Case is very different, for on an inclin'd Plane the Line of Direction falls out of the Base, and the *Æquilibrium* is lost; therefore the Carriage will roll till it finds it or meets with some Obstacle that reduces it to an *Æquilibrium*. This would do the same in one or two Wheels as well as in four, if one Wheel was broad enough to support it self. If this be call'd *driving*, a Cart, tho' it has but two Wheels, will have it as well as a Coach or Waggon.

C O R O L L A R Y II.

IT is not a good Objection to say, that the great Wheel being in Motion continues to move longer than the small Wheel, and so drives it; for upon the Ground the Resistance is much greater than such an Impulsion. For if we give by Force some Degree of Velocity to a Waggon on plane Ground, but such as may let it sink in ever so little, as in wet Weather or upon soft Ground, not so hard as Pavement, as soon as the Force ceases to act, the Waggon will stand quite still, which shews the Hind-wheels do not drive even in that Case.

C O R O L L A R Y III.

IF we consider the Thing upon a Pavement, be it even or uneven, or upon rough Ground with Rises and Falls; let the Waggon be so plac'd as to have the high Wheels upon the higher Ground, then indeed the Waggon will run down, and they will seem to drive the low Wheels; but then in rising again, the Fore-wheels will drive the Hind ones, and the Waggon will run backwards:

Lect. IV. wards: To say that the great Wheels turning more easily will drive the better, only shews that the Waggon will go better if all the Wheels are large, and that they will altogether roll better than if there were two little Wheels. So that little Wheels before do not (whatever Way you consider them) facilitate the Motion of a Carriage.

PROPOSITION XXVIII.

Great Wheels are always more advantageous for rolling than little ones, in any Case, or upon any Ground whatever.

THE Wheels of Carriages are consider'd according to the Velocity and Friction they have upon the Axle-tree, and likewise according to their Resistance or Sinking in upon the Ground. If we consider them according to the Friction, it is certain that a Wheel, whose Diameter is double that of another, will make but one Turn, whilst the little one makes two, for the same Length of Way, the Circumference (which is in proportion to the Diameter) being double: Therefore in respect to Friction, a Wheel of double the Diameter will have a double Advantage, there being but one Turn instead of two, which doubles the Friction in the small Wheel.

* Pl. 17. F. 12, and 13. The Wheel ABC* being twice as big as the Wheel DEF, will have twice the Advantage in respect of the Friction, the Holes of the Naves and the Axles being equal.

IF we consider the Wheels according as they sink into the Earth or fall into Holes, there will be the same Advantage for the one and Inconveniency for the other: If we consider the bearing, it is double in the great Wheel, therefore it will sink but half the Way; and if we consider Hollows, it will give the same Advantage in some Cases; but then in others (as for example, where the Holes are deep) the little Wheel will have much more Disadvantage; † for if it should fall into a great Hole as DE, || of a Diameter equal to that of the Wheel, it would wholly sink in, whilst the great Wheel would only fall in the Depth of its Segment AB, * which would not be half the Wheel, as is easy to be understood by the two parallel Lines AD and BE: † We may suppose the same to happen in marshy Grounds where the little Wheel would sink wholly in the same Hole that the great one would sink but in part.

† L. 3. Ann.
11. p. 173.
|| Fig. 13.

* Fig. 12.

† F. 12 & 13.

IF we consider an Eminence to go over upon even Ground as *Leſt. IV.* a Pavement, and that it is the ſame at B as it is at E, * the Segment or the Chord of the little Wheel will be one 3d nearer the Top than the Segment or the Chord of the great one, and there muſt be a third more Force to overcome the Rub. If the Rub be ſomething which muſt be broken or cruſh'd, wholly or in part, there will be the ſame Proportion and the Circumference of the Wheel making a fort of Wedge or inclin'd Plain, it will be ſhorter or leſs acute in the ſmall than in the great Wheel, ſo that the Effort muſt be greater to overcome it all at once. And if the Rubs are only Riſes and Falls of Ground, there will, for the ſame Reason, be more Difficulty for the *little* than the *great* Wheel. Conſequently great Wheels are better for rolling than little ones on any Occaſion or upon any Ground whatſoever.

C O R O L L A R Y I.

HENCE it follows that if a Wheel be only one Inch in Diameter, or in Height, more than another, it will have more Advantage; and that the higher Wheels are, the more advantageous they are in proportion, provided they are not too high, that is, not above five or ſix Foot high; for if they ſhould exceed that Proportion, they would themſelves become a great Weight, or if made light, then they would be too weak and ſubject to break on account of the great Length of the Spokes; beſides, with ſuch Wheels Horſes would be hinder'd from exerting their utmoſt Strength by having the Axle-tree higher than their Breaſt; ſo that they would draw downwards, eſpecially if the Horſes are not very tall; as in little Wheels the Draught is made more difficult by the Horſes drawing upwards: For to deviate from an horizontal Line of Direction by drawing either upwards or downwards, is inconvenient for the Horſes, as will appear more plainly by the following Experiments.

P R O P O S I T I O N XXIX.


CARRIAGES with four Wheels, as Waggonſ or Coaches, are much more advantageous than Carriages with two Wheels, as Carts and Chaiſes.

E e

W H A T

Le^{ct}. IV. WHAT we are to consider in Carriages is the Advantage which they have one more than another in rolling, and the manner of applying Horses or Oxen in such a way as they may be the least subject to tire, and that they may draw with the greatest Advantage. Now in applying Horses to a Cart with two Wheels, it is plain that the Tiller carries part of the Weight in what manner soever the Weight is in *Æquilibrio* upon the Axle; for in going down an Hill the Weight bears upon the Horse, and in going up Hill the Weight falls the other Way and lifts the Horse (that is, pulls him upwards) which takes away great part of his Force; and if to avoid this last Inconveniency, the worst of the two, the Weight be put forward, the Horse will the sooner tire for carrying as well as drawing: Besides, as in the Holes in the Road sometimes one Wheel sinks in and sometimes another, the Shafts strike against the Tiller's Flanks, which is the Destruction of many Horses.

MOREOVER as in a Cart the whole Weight bears intirely upon two Wheels, when one of them sinks into an Hole or Rut, half the Weight falls that Way, and to draw the Wheel out of the Hole, half the Weight must be drawn out; if it be upon soft Ground, where both the Wheels sink in and must be drawn out, the Labour is greater than when the four Wheels of a Waggon sink upon the same Ground, because the Weight being distributed upon four Wheels must make them sink less than if it was only supported by two; and in Holes where one Wheel only sinks in at a time, there is only a quarter of the Weight to be drawn out, but half when we use the Cart, in which case therefore a double Force must be us'd to draw out half the Weight: If two Wheels of a Waggon fall at once into an Hole, then only half the Weight is to be drawn out, but the whole Weight when we use the Cart; and in the Rises and Falls upon Pavement, as in crossing a Kennel in a Waggon, an *Æquilibrium* is often made between the Hind and Fore wheels, those in coming down helping these to rise as they are just got over the Kennel; and if this happens only on one side, there will be the same Help; but in a Cart it would happen otherwise, for one of the Shafts would strike the Tiller in the Flank. As for the Objection, that there is less Friction upon two Wheels than four (which very likely has been the Reason for preferring Carts to Waggons) it is wholly false, for we have shewn that there is as much Friction upon two Wheels as upon Four, if there be the same siz'd Hole in the Nave, and the Weight be the same.


same. On the contrary, there will be rather more in the Use of Lect. IV.
the Cart; because as all the Weight bears upon two Points the 
small Parts will be more liable to be torn off, the wearing being
double; and if the Load on the Waggon be not greater than on
the Cart, by making the Axels and Holes of the Naves less, it will
have still less Friction; but the Friction (or at least that Difference
of it) being but little when the Wheels are well greas'd, it is not
worth Notice. Besides, the Advantage shewn in the Use of four
Wheels, we must have Regard to the Till-Horses which carry as
well as draw in the Cart, but in the Waggon exert more Strength
to draw, and yet last longer because they are not bang'd on the
Sides. Therefore four-wheel'd Carriages, as Coaches and Wag-
gons, are more advantageous than Carts and Chaises.

P R O P O S I T I O N X X X .

*IT would be much more advantageous to make the four Wheels
of a Coach or Waggon large and nearly of a Height, than to
make the Fore-wheels of only half the Diameter of the Hind-
wheels, as is usual in many Places.*

IF there be some Conveniency for turning in making the Fore-
wheels of Coaches or Waggon as little again as the Hind-wheels,
there is a very great Disadvantage, because half the Force is lost
that would be effectual if they were large, according to the 26th
Proposition. The Jolts also are greater when we use little Wheels,
because they sink as low again in the Holes and Hollows of the
Pavement, and therefore must jump as high again; and this, no
doubt, has brought People to contrive Springs to avoid the jolting,
but at the same time it has made Coaches more apt to overthrow
by raising the Body of the Coach the higher to place the Springs
under it.

BESIDES these Disadvantages, Horses that draw upwards tire
sooner and grow more stiff in the Hams; and this is the Reason
that Horses that have been us'd to a Coach are no longer fit to
be ridden, tiring their Riders very much, which would not happen
if the Fore-wheels were high, and the Points where the Traces
are fix'd were as high as their Breast, so as to draw in Lines pa-
rallel to the Ground, as Cart-horses commonly do, and thereby gain
so much as to overcome the Disadvantages that Carts otherwise
have.

Le^{ct}. IV.  THIS Advantage of Carts would cease if the four Wheels of a Waggon were equal, and then one would have the above-mention'd Advantages also of four Wheels over two. Some object, that the Horses drawing upwards lift the Coaches out of the Dirt and ease the Weight; but if they do, then they carry a Part of the Weight; and as Horses one with another are able to carry but 200 lb, but can draw near 1000 lb on a Waggon, this Way must tire them more than fair drawing when the Traces are parallel to the Ground; therefore it is best to have all the Wheels of a Coach high and equal, &c.

S C H O L I U M.

THE following Experiments will confirm what I have been explaining. Let us make use of a little Waggon or Model of an Inch to a Foot represented in the seventeenth Figure of the seventeenth Plate, with the four Wheels of five Inches and nine Lines, and so contriv'd that one may put on Wheels of different Diameters; as for example, Four of 5 Inches, Two of 2 Inches 3 Lines, Two others of 3 Inches, and let them have Naves, Spokes, and Fellies in proportion, to represent the Wheels of a Coach or Waggon. Let them be chang'd one after another, the Waggon * DB being always loaded with the same Weight A of 5 lb, and drawn by means of a filken Thread running over a Pulley, with a little Bag or Scale of a Balance to put in Balls for the different Wheels, according as they are to run on even Ground upon Earth, Sand, or Pavement. The Board AF must be of Oak, three Foot long, plan'd on one Side, and carv'd on the other to imitate the Pavements and the Kennels of Streets: The paving Stones must be of 7 or 8 Lines instead of 7 or 8 Inches, reducing them from Inches to Lines, as the Wheels are reduc'd from Feet to Inches. It must be so contriv'd, that the Pulley may be turn'd to either Side of the Board. The whole being so dispos'd, the Experiments will answer to the following Table.

To represent a Cart we hang *in Aequilibrio* under an Axel-tree, the same Weight A of 5 lb, and a Pole only is made fast to the Axel-tree to tye the String to it, which makes the Cart three times lighter than the Waggon in making the Experiments; for the Waggon has an Axel-tree and two Wheels and Shafts more than the Cart; and the full Wheels of 5 Inches and 9 Lines weigh twice as much as the Five-inch Wheels with Spokes.

To

To draw the Load of 5 lb upon the smooth Side of the Board Lect. IV. laid level with the four great Wheels, each of 5 Inches 9 Lines in Diameter, there is occasion but for three quarters of a Ball.

For the Weight of Five Pounds upon the Waggon.

	<i>Balls</i>
With the Four Wheels of Five Inches _____	1
With the Two little Wheels before _____	2
With the Wheels of Three Inches before _____	1 $\frac{1}{2}$

For the Cart and the same Weight.

With the Wheels of Five Inches _____	2
With the Two little Wheels _____	3 $\frac{1}{2}$
With the Two Wheels of Three Inches _____	3

With the Waggon upon very moist Earth.

With the Four biggest Wheels, 2 Lines wide or thick _____	12
With the same narrower and almost cutting _____	16
With the Four of Five Inches, Three Lines wide _____	6
With the Two least before _____	12
With the Two of Three Inches before _____	8

For the Cart upon the same Earth.

With the Two great Wheels _____	9 $\frac{1}{2}$
With the Two least _____	18
With the Two of Three Inches _____	13

For the Waggon upon dry Land.

With the Four Wheels of Five Inches _____	28
With the least Wheels before _____	46
With the Wheels of Three Inches before _____	35

For

*For the Cart upon dry Sand.*

With the Two Wheels of Five Inches	————	Balls 40
------------------------------------	------	-------------

When it stopp'd with 39 Balls, I was oblig'd to add 10 more Balls to make it move from that Stop.

For the Waggon upon wet Sand.

With the Four Wheels of Five Inches	————	14
With the Two least Wheels before	————	28
With the Wheels of Three Inches before	———	17

For the Cart upon wet Sand.

With the Two Wheels of Five Inches	————	17
With the Two Wheels of Three Inches	———	24

To overcome an Eminence or Rub of Two Lines, for the Waggon.

With the Four Wheels of Five Inches	————	20
With the Two least Wheels before	————	30
With the Wheels of Three Inches before	———	25

Half the Number of Balls will do when only one Wheel goes over the Rub.

For the Cart to go over the same Rub.

With the Two Wheels of Five Inches	————	35
With the Two least Wheels	————	60
With the Two Wheels of Three Inches	———	48

For the Waggon to overcome an Height of one Line.

With the Four Wheels of Five Inches	————	15
With the Two least Wheels before	————	21
With the Two Wheels of Three Inches before	———	17

For the Cart to go over the same Rub.

	<i>Balls</i>
With the Two Wheels of Five Inches _____	27
With the Two least Wheels _____	38
With the Two Wheels of Three Inches _____	31

*For the Waggon to come out of a Hole, as if a paving Stone
was wanting under each Wheel.*

With the Four Wheels of Five Inches _____	9
With the Two least Wheels before _____	18
With the Two Wheels of Three Inches _____	14
If the Weight be laid upon the little Wheels before, to come out of the same Hole, we must put in	} 34
But if it be laid upon the Hind-wheels and the Ground even, only _____	} 3

For the Cart to be drawn out of the same Hole.

With the Two Wheels of Five Inches _____	18
With the Two least Wheels _____	34
With the Two Wheels of Three Inches _____	25

*For the Waggon to be drawn from an Hole, as out of the
Channel or Kennel of the Pavement.*

With the Four Wheels of Five Inches _____	4 $\frac{1}{2}$
With the Two least Wheels before _____	8
With those of Three Inches _____	6 $\frac{1}{2}$

For drawing the Cart out of the same Hole.

With the Two Wheels of Five Inches _____	10
With the Two least Wheels _____	15
With the Two of Three Inches _____	11

For

For the Waggon upon the Pavement.

	Balls
With the Four Wheels of Five Inches _____	2 $\frac{1}{2}$
With the Two least Wheels before _____	4 $\frac{1}{2}$
With the Wheels of Three Inches _____	3 $\frac{1}{2}$
When the Wheels of Three Inches are behind, and those of Two before _____	4 $\frac{1}{2}$

For the Cart upon the Pavement.

With the Two Wheels of Five Inches _____	4 $\frac{1}{2}$
With the Two least Wheels _____	8
With the Two Wheels of Three Inches _____	6 $\frac{1}{2}$

If the Board be rais'd an Inch at the End where the Pulley is,

For the Waggon.

With the Four Wheels of Five Inches _____	4 $\frac{1}{2}$
With the Two least Wheels before _____	6 $\frac{1}{2}$
With the Two Wheels of Three Inches before _____	5 $\frac{1}{2}$

For the Cart in the same manner.

With the Two Wheels of Five Inches _____	6
With the Two least Wheels _____	11
With the Two of Three Inches _____	8

IF we make use of the Four great Wheels of Five Inches and nine Lines, about one quarter more of Force is requir'd than for the Wheels of Five Inches which are three Lines wide, and that as well for the Waggon as the Cart; because as those largeſt are very narrow and almoſt cutting, they run into the Separations of the Pavement, and after ſliding down between Two Pavements, to riſe up again there is more Force requir'd, and they go leſs ſwift than the broad Wheels, even when that Quarter of Force is ſuperadd'd and tho' they are ſo much bigger; but upon plane and ſmooth Ground, where they don't ſink in, they go much eaſier, and have more Advantage than the others.

C O R O L L A R Y I.

WHENCE it is easy to judge how much those Carters are deceiv'd who would have the Irons or Shooings of the Wheels to be made very narrow, that they may the better come out of the Ruts, and cut the Ground the more easily: For if the Wheels have no Friction on the Sides of the Fellies, being narrow they sink deeper and spoil the Ruts the more; and if they go where there are no Ruts, as on Earth, they will tire the Horses much more, one fourth part more Force being requir'd. Such kind of Wheels are very disadvantageous to every Body, because they cut the Ruts the deeper.

C O R O L L A R Y II.

THE same Inconvenience happens upon Pavement, and the Irons of the Wheels being narrow they wear out the faster, bearing in some measure but upon one Point; and as the Iron Plates wear, they grow round, and slide more strongly between the Pavement, which also breaks them easier than those that are wider.

C O R O L L A R Y III.

By the Experiments upon the smooth Board, it appears that the Friction upon the Axel-trees is inconsiderable; for with the Waggon that had the four Wheels of five Inches and nine Lines, one Ball, twenty of which weigh a Pound, draws a Weight of six Pounds, or a Weight of five Pounds, with the Wheels of five Inches: The Waggon together with the Wheels weighing besides, about one Pound and a quarter, which makes the whole Weight taken together, equal to 130 Balls. So that in this Case, one Pound would draw 130 Pounds; or, what amounts to the same thing, the Friction on the Axel would be only equal to the 130th part of the Force with Wheels in this Proportion.* For when the Ground is perfect-^{* Ann. 4}ly even, all the Resistance arises from the Friction, which is but small in comparison with the sinking in of the Wheels in Earth and Holes, from whence they must be raised.

Lect. IV.



C O R O L L A R Y IV.

By the Experiment of the Cart whose Weight bears only upon two Wheels, it appears that the Friction is double when compar'd with that of the Waggon with four Wheels; for there is requir'd double the Weight for the Cart, and even then it does not go till it be put into Motion, whereas the Waggon goes of itself; and double the Weight, or thereabouts, is requir'd for Wheels of half the Bigness. This perhaps is not altogether owing to the Friction, for it ought to be double in the little Wheels compar'd with the great ones, since they go twice round while the great ones go but once round; and tho' the little Wheels in the Cart must be set a going as well as the great ones, yet they go a little faster than the two great ones, and the two great ones in the Cart faster than in the Waggon, tho' they move in zigzag: This may also partly be owing to their not being perfectly round, nor in *aquilíbrio* upon the Axel, which is not sensible in the Waggon: And this shews that it is a great Disadvantage to make use of Carts, even in respect of Friction.

C O R O L L A R Y V.

It is easy to conceive, from the Experiments made upon hard Clay and upon Sand, that half the Force is generally lost in a Waggon when little Wheels are before instead of great ones; for tho' it does not appear that half is lost upon the Ground when the Earth is firm, there would be much more than half lost, if the Earth was soft, as we see more is lost on dry Land. Besides, one would often be mir'd with little Wheels in those Places where great ones would go through.

C O R O L L A R Y VI.

AND tho' half the Force be not lost upon the Pavement, especially when a Waggon or Coach is drawn by the Horses in a Trot, because as the Wheel goes down the Declivity of one Pavement, it acquires a Force to rise up the next; but yet if we observe the Horses drawing, we shall see that they grow heavy or stiff in the Hams drawing upwards, and on that account in these Circumstances we may also reckon half the Force even upon a Pavement; but there will be more lost on stiff Clay and Sand.

C O-

C O R O L L A R Y VII.

IF we add to this the Consideration of the hinder Axel being bent in a Coach, which makes the Wheels be less free, we need not wonder that vigorous Coach-Horses, that are well look'd after, will be fatigu'd when they have drawn two or three Hours in the Streets, and have gone four or five Miles upon the Pavement: And for the Country we must use four or six Horses, and we find that if they be back'd a few Steps they'll be out of Breath, by reason that the End of the Pole is low, and has a Tendency to break in such a low Direction; whereas the Force and Direction would be wholly employ'd in going forward or backward, if the Pole was as high as the Breast of the Horses, by having high Wheels before.

C O R O L L A R Y VIII.

HENCE we may find that Horses pay very dear for the Convenience of a short Turn, and that it would be better to go to the End of a Street for the Convenience of turning; for since half the Streets are inconvenient even for a short Turn, what would it signify if the Wheels were high, and the Pole had not the Crane-neck or arch'd Piece behind, to go a little farther, or cross a few more Streets? It would be better for the Horses, and even for the Coachmen, who would be less fatigu'd with the Jolts, and that would be sufficient amends for a little more Trouble to get up into their Boxes, which happens but once, whereas they are shak'd with a thousand Jolts. Their Masters would be less interrupted in their Business, by such Jolts as happen in the short Turns, and the Coach itself would be less liable to overthrow.

C O R O L L A R Y IX.

THEREFORE the Contrivance of a circular Piece or Crane-neck for turning, which obliges one to have little Wheels, has much more Inconvenience than Usefulness, both for the Horses and Coachmen, besides the greater Expence of the Crane-necks and the Springs, which are often the occasion of overtuning by raising up the Coach too high, as has been said.



C O R O L L A R Y X.

CONSIDERING these Experiments made with great and little Wheels, it is not hard to conceive that the Berlins are harder for the Horses than Coaches; besides the Lowness of the Fore-Wheels, the Shafts do not yield, and the Pole that bends sends back the Wheel a little when there is a Rub to be overcome, and then draws it on more swiftly; this makes the Wheel act backwards like a Wedge: If Berlins are less subject to overfet, when they do, the Fall is the greater: If they cost less, the Wheels must be oftner repair'd, because in the Berlins the Wheels cannot have long Naves; and then the Shafts break oftner than Poles.


C O R O L L A R Y XI.

THO' it may seem that there is not so much Advantage in going up and down for great Wheels as for little ones, because as they roll easily they are troublesome to the Horses when they are going down, and that the little ones not rolling so easily are not so difficult to stop; and besides, that in going up Hill, the Fore-Wheels are not so much loaded, and therefore less Force is requir'd in proportion to the great ones than upon plain Ground: Yet it appears that in going up there is always more Force requir'd for little Wheels than for great ones, and that it will always be so proportionably; for as the Direction is lower, the Horses are thereby more tired, and tho' the great Wheels are more rolling, the Horses too have the full Advantage of their Strength to stop them, the Pole being then Breast high, whereas it is very low when we use small Wheels, and it tends to break, as often happens in going down Hill: So that all being well weigh'd, there is at least as much Advantage in proportion for the great Wheels as the little ones in going up and down Hill: Besides, in travelling we go an hundred Steps upon plain Ground for one up or down Hill.

C O R O L L A R Y XII.

THERE is another Disadvantage for little Wheels, which is, that they break the Pavements and spoil the Ways more than great ones: Besides, they bear more Weight, and having less Bearing they sink deeper and jump up higher; which hurts the Houses which they shake

shake as they go by; they make more Noise and also splash more. Lect. IV°

*C O R O L L A R Y XIII.*

ONE may see the Difference of stiff Clay, Sand and Pavement, that it is always most advantageous to draw on the Pavement, and that Horses must needs tire very much on Sand; but that in Rainy Weather it is often better to go on the Sands than the common Earth when it is a stiff Clay. But in dry Weather Earth is better than Sand, Sand being more easy to draw upon when Earth is the most difficult, &c.

C O R O L L A R Y XIV.

ACCORDING to the Experiment of bringing forward the Weight upon the little Wheels, where twenty-four Balls were requir'd to draw the Load out of a Hole, instead of three when the Weight was behind and not in a Hole; it appears that there would be requir'd much the same proportional Force, on Pavement or Earth. This shews that going into the Country, it is better to put Boxes, Portmanteaux, and Footmen behind than before, which is the Reverse of what most Coachmen do, imagining, according to their Notion of *great Wheels driving* that the Coach will roll the better the more it is loaded before. Whereas the Pages who are before fatigue the Horses twice as much as the Footmen that are behind.

C O R O L L A R Y XV.

BY observing the Waggon or Cart stopp'd on the Sand, for which a quarter of Force must be added to each of them when they had time to sink, one may conclude that if a Carriage be mir'd, and the Horses are baulk'd in their drawing; we must not stand long, but put on the Horses behind to draw it out and then go thro' another Place if possible; if not, the Horses must from some little Distance be driven briskly, that the Wheels may not have time to sink, and the Horses may have acquir'd some Swiftnefs, as when we go to jump over a Ditch. In making the Experiments, we must not give the Wheels time to sink on the Sand or Earth, but lift them up every time when we put Balls into the Bag if there are
not

Leſt. IV. not enow. One may obſerve that even upon Pavement we muſt uſe more Weight when the Load has ſtood ſome time, the adheſion of Parts (or rather *taking* like the Teeth of Wheels) becoming greater both on the Axel-tree and on the Pavement, as has been obſerv'd in the Conſideration of Friction.

C O R O L L A R Y XVI.

It is alſo well when we travel upon Sand, either in dry or in wet Weather, to go in the Ruts, whereby we may avoid the Friction on the Sides of the Fellies or Curves of the Wheels, and have no Earth to turn up, and alſo the Ground is firmer there. It appear'd by the Experiment, that when the Waggon or Cart had gone twice thro' the ſame Ruts, if it was then drawn in the ſame Rut the third time, it requir'd ſcarce half the Force this laſt time, whether upon Sand or upon Clay, becauſe the Wheels then did not ſink above half a Line: Therefore in making Experiments to be compar'd together, we always fill'd up the Ruts of the Sand and of the Clay, that the Difficulty might be the ſame in the Caſes compar'd. For without that, when the Cart went after the Waggon, it always had the Advantage in the Experiments on Sand: And the Waggon would ſtill have loſt more if it had gone before the Cart upon the Clay.

C O R O L L A R Y XVII.

SEVERAL other things might alſo be obſerv'd concerning little Wheels, as to Holes, Heights to be overcome, and other Caſes, wherein there will always be found a great Diſadvantage: If it be not ſo great in the Heights to be overcome, becauſe there appears to be only the Loſs of a third Part; as the little Wheels then do not ſink deeper than the great ones; yet in other Caſes more than half the Force is loſt: So that which way ſoever we conſider that Matter, there will always be more Diſadvantage than Convenience.

C O R O L L A R Y XVIII.

As to the Carts with two Wheels, we ſee ſufficiently what Diſadvantage they have when compar'd with Waggonſ of four equal Wheels; and if they have any apparent Conveniencies, as that of loading

loading and unloading more easily, much more is lost than gain'd Lect. IV.
by saving that Labour, which ought not to be consider'd, any more
than the greater Ease of going into a Chaize with low Weeels;
for we suffer for it in another Case by the rough Jolts, and the
greater Labour of the Horse that draws.

C O R O L L A R Y XIX.

THIS Reason of loading more easily might at most obtain in Cities, where we use small Carriages and often load and unload; but for Carriages that hold their Load a Week or two without unloading, the greater Ease of loading should be considered. At that rate we might alledge that we should only use Sledges, rather than load upon a Waggon with four high Wheels: But what would be the Difference of the Profit?

C O R O L L A R Y XX.

Do not those Carters who use Carts to carry Wines with very low Wheels, that they may load with the more Ease, and then also ride upon the Horse, deserve to be made to draw the Cart themselves instead of killing Horses by needless Labour that might do great Service if work'd moderately? That Laziness is so much the more to blame, because a Turn or two more of the Handle of the Jack, or of the Windlass, would raise up a Vessel of Wine into a high Carriage, with spending very little more time, and scarce taking any more Pains.

C O R O L L A R Y XXI.

IF the Cart has some Advantages on account of the Direction of the Fore-Horses, and the Height of the Wheels, it has a great Disadvantage for the Tiller, especially upon Pavement; one Wheel coming down from a high Pavement drives one Shaft against the Horse's Flanks, then the other Wheel falling drives the other Shaft against the Horse's other Side, so that the poor Horse being bang'd about, especially in great Jolts, is soon worn out or kill'd: So that there must be a great Force to draw the Cart upon the Pavement on this account; and even when it is drawn by a String over a Pulley it goes in zigzag.

C O R O L



C O R O L L A R Y XXII.

BESIDES this Disadvantage, the Tiller carries part of the Weight; as he goes up the Weight falls back and pulls him, and in going down Hill the Weight comes upon his Back; besides he is forc'd to stop alone an immense Weight that is laid upon the Cart; so that it is a Wonder no more Tiller Horses are kill'd, tho' People generally make Tillers of the strongest Horses they can get. A Horse thus harra's'd every way cannot employ so much Force to draw as if he was before, or between the Shafts of a Waggon, where he is not bang'd upon the Sides, and has nothing to carry.

C O R O L L A R Y XXIII.

WHEN we consider how much of the Tiller's Force is lost, how much deeper the Wheels sink in than in the Waggon, how much more Force is requir'd upon Sand, Clay and Pavements; we may judge, that there is a double Advantage, or nearly, to make use of a Waggon with four equal Wheels and as high as those of Carts; for the Direction for the Horses would then be the same; and the Experiment shews us how much more Force is requir'd for a Cart upon Sand and Clay, where it goes freight like a Waggon.

C O R O L L A R Y XXIV.

SOME Carters, when they have new Wheels in their Carts, falsely attribute to the Friction of the Naves the tiring of their Horses, which are twice as much fatigu'd without appearing to do any more Work; for when the Wheels have had a few Turns they go free upon the Axel-trees; and when they are well greas'd they go as easy the second Day as any time afterwards; besides we have shewn that the Friction on the Axel-tree is very little. But the Cause of this Labour to the Horses is the quantity of Nails in the Iron Plates round the Wheels, and the great Height of their Heads, which is about an Inch. This Difficulty of rolling along when there is a Space between the Nails, is in the Experiment represented by an height of one Line to overcome, reducing the Feet of the Wheels to Inches, &c. And since it appears that there must be twenty times more Force, or thereabouts, to overcome this Height of one Inch; tho' the Nails should be but half an Inch high, one may
gues

gues how much Labour must be employ'd to raise the Weight so Lect. IV.
 at every Nail when the shoeing of the Wheels bears upon the Pavement, and then the Wheel is rais'd up upon the Head of a Nail all the while the Wheel goes round, and especially where there are narrow Gutters. This is the true Cause of the extraordinary Labour of the Horses.

C O R O L L A R Y XXV.

THIS shews that it is necessary the Wheels should be round, and the Carters who think they save Money in using many great Nails (because the Iron Plates or Shoeing does not wear so fast when there are such Nails) are in an Error, and lose double by doing less Work and fatiguing their Horses.

C O R O L L A R Y XXVI.

THESE Nails also have the same Effect as narrow Plates, they slide between the Pavement and from thence they have a greater Height to rise in lifting the Weight up, and sometimes sliding down again, and that wears the Plates round, which makes them slide more between the Pavements and into the Ruts which the Kennels make in the middle of the Streets; and it is also upon this account that narrow Plates soon grow round, and being round tire the Horses almost as much as if there were Nails.

C O R O L L A R Y XXVII.

THIS shews plainly enough, that it is advantageous to have the Plates upon the Fellies of the Wheels wide, as well on the Pavement as on the Ground. as it appear'd by the Experiment of the great narrow Wheel upon the Pavement, and also considering the Strength that the Horses must exert to draw out the Wheels from the Kennels, the Damage that is done to the Pavement, and the quick wearing of narrow Plates, which also soon become round and inconvenient. It wou'd be better to have the Kennels like Troughs, wide and shallow, * as IE Fig. 14. and not like LM Fig. 15, which is the common Make of the Kennels in the Streets.

* Plate 17.
 Fig. 14 & 15.



C O R O L L A R Y XXVIII.

IF Kennels were made in this Shape several Advantages wou'd be gain'd: The Plates or Shooings of Wheels would not so soon grow round, nor break the Paving so easily: The Pavement of the Kennel itself wou'd last longer because it wou'd not receive such great Shocks, every part supporting more equally; whereas now the lowest Place bears much a greater part of the Weight, whereby it sinks most and sooner breaks. This sinking in makes those Places lower when they are mended, and the Rain-water stays in them so that the Wheels or the Horses Feet striking into them, splash the People that walk along, whereas the Water wou'd always run down into the middle of the Kennel, were it not for these Inconveniencies: neither wou'd the Stream of the Kennel-water grow so wide as it does in great Showers, because in the Form we propose, the Kennel is wide enough at bottom to carry off the Water; and it would be easier in walking to step over them.

C O R O L L A R Y XXIX.

THUS also might the little Channels, that bring the Water from the Houses to the Kennels in the middle of the Street, be made less: Nay it wou'd be better to be without them, only making a gentle Declivity from the Houses to the middle of the Street. These little cross Kennels are very inconvenient for those that walk the Streets, nay for the very Horses and those that go in Coaches: People don't consider that one must stop a little at every one of these Channels and go irregularly, taking first a little Step and then a great one to stride over; and a little Stop of the Center of Gravity or of a Body in Motion tires one very much: This is the Reason why People are more tir'd to walk an Hour in *Paris* than two or three in the Country even upon Pavement, because there the Body always continues its Motion without Interruption. The same might happen in *Paris* if there was only one Kennel pretty wide and shallow in the middle of the Streets; then also Persons that go in Coaches wou'd be less jolted.

C O R O L L



C O R O L L A R Y XXX.

THE Use of Waggon's with regard to pav'd Streets and to publick Roads would also be of publick Advantage, instead of Carts, which are very weighty in proportion to their Bigness, and which are generally loaded so heavy as to endanger the breaking of all the Geer. The vast Weight which they carry being supported on two Points, as they have but two Wheels, sinks deep and makes great Ruts in the Ground; and upon Pavement when the Wheel is born up on a Nail Head and comes to fall down again, it presses down or breaks that Pavement worse than if ten or twenty Waggon's with as heavy a Load had pass'd over that Place. For a Waggon Wheel, that falls upon the same Place and from the same Height, comes down but with half the Weight that the Cart Wheel does, and therefore strikes it but with half the Force. Now the Pavement being something of the Nature of Glass, will not break unless it receives a sufficient Blow, and an hundred Blows, if each of them be less than that sufficient Blow, will not break it. The Blow from the Cart in those Places is often sufficient to break the Pavement where an hundred Blows from a Waggon Wheel, as they strike but with half the Force, will not be able to hurt it: So likewise will a Waggon pass over as often as you will where a Cart would sink in. Carts therefore, little Wheels, narrow Plates for shoeing the Fellies, and great Nails, are to be avoided as much as can be.

C O R O L L A R Y XXXI.

By the Experiment made upon Clay and upon Sand, we may see how useful it is to have the Roads firm and solid, since there is so much Difference between the one and the other. And certainly the Carriage of Goods, by Land or Water, contributes much to the artificial Riches of a Kingdom or State: And all things well consider'd, we shall find that the Time and Expence employ'd in mending Roads, is the most for the publick Advantage.

C O R O L L A R Y XXXII.

In this and the other three following Corollaries Mr. CAMUS finds fault that the Laws and Orders concerning mending Roads and Matters relating to Carriages are no better observ'd, and

Leſt. IV. *takes notice that a Farmer that uſes many Horſes to draw a large Cart very heavy loaded, may ſpoil the Roads for many Miles, worſe than an hundred others with the common Carriages, and therefore propoſes in*

C O R O L L A R Y XXXVI.

T H A T Orders ſhould be made for the publick Good, That only two Horſes ſhould draw in a Cart and four in a Waggon, which Four would do more Work than ſix in a Cart.

C O R O L L A R Y XXXVII.

T H A T Wheels of Carriages never be made leſs than four Foot and a half, or five Foot in Diameter ; that the Plates of Iron never be leſs than three Inches wide, the Fellies three Inches and a quarter deep, at leaſt for Coaches, and four Inches for Waggons or Carts, which by that means would be ſtronger as they are broader.

C O R O L L A R Y XXXVIII.

T H A T the Nails for the Plates be made without Heads, and that a great many be made long enough to go quite thro' to be rivetted, which will keep the Irons from riſing up. For two Nails ſcrew'd behind, or with a Counter-Rivet will hold the Irons on faſter than ſix with Heads. One may as well make the Plates three Inches wide as two, and as they wear out more at the Edge, they may be made thinner in the Middle next the Fellies, in the Shape * Pl. 17. F. 16. ſhewn in the Section, *Fig. 16.* at B C, * which may be eaſily done in the forging by help of a Swage of that Shape. Theſe Plates being put on hot will preſerve the Fellies more from cracking, laſt as long as others, and neither be heavier or coſt more. By this means the Roads would not be ſo ſoon ſpoil'd, and all that uſe Carriages would find their Advantage.

C O R O L L A R Y XXXIX.


T H E Naves of Coach Wheels ſhould be made a little thicker in the middle, and not be bor'd quite thro' where the Spokes are let in, for then their Ends wou'd not receive the Greafe, and thereby be ſo apt to get looſe: The Naves might alſo be made fifteen or ſixteen Inches long,

long, with the great End a little less, since the greater Length of the Nave does not encrease the Friction, and it holds the Wheel straighter. The Spokes should also be made an oblong Square at Bottom, to have a Shoulder to keep them firmer than when they are round, as they are usually made. The Spokes thus fix'd would not be so apt to fly out nor make that rattling which they do, especially in Summer-time. The Workmen may work out these Shoulders with a Bevil at the same Angle that the Spokes are let in, to make the Wheels dishing out, after the manner the Joiners make their Skew Tenons, and the same Instrument would direct them in cutting the Mortaise, &c.

Mr. CAMUS ends his Considerations upon this Subject with observing that a Post Chaise with two Wheels has all the Inconveniencies observ'd in Carts, and besides has the Inconveniency of the second Horse which draws on one Side; therefore, for the Ease of the Horses as well as the Travellers, he would have all those Chaises to have four Wheels, and the Driver not to ride, but sit on a Box like a Coachman, and the Fore-Wheels as high as the hind ones.

BESIDES the Friction already explain'd, there is another Impediment to Motion in several Engines, and that is the Difficulty with which Ropes are folded, which encreases according to the Bigness and Stiffness of the Rope, the Weight which they bear, and the Smallness of the Diameters of the Bodies about which they wind. This Impediment we shall also call Friction, because we must make Allowance for it as well as for the rubbing of the Parts of the Machine, otherwise we shall always find the Effect of Powers by means of Engines to be less than we expect.

Monfieur *Perault*, in his Comment upon *Vitruvius*, describes an Engine of his contriving, whereby he thinks to avoid all Friction. This Machine is an *Axis in Peritrochio* applied a new Way. But as he is not aware of the Friction or Hindrance that arises from the Difficulty of bending the Ropes, the Effect will by no means answer as he proposes; for upon Tryal, his Machine appears to have more than double (sometimes than triple, or quadruple) the Friction of the same Engine us'd in the common Way, when the Pivot or Iron Axis is in Diameter the 12th Part of the Roller or wooden Axel. He does indeed say that the Machine was try'd and succeeded;


 Lect. IV succeeded; but having carefully examin'd it, I found it impossible to answer according to the Description; and lest at any time any body should be at the fruitless Expence to make the Machine in large, I demonstrated the Disadvantage of it before the *Royal Society*, and shew'd that besides the Friction (overlook'd by its Author) there was a great Inconvenience in the Application: And this I confirm'd by Experiments made on a Model of an Inch to a Foot, the large Pulley or Wheel being suppos'd of Five Foot.

Monsieur *Perault's* Account of his Engine is as follows: "In
 " Imitation of the (modern) Crane, I have invented two Engines
 " for raising Weight. The first is made of that Organ which is
 " the most advantageous of any in Mechanicks for facilitating Mo-
 " tion; because it is free from that Inconveniency which we meet
 " with in all others; namely, the Friction of the Parts of the Ma-
 " chine, which renders their Motion more difficult. This Organ is
 " the Roller, which *Aristotle* prefers to all other Organs, because
 " all the others, as Wheels, Capstanes, and Pulleys, must necessarily
 " rub in some of their Parts. But the Difficulty was to apply the
 " Roller to an Engine that raises Weights, its Use having only been
 " hitherto to cause them to roll on an horizontal Plane. The Engine
 " which I propose has a Base AAB,* (*Plate 17. Fig. 17.*) some-
 " thing like the Crane; This Base has in its upper part the horizon-
 " tal Piece B, which clasps an upright Shaft CO, supported under
 " its Pivot O, on which the whole Engine moves in the same man-
 " ner as the Crane, when the Weight is to be lower'd. This Shaft
 " supports on its Top a cross Piece DD, to which are fasten'd the
 " Ropes EE, which wrap round the Barrel, Axel, or Roller F,
 " which has another Rope G, that also wraps or winds round one
 " of its Ends. This last Rope is that which raises the Weight. At
 " the other End of the Axel there is a great wooden Wheel like a
 " Pulley HH, about which is wound a long Rope N. To work
 " this Engine, one must pull the long Rope N, which causing the
 " great Wheel to turn, does also carry round the Axel or Barrel,
 " which is made fast to it. This Axel, as it turns round, causes the
 " Ropes EE to wind about it, and therefore the Axel and the Wheel
 " rise, whilst the Rope F, to which the Weight is fasten'd, does al-
 " so wind itself upon the Axel the contrary way; and this double
 " winding up of the Ropes makes both the Burthen and the Axel
 " and Wheel to rise at the same Time. Now it is evident, that all
 " this Rise is perform'd without the Friction of any Part, and conse-
 " quently,

* Plate 17.
 Fig. 17.

“quently, the whole Power which draws the Rope N, is employ’d Lect. IV.
“without any Hindrance; which cannot be in other Engines.”

“It may be objected that the Power which acts at N, must, besides the Weight, raise also the Axel and great Wheel, and that their Weight is one of those Obstacles which *Aristotle* says all Engines are liable to; and that this Obstacle is equivalent to the Friction which is in other Organs. But it may be answer’d, that Friction is an Obstacle wholly unavoidable in all other Organs; but that it is easy to remedy the Obstacles of this, which is done by means of the heavy Body M, taken equal in Weight to the great Wheel and Axel, which it sustains by means of the Rope II, which running over the Pulleys L L, is fix’d to the Ring or Collar K that goes round the Axel F. For the Axel and the Wheel being counterpois’d by this Weight, the Power which acts by drawing the long Rope N, acts for raising the Weight only. The Experiment which was made with this Engine has confirm’d the Truth of this Problem, by comparing its Effects with those of a Crane, in which the Proportion of the Bigness of the Axel to the Circumference of the Wheel, was the same as in my Machine. For it happen’d that in the Crane, a Weight of *One* hanging at a Rope going about the Wheel, drew up a Weight of *Seven*, when it had one Half added to make it preponderate, or give Motion to the Power: And when the Weight to be rais’d, and the Weight which serv’d as a Power, were proportionably encreas’d, there was also a Necessity to encrease the additional Weight, which made the Power preponderate in the same Proportion: So that as it was requir’d to add one half to the Power when the Weight was *Seven*, the Addition to the Power became *One* for a Fourteen Pound Weight, *Two* for a Twenty-eight Pound, *Four* for a Fifty-six Pound, and so on; because the Resistance from Friction encreases nearly in the same Proportion that the Weights are encreas’d. But this did not happen in my Engine, in which *one Quarter* was alwas sufficient for the Draught (or to make the Power preponderate) not only when the Weight was *Seven*, but also when it was Fourteen Pounds, twenty-eight Pounds, Fifty-six Pounds, &c. which evidently shews, that this Engine acts without Friction.”

THUS far *Monsr. Perault*. But however plausible this Description may appear, a little Attention will shew, that if this new Engine

Leſt. IV. gine had no Friction, yet it is more inconvenient than an *Axis in Peritrochio*, with the ſame Proportions; and likewiſe that it has
 * Pl. 18. F. 1. more Friction than the ſame Machine in the common Uſe. A C E * (Fig. 1.) is a common *Axis in Peritrochio*, which has the Wheel A E, five times bigger in Diameter than the Axle; ſo that A C the Radius of the Wheel (which is the Diſtance of the Power) is to C B the Radius of the Axle (the Diſtance of the Weight) as 5 to 1: Conſequently *One* (for Example one Ounce, as in our Experiment) will keep *Five* in *Æquilibrio*. Now tho' the Friction of the Gudgeon at C is unavoidable, yet it may be diminiſh'd by diminiſhing the Diameter of the Gudgeon †. Provided it remains ſtrong enough to ſuſtain the Machine and its Burthen. Here one Penny Weight, or $\frac{1}{2}$ of the Power added to it, makes it preponderate, and give the Machine Motion with a due Velocity.

† Ann. 4.

Now this very Engine made Uſe of in *Monſ. Perault's* Way, does ſo alter the Diſtances of the Weight and Power, that inſtead of *One* for our Power, we muſt have *Two and a half* to keep the very ſame Weight *Five* in *Æquilibrio*, as may appear by a Sight
 ** Pl. 18. F. 2. of the ſecond Figure,** where ſince in the Action of the Machine, when we pull the Rope P A, we make the Axle D B to wind itſelf up upon the Rope H D, it is evident that D is now become the Center of Motion, D B (the whole Thickneſs of the Axle) the Diſtance of the Weight = 2; and the Diſtance of the Power is reduc'd to A D = 4. So that if two Men, having been employ'd in the common Way to raiſe Weights equal to the Strength of ten Men, an Engineer ſhould alter the manner of working, and fit up the *Axis in Peritrochio* in *Monſ. Perault's* Way, inſtead of gaining an Advantage, he muſt call in three more Men to perform this Work. †† If it be answer'd, that what is loſt in Strength will be gain'd in Time, it may not only be ſaid, that one cannot always call in more Help on a ſudden, but that even then, tho' we ſhould not call this an Inconveniency, yet there will be ſtill more Friction in this than in the common Method; for the Roller or Axle will find a Difficulty to wind on the Ropes, becauſe they are not perfectly pliable, and the leſs ſo, the greater the Weight is that ſtretches them. This, together with the Friction of the Collar of the Rope of the Counterpoiſe to the Engine, makes the Hindrance greater than in the common Way. For it appears by my Experiments, that when the Power is become equal to $2\frac{1}{2}$ to keep the Weight 5 in *Æquilibrio*, there muſt be added $\frac{1}{2}$ (here four-penny Weight) to put the Power in Motion. And

†† Ann. 5.

AND, to shew that this Friction of the Ropes is not always the same as Monf. *Perault* supposes it; when P (or the Power) is made only one Ounce, and W (or the Weight) 2 Ounces, then to make the Power preponderate, only 2 Penny-weight and 18 Grains was sufficient. But When P is $= 2\frac{1}{2}$, and W = 5, the additional Weight mark'd $\frac{1}{2}$ was 4 Penny-weight and 2 Grains.

It is plain from this, that Monf. *Perault's* Experiments were very inaccurately made, and therefore not to be depended upon.

I have been the more particular here; because we are apt to be led into an Error by the Over-sights of a Man of great Reputation, whom we don't easily suspect of a Mistake.

THO' it be as difficult, at least, to give a certain account of the Forces requir'd to bend Ropes of different Diameters, (stretch'd by different Weights, in making them go round Bodies of different Bignesses) as to give an exact Theory of Friction: Yet to consider nothing of the Loss of Motion occasion'd thereby, would be as prejudicial to the Practice of Mechanicks, as it would be to overlook the Friction of the Parts in Engines. Therefore, tho' the different Materials of which Ropes are made, their different Stiffness, according as they are more or less twisted, and sometimes the Temperature of the Weather (as to Moisture and Dryness) at the Time that they are us'd, makes it very difficult to be exact in our Conclusions; yet we think it is of great Use to give the best Theory we can, and mention some of the Experiments, at a Medium, made upon Ropes pretty good in their kind and moderately twisted; because if any part of a Rope of any Length of equal Thickness and even Twist from End to End, be stretch'd by a known Weight round a Cylinder, Roller, or Pulley, and it be observ'd what Force will bend it about a Roller of a given Diameter, we may know, what other Force will be requir'd to bend it round any other Body, and when stretch'd with a different Weight: and besides, in new well made Ropes, the Difficulty of bending, *ceteris paribus*, is pretty near as the Diameters (not the *Solidities*) of the Ropes.

EXPERIMENTS, *Plate 18. Fig. 3.*

To two immoveable Hooks R R I fix'd the two Ropes R r, R r, at the Distance of about 8 Inches from each other; and at the lower

H h

End

Lect. IV. End of the Rope I hung the Scale *SS*, on which I plac'd the Weights *W* to stretch the Ropes. Then I took successively three Cylinders like *CC*, each one Foot long, (one of half an Inch, another of an Inch, and the third of an Inch and an half Diameter) and having wrapp'd the two Ropes about one of the Cylinders, as may be seen in the Figure, by putting Weights in the little Scale *s* by means of the Ribbon *m* I brought down the Cylinder towards *W*, always taking care that the Parts of the Ropes did not rub against each other, and rolling the Cylinder up and down two or three times before I settled the Weight that I observ'd to bring down.

Pl. 18. F. 3.

N.B. The Weight includes the Weight of the Cylinder and Scale.

† Pl. 18. F. 4.

IN the 4th Figure † *C* represents the Section of the Cylinder or Roller about which the Rope *R r* is wound, and *KL* its Diameter, and *ms* the little Scale and Ribbon as before.

THE Weight *W* may be always look'd upon as the Weight stretching the Rope; because (tho' it stretches the two Ropes, and consequently each Rope is stretch'd but by half of it) the Cylinder *CC* * brought down by the Weight *s* bends or causes a folding in the two Ropes at *C* and *C*, which gives it the same Difficulty of Descent, as if only one Rope bearing the whole Weight was folded about it.

* Pl. 18. F. 3.

A TABLE of EXPERIMENTS, shewing what Forces were requir'd to bend Ropes of different Diameters, stretch'd by different Weights, round Rollers of different Bignesses. Left. IV.

<i>The Quantity of the Weight W, by which the Rope is stretch'd, express'd in 16 Averdupoids.</i>	<i>The Resistance of the Rope about a Cylinder of half an Inch Diameter, express'd in 3 Averdupoids.</i>	<i>The Resistance of the Rope about a Cylinder or Roller of one Inch Diameter in 3 Averdupoids.</i>	<i>The Resistance about a Roller of one Inch and half Diameter in 3 Averdupoids.</i>	<i>Diameters of the Ropes of 3 Strands or Twists, express'd in tenth Parts of an Inch.</i>
60 lb—	225 $\frac{3}{4}$ 90 — 45 —	112 $\frac{1}{2}$ $\frac{3}{4}$ 45 — 22 $\frac{1}{2}$ —	75 $\frac{3}{4}$ 30 — 15 —	— 0,5 — 0,2 — 0,1
40 lb—	150 — 60 — 30 —	75 — 30 — 15 —	50 — 20 — 10 —	— 0,5 — 0,2 — 0,1
20 lb—	75 — 30 — 15 —	37 $\frac{1}{2}$ — 15 — 7 $\frac{1}{2}$ —	25 — 10 — * —	— 0,5 — 0,2 — 0,1

* *The Experiment could not be made here, because the Cylinder of 1 $\frac{1}{2}$ Inch Diameter, which should have been us'd here, weigh'd above 8 Ounces; and the Weight requir'd to bend the Rope, appears by Analogy to be but 5 Ounces.*

OBSERVATIONS on the foregoing Table.

THE Experiments mentioned here are taken at a Medium from a great many that I made, the Difficulty of bending the Ropes being sometimes a little less, and sometimes a little greater.

ONE may from this Table make a more extensive one, by putting in Numbers analogically for several other Thicknesses of Ropes and larger Diameters of Cylinders or Rollers; but in Ropes of a Diameter bigger than half an Inch, a Roller of half an Inch is too little and hardly ever us'd in Practice; nay, in such Cases the bending of the Ropes is more difficult than in Proportion to the other Experiments; and in my Tryals, the Rope of 0,5 of an Inch Diameter generally requir'd more than here set down.

Lect. IV. I FOUND that a woven Clock-line of 0,1 Inch Diameter requir'd more Force to bend it than a twisted Rope of the same Thickness which seem'd much stiffer; but if we consider that the woven String flattens as it folds round a Cylinder; that has the same Effect as if the Cylinder was become less, one Fold of such a Line making by its central Parts a less Circle than a twisted Rope, which does not become flat.

MONS^r. *Amontons* made several Experiments of this kind, mentioned in the Memoirs of the *Royal Academy* of Sciences at *Paris*, for the Year 1699, and calculated a Table of the Force requir'd to bend Ropes, which Table I cannot recommend, because it is built upon a Mistake; for he says, that the Difficulty of bending a Rope of the same Thickness, and loaded with the same Weight, decreases when the Diameter of the Roller encreases, but not so much as that Diameter encreases; but I have found by many repeated Experiments, that the Difficulty decreases directly as the Diameter of the Roller encreases. That is, *The Difficulty of bending a Rope round a Roller, is, cæteris paribus, inversely as the Diameter of the Roller.* N. B. I believe Mons^r. *Amonton's* Mistake arose from the Parts of the Rope rubbing against each other, which I always took care to avoid.

WHEN a Rope is carried under or over the Wheel or Sheave of a Pulley, the Difficulty of bending is as great as if it went quite round a Roller, which will appear by observing the 4th Fig. of *Plate 18*; For while the Rope is folding about the Cylinder C in the Direction *r* K L, the under Part L o K unfolds of itself without any Difficulty.

I HAVE in this Lecture (Page 187, 188, 189) given Rules for the Friction in a compound Engine, and shew'd the Application of those Rules by an Example; but did not then take in the Difficulty of the folding of Ropes. But now I will give another Example wherein that also is consider'd, and shew how near the Theory agreed with the Experiment perform'd with a Machine made as nicely as possible, to make the Comparison the juster.

The Machine consists of three Pulleys (two *upper* and one *lower*, or a *Tackle of three*) whose Diameters are exactly as follows, *viz.* two Inches, one Inch and a quarter, one Inch and an half; and

and all the Center-Pins of one quarter of an Inch Diameter exactly, and the Rope of one tenth of an Inch Diameter. The Weight is 18 Pound *Averdupois*, and consequently the Power to keep it in *Æquilibrio* must be 6 Pounds, and a very little more must make the Power raise the Weight, if there was no Friction; but here no less than 20 Ounces are requir'd, tho' the Machine be very exact.

I HAVE shewn (in *Rule 2. Page 187.*) that two thirds of the Power are equal to the Friction of a Cylinder whose Surface moves as fast as the Power, and whose Gudgeons are equal in Diameter to the Cylinder. Now as the Diameter of the first Pulley 1 is eight times bigger than its Pin, its Friction must be 4 Pound divided by 8, or 8 Ounces; because as the Surface of its Circumference moves with the same Velocity as the Power, its rubbing Surface on the Pin must move eight times slower.

THE second Pulley 2, whose Surface moves as slow again as the Power, and whose Pin is six times less in Diameter than itself, must of consequence have its Friction only of $5 \frac{1}{3}$ Ounces; because two thirds of the Power, or 64 Ounces, must first be divided by 2, by reason of the Velocity of the Pulley's Surface being but half that of the Power; and then again by 6, because the Pin being six times less, the Parts rubbing on the Pin must still move six times slower. So that $\frac{64}{2} = 32$, and $\frac{32}{6} = 5 \frac{1}{3}$ Ounces.

THE third Pulley 3, moving with a third of the Velocity of the Power, 64 Ounces, must be divided by 3, and that Quotient again by 5, because the Pin is here $\frac{1}{5}$ of the Diameter of the Pulley; so that the rubbing Parts of this Pulley have their Velocity one 5th of a 3d, or $\frac{1}{15}$, of the Velocity of the Power; and therefore $\frac{64}{15}$ will give 4,26 $\frac{2}{3}$ Ounces. Now the Sum of all these Frictions (*viz.* 8 Ounces + 5,333 $\frac{1}{3}$ Ounces, + 4,26 $\frac{2}{3}$ Ounces) makes 17,6 Ounces, which is the $\frac{5}{10}$ th & $\frac{4}{10}$ part of the Power. This Addition to the Power will so encrease the Friction, as to require a Super-addition of the $\frac{5}{10}$ th & $\frac{4}{10}$ part of that first Addition, and so on in this Series, 17,62 Ounces + 8,8 Ounces (which is $\frac{37,42}{2}$) + 0,59 Ounces (which is $\frac{37,42}{64}$) $\frac{37,42}{64}$ in all 21,41 Ounces.

To his must be added the Friction or Resistance on account of the Difficulty of bending the Ropes, which by the last Table may be found in the following manner.

SUPPOSE

Lect. IV. SUPPOSE the Part of the Rope, which is at 1 the Side of the first or upper Pulley to be fix'd, then will the 3 Ropes D E, 3 B, and 2 A sustain together the whole Weight W, * which (together with the Block) weighs 18 Pounds, so that we may consider each Rope as stretch'd by six Pounds, and folding round different Cylinders of the respective Diameters of the Pulleys.

*Pl. 18. F. 5.

FOR the first Pulley, we look in the Table for a Rope of one tenth of an Inch, and find that when stretch'd with the Weight of 20 Pounds it requires 7,5 Ounces to bend it round a Roller of an Inch, therefore we must first make Use of this Analogy.

As 20 lb stretching a Rope of one tenth of an Inch in Diameter:

Is to $7\frac{1}{2}$ Ounces, the Force requir'd to fold it on a Roller of one Inch Diameter::

So is 6 Pounds, when so much only stretches the Rope :

To 2,25 Ounces (that is, two Ounces and a quarter) the Force able to fold the same Rope, only stretch'd with 6 lb, round a Cylinder of one Inch Diameter.

BUT as the first Pulley is not of one but two Inches Diameter, we must diminish the Force necessary to fold the Rope in a reciprocal Proportion of those Diameters, by the next Analogy, which will also serve for the two other Pulleys.

As the Diameter of the Pulley, where we want to know the folding Force :

To the Diameter of the Pulley where the Force is known::

So is the folding Force before found:

To the folding Force requir'd.

That is, as 2 : 1 :: 2,25 : 1,125 Ounces, for the 1st Pulley ;

And, as 1,5 : 1 :: 2,25 : 1,5 Ounces for the 2d Pulley ;

And lastly, as 1,25 : 1 :: 2,25 : 1,8 Ounces for the 3d Pulley.

By such Analogies we may encrease the little Table above to any other Proportions, so as to shew by Inspection the Force requir'd to

* It is true, that in Motion, one Rope bears more of the Weight and another less of it ; but upon the whole, the different Difficulties of folding the Ropes taken together, will amount to the very same. So that this is the best, because the easiest way of considering it.

to fold Ropes in most Cases; or most Cases may be deduc'd from **Lect. IV.**
this Table itself.

BUT to go on with the Theory of the Friction of our Machine: These three last Frictions, or Resistances of Ropes added together, make 4,425 Ounces, which added to 21,41 Ounces the Friction above found, gives in all 25,835 Ounces, a Friction greater by near six Ounces than the Experiment gave. But I have demonstrated in the Annotation to my third Lecture (Page 164) that, *When a String or Rope runs over a single Pulley or Roller by the Descent of the preponderating Weight (the other Weight rising at the same time) the Pressure on the Axis of the Pulley is always equal to the Quadruple of the Product of the Weights multiplied into one another, and divided by the Sum of the same Weights.* And that Pressure being always less than the Sum of the two Weights, when they are unequal, so much as it is less must be taken out of the Account of all the Frictions, and the Experiment then will be extremely near the Theory. But to omit nothing in our Calculation, we will also examine what this Diminution of Pressure is.

THE Power (6 Pounds) together with what we have found necessary to be added to it on account of all the Frictions and Resistances of the Ropes, is to be look'd upon as the preponderating Weight in the Case of the Proposition abovementioned, that is, 6 Pounds and near 26 Ounces, or 122 Ounces: and 6 Pounds without any Addition is to be considered as the Weight overpoiz'd on the other side of the first Pulley 1, which is an upper Pulley. Let the two Weights be multiplied together, that is, 6 Pounds or 96 Ounces \times 122 Ounces, the Product of which is 11712 Ounces; which again multiplied by 4, or 11712×4 gives the Product 46848; then dividing the last Product by 218 (the Sum of the Ounces in both Weights) or $\frac{46848}{218}$, the Quotient will be 214,9, which subtracted from 218 the Sum of the Weights, will give 3,1 Ounces the diminished Pressure, or that part of the Pressure which the Pin of the Pulley 1 is freed from when the Power preponderating runs down.

NOW as there is another upper Pulley over which the Rope also runs, there must be likewise taken off on that account 3,1 Ounces: So that if these two last Sums, or 6,2 Ounces be taken from 25,835 Ounces

Leſt. IV. Ounces found by Theory equal to all the Frictions, there will remain 19,635 Ounces, the Addition neceſſary to make the Power over-balance the Weight with the leaſt Augmentation poſſible; and that in the Experiment is 20,365, or a little more than one third of an Ounce; for 20 Ounces added to the Power 6 Pounds makes it run down.

N.B. *Nothing was here allow'd as an Encrease of Friction on account of the Weight added to bend the Ropes, which would ſtill bring the Experiment nearer the Theory.*

IN Practice we need not make this laſt Allowance, or conſider this Diminution of Preſſure, eſpecially in Tackles of many Sheeves, becauſe there are generally ſome Hindrances more than the common Friction; as for example, when Sheeves rub againſt the Sides or Cheeks of the Blocks, or when their Hole wears bigger, which encreaſes the Friction as much as if the Pin was become ſo much bigger.

BEING willing to try how far the Theory of Friction and bending Ropes would agree with ſuch Tackles of Pulleys as are commonly uſ'd in Building, and conſequently be uſeful to direct our Practice, I made the two following Experiments.

EXPERIMENT I.

I took a Tackle of five Braſs Sheeves in Iron Frames or Blocks; that is, three Sheeves in the upper Block and two in the lower. Having made an *Equilibrium* by hanging one hundred and a quarter at the lower Block, and a quarter of an hundred at the running Rope, I added 17 Pounds and an half before the Power could go down and raiſe the Weight.

EXPERIMENT II.

Two hundred and an half being balanc'd by half an hundred, the Addition of 28 Pounds made the Power raiſe the Weight.

N.B. *The Sheeves were 5 Inches Diameter, the Pins half an Inch, and the Rope three quarters of an Inch.*

IN the firſt Experiment 17 lb and an half exceeds by $4\frac{1}{2}$ lb the Sum of the Frictions and Reſiſtances deduc'd from the Theory.
But

But in the second Experiment 28 lb exceeds the Sum of the Friction, &c. not quite 1 lb. The Reason of this appear'd to be, that the Rope at first was too big for the Cheeks that held the Sheeves; but in the second Experiment, where the Rope was more stretch'd, its Diameter became something diminish'd, and so brought off the Rope from rubbing so hard against the Cheeks. Lect. IV.

FROM knowing the Quantity of Friction in such large Tackles, we may know what to expect in Practice. For if one Man, who for a small time can exert the Force of 100 lb, thinks that he may draw up a Stone or a Roll of Sheet Lead, or any other such Weight to the top of an House with a *Tackle of Five* (because this would seem feasible from Mechanical Principles) will find himself mistaken on account of the Friction, which will not be surmounted without an additional Force of 50 lb.

I hope this Account which I have given of Friction and Hindrance to Motion in Mechanical Engines (however imperfect it is) may be of considerable Use to direct such Persons as concern themselves with Engines and Manufactures. And to afford all the Help I can upon this Subject, I will give him some Considerations of the comparative Strength of Men and Horses, as well as the best way of applying their Forces, being the Result of many Years Observations of my own, as well as what I have found in Authors who have treated of these Things.

AN Horse draws with the greatest Advantage, as we have already shewn from Mr. CAMUS, when the Line of Direction (being parallel to the Plane on which the Weight moves) is level with the Horse's Breast, and is able in such a Situation to draw 20 lb eight Hours a Day, and walking about two Miles and an half an Hour, which is about three Foot and an half in a Second. And if the same Horse is made to draw 240 lb he can work but six Hours a Day, and cannot go quite so fast; and in both Cases, if he carries some Weight, he will draw better than if he carried none. We don't mean by this what an Horse can draw upon a Carriage; because in that Case Friction is only to be overcome, so that a middling Horse well applied to a Cart, will often draw above 1000 lb; but so much as an Horse could draw up out of a Well over a single Pulley or Roller (made to have as little Friction as possible) is properly what an Horse can draw; and Horses, one with another,

Le^{ct}. IV. draw about 200 lb in such a Case as we said before. To this may be referr'd the working of Horses in all sorts of Mills and Water-works, where we ought to know as near as we can, how much we make every Horse draw, that we may judge of what the Effect will be when proper Allowance shall have been made for all the Frictions and Hindrances, before we cause any Machine or Mill to be erected.

WHEN an Horse draws in a Mill, Water-work, or Gin of any Kind (in which the Horse is made Use of to draw round a Capstane or *Axis in Peritrochio*) great Care should be taken that the Horse-Walk be large enough in Diameter, otherwise the Horse cannot exert all his Force as he goes round; for in a small Circle or Horse-Walk, the Tangent (in which the Horse should draw) deviates more from the Circle in which the Horse is oblig'd to go, than it does in a great Circle. The Horse-Walk should not be less than 40 Foot in Diameter, when ever there is Room for it; and the same Horse loses of his Force considerably in a small Walk, because he pulls in a Chord of the Circle, drawing the horizontal Beam behind him at acute Angles, so much, that in a Walk of 19 Foot Diameter, I have known an Horse lose 2 fifths of the Force that he exerted in a 40 Foot Walk. Most of the Mill-wrights in *London* (and I believe in most great Cities) do not love to make large Horse-Walks, even when they have Room; because, as there is generally want of Room where they have been oblig'd to set up Works, they have accustom'd themselves to make their Geers for small Horse-Walks, and think it enough to give the same proportional Velocity to the Power and Weight as is done in a larger Horse-walk (because if the Cog-wheel be so much less in the Diameter as the Horse draws nearer to the Center, the difficulty of drawing, were it not for the twisting of the Horse, would always be the same) not considering the Strain put upon the Horse; or when by Practice they have found how much a Horse may easily draw, with all the Disadvantages which the sudden turning gives him, they won't take the Advantage which more Room might give in removing that Difficulty, because they don't care to go out of the Way which they have been accustomed to. But such Mill-wrights, as have work'd at Coal-pits and Mines know better, as they have been us'd to large Horse-walks in Coal Fields, &c.

I HAVE

I HAVE often found that five Men are equal in Strength to one Leet. IV. Horſe, * and can with the ſame Eaſe push round the Horizontal Beam in a 4 Foot Walk; but three of the ſame Men will push * Ann. 6. round a Beam in a 19 Foot Walk, which an Horſe (otherwiſe equal to five Men) can but draw round.

THE worſt way of applying the Force of a Horſe is to make him carry or draw up Hill; for if the Hill be ſteep, three Men will do more than a Horſe, each Man will climb up faſter carrying 100 lb Weight, than a Horſe that is loaded with 300 lb. This is owing to the Poſition of the Parts of a Man's Body, which are better adapted to climb than thoſe of a Horſe.

IT follows from this Obſervation, that thoſe who have thought to gain great Advantage from the Weight of a Horſe by applying it to an Engine to work the Forcers of Pumps, have not in the Execution found what they expected from a Calculation of the Weight of that Animal, becauſe at every Step the Horſe is really climbing up Hill.

As a Horſe from the Structure of his Body can exert moſt Force in drawing horizontally in a ſtrait Line, a Man can exert leaſt Force that way; as for Example, if a Man weighing 140 lb walking by a River or Canal ſide, draws along a Boat or Barge by means of a Rope coming over his Shoulders, or any how faſtened to his Body, he cannot draw above 27 lb, or about the ſeventh part only of what a Horſe can draw in that Caſe; for the whole Force that a Man exerts in that Action intirely depends upon his Weight, and not his whole Weight neither, only about $\frac{2}{14}$ parts of his Weight, acting obliquely too, pushing him forwards as he ſtoops, produce the whole Force whereby the Man draws the Barge along, as has been demonſtrated by Mr. *De la Hire*, in a *Memoire* which he preſented to the *Royal Academy of Sciences at Paris*, in the Year 1699; of which I have given a Tranſlation in my Notes. †† Ann. 8.

HIS other Reasoning about the Application of the Force of a Man is juſt; but his *Data* not being true, ſome of his Concluſions tho' truly drawn from his *Data*, are not true in Faet; therefore I have in the ſame Note given Remarks upon what he has ſaid.

Lect. IV.



IN drawing a Barge in the manner above-mentioned, a heavy Man (provided he be not unwieldy) will do more than another, unless he carried Weight proportionably, and the higher the Weight is carried, the better.

WHEN a Man turns an horizontal Roller or Windlass by a Handle or Winch, he should not have above 30 lb Weight acting against him, if he is to work ten Hours a Day, and raise the Weight about 3 Foot and an half in a Second, which is the common Velocity that a Horse draws with. I say 30 lb, supposing the Semidiameter of the Windlass equal to the Distance from the Center to the Elbow of the Handle; for if there be a Mechanical Advantage, as there usually is, by having the Diameter of the Axle, on which the Rope winds, four or five times less than the Diameter of the Circle describ'd by the Hand, then may the Weight (taking in also the Resistance on account of the Friction and Stiffness of the Rope) be four or five times greater than 30 lb, that is, so much as it rises slower than the Hand moves.

IN this Operation the Effect of a Man's Force varies in every part of the Circle describ'd by the Handle. The greatest Force is when a Man pulls the Handle upwards from about the Height of his Knees, and the least Force when (the Handle being at top) a Man thrusts from him horizontally; then again, the Effect becomes greater as a Man lays on his Weight to push down the Handle; but that Action cannot be so great as when a Man pulls up, because he can lay on no more than the whole Weight of his Body, whereas in pulling he can exert his whole Strength: Lastly, a Man has but small Force to pull the Handle horizontally towards him when it is at lowest. Let us, as *Mons. de la Hire* does, suppose a Man of moderate Strength; to weigh 140 lb, he may in the four principal Places of pushing and pulling in the whole Circumference of Motion exert the following Forces, *viz.* in the strongest Point a Force equal to 160 lb; in the weakest, a Force equal to 27 lb; in the next strong Point, 130 lb; and in the last or second weak Point, 30 lb. Let us add all these Forces together, which will make 347, and divide them by 4, and we shall have 86 lb $\frac{3}{4}$, and this gives us the Weight that a Man might lift by a Winch, if he could exert his whole Force continually without stopping to take Breath; but as that cannot be, the Weight must return and over-power at the first

first weak Point, especially when the Handle moves slowly, as it must if a Man was to exert his whole Strength all round. Besides, for raising such a Weight we must suppose the Man to act always along the Tangent of the Circle of the Motion, which does not happen in the Operation. Then there must be a sufficient Velocity given, * that the Force applied at the strong Points may not be spent before the Hand comes to the weak ones, so that it is difficult for a Man to continue that irregular Motion; and therefore when there are no other Advanges, the Resistance ought to be but 30 lb; and even that could not be supported at the weakest Point, were it not for the Force remaining from the strong Point.

Lect. IV.

* Ann 9.

IF two Men work at the End of a Roller or Windlass to draw up Coals or Ore from a Mine, or Water out of a Well, they may more easily draw up 70 lb (still supposing the Weight and Power to have equal Velocities) than one Man can 30 lb, provided the Elbow of one of the Handles be at right Angles to the other; for then one Man will act at the strong Point, when the other acts at the weak Point of his Revolution; by which means the two Men will mutually and successively help one another. The common way is to put on the Handles opposite to one another, which cannot give the Advantage above-mentioned; tho' there is some little Force gain'd even in that Position, because one Man pulling while the other thrusts, works at the strongest of the two weak Points, whilst the other works at the weakest, and so helps him a little.

THERE is indeed a Way to make a Man do a third part more Work with a Windlass when the Motion is pretty quick, as about 4 or 5 Foot in a second, † and that is by the Application of a Fly, † Ann. 9. which is a Cross with Leaden Weights at its Ends, or rather (what is much better) a heavy Wheel at right Angles to the Axis of the Windlass or Roller. By this means, the Force of the Power, which the Man would lose, is kept in the Fly and equally distributed in all the Parts of the Revolution; so that for a little while a Man may act with the Force of 80 lb, that is, overcome a continual Resistance of 80 lb; and work a whole Day when the Resistance is but 40 lb. ††

WHEN

†† The Fly may be applied to several Sorts of Engines whether mov'd by Men, Horses, Wind or Water, or any animate or inanimate Power; and is of great Use in those Parts of an Engine which

Lect. IV.

WHEN a Man carries a Weight or a Burthen upon his Back, he exerts a great Force very effectually, many Muscles at once being employ'd in that Operation; the Muscles of his Neck, Back and Loyns keep his Body and Head in the proper Position to sustain the Weight; those of his Shoulders and Arms help to keep it in its Place; and the Muscles of the Legs and Thighs raise the Weights of all the Body and Burthen as the Man walks along. In this way of working, three Men do much more than an Horse, and two oftentimes do as much, nay even more, as may be observ'd in the daily Labour of the *London* Porters. * A Porter will carry 200 lb, and walk at the rate of three Miles an Hour: A Coal-heaver or Porter that carries Coals, will carry 250 lb, but then he does not go very far before he lays down his Burthen; tho' on the other hand he will often go up stairs with that Weight: Chairmen do not act with all the very same Muscles as Porters, but as they have Straps brought down from their Shoulders to the Poles of the Chair, the Muscles of the Loins and Back are concern'd, and likewise the Extensors of the Legs and Thighs: Two of them will walk very fast with 300 lb (that is 150 lb each) at least at the rate of four Miles *per* Hour. Whereas a Carrier's Horse,

* Ann. 10.

which have a quick circular Motion, and where the Power or the Resistance act unequally in the different parts of a Revolution. This has made some People fancy that the Fly adds a new Power; supposing that a Fly join'd to an Engine that is to move round does help to carry it about. But tho' it may be said in some way to facilitate the Motion, by reason that it makes it more uniform and equal, yet upon the whole it causes a Loss of Power, and not Increase. For first, It requires a continual Supply of Power to put the Fly in Motion to a certain Degree of Velocity, and to keep and maintain it in that Velocity; for that the Fly has no Motion of its own, but what it receives from the impress'd Force. Secondly, The rubbing and wearing of the Pivots or Gudgeons of the Axis do still hinder and lose the impress'd Motion: And thirdly, The Air thro' which the Weights at the Ends of the Fly do move, do also hinder the Motion thro' it (tho' less when the Fly is circular) and both these Impediments together, if the Fly be not still supplied with new Power, will make it stand still and be at Rest.

So that the Fly can of itself add no new Power to the Motion of the Engine to which it is applied, more than what is receiv'd from the first Mover that impress'd the Motion on it, but loses even some of the first Motion.

But the reason how it becomes convenient and useful in many Engines (as we have shewn in the Windlafs or horizontal Axis in Peritrochio) is this——That, whereas either the Powers exerted by the Engine are intermitted or unequal, and so the Motion is more difficult in one part of the Revolution than another, or perhaps the Strength of the Man, or any other Power to be supplied, cannot be so well applied to one part of the Revolution as it can to another: In these Cases the Fly becomes a Moderator, and makes the Motion of Revolution almost every where equal, tho' the Resistances are unequal, and the Forces impress'd are unequal; for that it has accumulated in itself a great Degree of Power, which it equally and gradually exerts, and as equally and gradually receives; whence making the Revolution in all parts pretty near uniform, it becomes more pleasant, easy, and convenient to be acted and mov'd by the impelling Force; which is the whole Benefit which is procur'd by this Mechanical Engine, this way applied. But I shall speak of the Fly and some other of its Uses in another Place.

Horse, that goes but about 2 Miles *per* Hour, carries only 224 lb; Lect. IV. or sometimes, when the Roads are very good, and the Horses strong, * 270 lb.

* Ann. II.

Mr. *Richard Newsham* Engineer of *Clotb-Fair* near *Smithfield*, has contriv'd his Engines to put out Fires in such a manner, that part of the Men that work them exert their Force by treading, which is more effectual than any other way that Men can work at such Engines, the whole Weight of the Body being successively thrown on the Forcers of the Pumps; and even part of a Man's Strength may be added to the Weight by means of horizontal Pieces to which he can apply his Hands when he is treading: Whereas, by applying the Hands to move Leavers or turn Winches, the Power must act very unequally: This is the Reason why with the same Number of Men he has generally thrown Water farther, higher, and in greater Quantities, with the same siz'd Engines, than other Engineers who have try'd their Engines against his.

N.B. *His Engines have several Conveniencies peculiar to them, which makes them preferable to all others that I ever saw for extinguishing accidental Fires; but I reserve their Description for another part of my Book.*

THE last and most effectual way of a Man's working, is the Action of *rowing*; wherein a Man acts with more Muscles at once for overcoming the Resistance, than in any other Position; and as he pulls backwards the Weight of his Body assists by way of Leaver.

C O R O L L A R Y.

FROM the Consideration of the several Ways of a Man's acting in the way of Labour, compar'd with the last, we may see how much People are mistaken who think to row a Galley, Boat, or Barge, by vertical Oars fix'd on a horizontal Axis like a Mill-Wheel, the Men working this Machine by *beaving* at a *Capstane*; or turning Winches within the Vessel. For this will always be bringing the Men from easier and more effectual, to harder and less advantageous Work; as has been found by a great many more Engineers than will own it, and will be found by all that shall ever try it, be the Machine made in any Shape whatever, unless when the Men work in a rowing Posture.

A PRO-

Lect. IV. A PRODIGIOUS Force may be exerted by the Muscles of the
~ Legs and Thighs to raise an immense Weight a small Height; but
as that Operation cannot be continued and brought to daily Labour, I refer the curious Reader to the Notes where that is more
* Ann. 7. fully explain'd: * neither do I here take notice of digging, hammering, cleaving of Wood, or any of the laborious Operations of Handy-craft Trades; because some Men are much more dextrous than others; and the same Man by long Use becomes so perfect in one way of working, that by an acquir'd Sleight he shall do twice the Work that an unexperienc'd Person can do, and yet not employ half so much Strength. But this is properly *Craft* and not *Labour*, which last was all I meant to consider here.

ANNOTATIONS on Lecture IV.

1. **P**AGE 184. — [*The Friction is equal to about one third of the Weight, &c.*] There are some Cases wherein the Friction does not come up to a third Part of the Weight of the Body which rubs; but as in most Cases it does, I chose to give that Proportion as the Foundation of the Calculation of the Friction of an Engine compos'd of many Parts before we erect it, especially in regard to the Manufactures, because the wearing of the Parts of Machines will in time alter their Figure and encrease the Friction. And it will be better in Practice to find the Friction to be less than in the Theory, tho' that seldom happens, when calculated from a third of the Weight. Annotat. Lect. IV.

2. [Page 186. — *It would require one third of the Weight of the Sledge, &c.*] In the Table of the Frictions of Sledges (Page 193, 194 and 195) quoted from *Monf. Camus's* Experiments made on small Models, there are more Cases wherein the Friction is less, than where it is more, than one third of the Weight; but it is to be observ'd, that in all his Experiments the Sledge is in Motion; but as I mention'd in my last Note, I rather chose to continue the Proportion of a third on account of Accidents, beginning to draw from Rest, [and meeting with Rubs and Unevenness in the Streets, &c.]

3. [Page 201. — *We cannot expect that any Carriage to bear Weight can have so little Friction, &c. — the manner of remedying Friction, &c.*] If the Axels in some Carriages being made of Iron, and in others only clouted (that is cover'd) with Iron, were to run in brass Boxes fix'd in the Naves of the Wheels, they would go so much easier, and wear so much longer, without danger of firing, as to make full Amends for the extraordinary Expence. Where People are curious and don't value Cost, as in Chaises and some Chariots, an Iron Axel being less in Diameter will have less Friction in proportion as it is smaller, and last very long if it turns in Boxes of cast Brass; but the Gudgeon in the Box must be sufficiently long, which it may be without encreasing the Friction, as has been already prov'd, but will be more clearly evinc'd by Experiments made on a Machine which I shall describe in this Annotation. If the Gudgeons are but two or three
K k Inches

Annotat. Inches long as some have made them, thinking thereby to diminish the Friction, they will only wear two or three times faster than if they had been made four or six Inches long.

A Wheel of an Engine whose Axis is fix'd to, and turns with, the Wheel, may have the Friction of its Gudgeons diminish'd in any Proportion. As for Example, under the Iron Gudgeon G g, † which we will suppose here of an Inch in Diameter, let there be two brass Rollers A B of eight Inches Diameter each, whose Axes C, D, are horizontal and parallel to the Axis of the Wheel: the Rollers which consequently are vertical (as may be seen in *Fig. 7.* †) are about an Inch thick (or more if you will) and not in the same Plane, but one a little before the other, and parallel to it. Upon these Rollers is supported and turns the Gudgeon G g of the great Wheel. *In this Case the Friction of the Gudgeon becomes eight times less than if it mov'd in the common Brasses;* for if we suppose the Gudgeon to move in the Direction G g, it will not quit the Part g of the Wheel A on which it bears, to go to another bearing with its touching Part, as happens when it turns and makes a Friction in the common way, but it brings along with it the Circumference of the Wheel or Roller A, turning it about in the Direction A g, whilst its other touching Part G in its turning does also carry round the Roller B in the Direction G B, and these Rollers wou'd follow that Motion without any Friction, were it not for the bearing of their own Iron Axes C, D, in their Brasses. So that the Friction is transferr'd from the Gudgeon G g to the Axes C, D, where the Velocity of the rubbing Parts being eight times slower than it wou'd have been at G g, the Friction must be eight times less, as we have † shewn already. Now tho' these Rollers have four bearing Points, and the other two which support the Gudgeon of the great Wheel's Axis have also four bearing Points, the Friction is no greater, than if there was but one bearing, because each of these Points sustains but an eighth part of the Weight. *Therefore the Friction is made eight times less by means of these Rollers: which was to be prov'd.*

† Page 185,
186.

S C H O L I U M.

If the Axes of the Rollers are made but of half the Diameter of the Gudgeon, as at E and F, then the Friction will be 16 times less; and yet they will be sufficiently strong; for as a Cylinder of half an Inch Diameter is equal in Strength to the fourth part of one of an Inch, the four Ends of the Axis of two Rollers at one End will be equal in Strength to the Gudgeon, and so at the other End.

C O R O L L A R Y.

Hence follows, that if the Ends of the Axes of the Rollers were each of them supported by two other such Rollers, the Friction would again be diminished 16 times more. Let us for Example suppose a Wheel to be fix

Foot

Foot in Diameter and to weigh 648 lb, the third of that Weight for Annotat. Friction is 216 lb, which must be divided by 72, because the Diameter of the Gudgeon is so many times contained in the Diameter of the Wheel, and we shall have 3 lb for the Friction of the Wheel upon common Brasses; but this Number must be divided by 16 times 16, that is, by 256, on account of the Rollers bearing on Rollers, which will thus reduce the Friction to $\frac{3}{256}$ lb, or little more than one 85th part of a Pound, or 3 Drains of 16 to the Ounce.

4. [Page 217. —*the Friction on the Axel, &c.*] Since, as we have said and already prov'd *, the Friction arises from the Weight that presses the Parts together, and not from the Number of Parts that touch; there is no Occasion for shortening the Axes of Wheels, either in Carriages or any other kind of Wheels in order to diminish the Friction; for that will not only fail of the desir'd End, but make the rubbing Axis wear out much faster; and in Clock-work it is of bad Consequence, because when the Holes are counter-sunk to shorten the bearing, the Holes soon wear too big, whereas, if they were made only cylindrical the thickness of the Frame-Plate, the Friction wou'd be no more, and the Pivots wou'd wear much longer; and indeed it is now what all good Clock-makers practise; for if they counter-sink, it is but a small Depth to hold Oil. Experiments on the following Machine will make this more sensible and evident.

Plate 18. Fig. 8.

On the flat brass Plate A B C (which is here represented almost as big as the Machine) are fix'd two upright Plates D and E, with a Slit in one of them at D, and a Hole in the other between the Letters K and L, to receive the small Pivots at the Ends of the Axis D K of the Wheels Z F L G. But those Pivots, which are but about one 30th of an Inch in Diameter, do not bear on the Hole at K and the bottom of the Slit at D; for they are supported upon two circular Plates or Rollers at each End of the Axis, viz. the vertical Plates M 1, M 2, M 3, M 4, in the manner describ'd in the last Note, and represented at G g Fig. 6. † and the Secti-
on of one of the Rollers or Plates is shewn by Fig. 7. So that when the Wheel turns round one way, all the rolling Plates turn the contrary way as freely as if the Pivot had been a Pinion, and the Wheels or Plates had been tooth'd, because the Axes of the Plates have very small Pivots which turn in very smooth Holes, that are made and polish'd in the four upright fix'd Plates or Cocks N 1, N 2, O and P, which last Cock has only the Corner of its Basis visible in the Figure, the Wheel M 2 hiding it in this Situation of the Machine. The little Cocks, such as D, d, d, serve for the Ends of the Pivots to bear against (both for the great Wheel and the four little Wheels, there being ten of them of which only three are visible here) and by that Resistance against the Ends, the Shoulders as C C, Fig. 7. * * Pl. 18 F. 7. will never rub at all.

K k 2

By

Annotat. By this means the great Wheel has so little Friction, that if we apply
 Left. IV. a Finger to its Circumference to put it briskly in Motion, any Point of
 its Circumference will go more than the space of a Mile before the Wheel
 stops; for tho' one cannot count the Number of Revolutions of the great
 Wheel by looking at it, yet one may know their Number by looking at
 the Holes which are left in the small Wheels (one in each) for that Pur-
 pose; for they being two Inches Diameter turn but once whilst the great
 Wheel (whose Pivots gives them Motion) turns sixty times.

The Friction of the great Wheel thus becoming so little, as to be in a
 manner insensible, it is fit for the Purpose intended. Then on the top of
 the Cock or upright Piece Q R, screw'd down fast by its Base or Foot Q,
 is fasten'd by a Screw at R one End of the Spiral Spring S₁, S₂, S₃, S₄,
 whose other End is fasten'd to the Axis of the great Wheel near S₄.
 Now if the Wheel be made to turn on its Pivots, by bringing the Point Z
 of its Circumference towards the Index Y, which points to Degrees on
 the Edge of the Wheel (but not represented here except by the Dots
 mark'd from Z towards Y) as soon as we let it go it will return towards Z
 and make several Vibrations backwards and forwards, like the Balance of a
 Watch, for a long time; but if any thing bears against the Axis H I D,
 which is truly cylindrical and one 4th of an Inch Diameter, then there will
 be fewer Vibrations in proportion to that Friction. Now to shew that the
 Friction is proportional to the Weight that presses on the Axis, and not the
 Surface, the following Experiments are made.

EXPERIMENT I.

- † Pl. 18. F. 9. Take the Piece of *Figure 9* † weighing half an Ounce (being made in
 the Form of a flat Cross, and after filing, ground smooth upon an Oil-stone
 on the flat Under-side of V, with a little Hole at the End T, and a small so-
 lid round Piece hanging down at the opposite End X) and lay it over the
 Axis of the Wheel betwixt I and D, so that the Weight of the hanging
 End X, drawing down the flat V upon the said Axis, makes the End T rise
 with its little Hole against the Point or lower End of the Screw T of the
 Cock † T, which keeps it in its Place when the Axis turns round under it;
 then having brought Z to Y, or drawn back that Point against the Bent of
 the Spring about 90 Degrees, observe the Number of Vibrations that the
 Wheel makes before it stands still by the Friction of the Cross in the Situa-
 * Pl. 18. F. 8. tion represented by faint Lines, at T V X *Fig. 8.* * Suppose the Num-
 † Pl. 18. F. 10. ber of the Vibrations be 50; then the Cross of *Fig. 10.* † be put on in
 the same manner and at the same time just by the other, and kept in its
 Place by the bottom of the other Screw †; as it is exactly the same in Fi-
 gure and Weight as the first Cross, it will add as much more to the Friction
 of the Axis, the Weight pressing as well as the Surface rubbing being dou-
 bled; which appears by giving Motion to the Wheel as before, because
 then the Number of Vibrations will be but 25.

E X P E-

EXPERIMENT II.

Annotat.

Instead of the two Crosses abovementioned, put on the Cross of *Fig. 11*, *Lect. IV.*
 * which weighs one Ounce (that is as much as the two others) but has **Pl. 18. F. 11.*
 the Surface under V exactly smooth'd and polish'd like the others: Then
 put the Wheel in Motion as before, and the Vibrations will be but 25 in
 Number, tho' the Surface rubbing is but half, because the Weight is the
 same. And this is further prov'd by

EXPERIMENT III.

In which, the Piece X must be unscrew'd from its Place and screw'd a-
 gain in the same Hole on the other side of the Cross to bring the little Flat
 of the Prim at V to bear upon the Axis of the great Wheel instead of the
 broad Part of the Cross. Then also will the Wheel lose its Motion after
 25 Vibrations, tho' the bearing Surface be above twenty times less than
 when both the first Crosses were on; because the Weight is the same.

5. [Page 232. — *shou'd alter the manner of working, &c. — instead of
 gaining an Advantage he must call in more Men to perform the Work.*

Some have endeavour'd to render this Engine more useful by causing it
 to roll upon an inclin'd Plane, instead of making it rise directly up in the
 manner describ'd, and condemn'd in my Account of it. I thought proper
 to shew here, what must be the Loss of the Power in proportion to the
 Inclination of the Plane.

I say therefore, *That in every Inclination of the Plane, if the Sine of
 the Angle of Inclination be taken in Parts of the Radius of the Axel, or
 Roller, The Power will be to the Weight :: as the Radius of the Roller +
 the Sine of Inclination : to the Radius of the Wheel, — the said Sine of
 Inclination; that is, in the Fig. P (= 1) : W (= 3) :: dk : a k. † Plate † Pl. 19. F. 1.*
 19. Fig. 1.

In the present Experiment BE is an inclin'd Plane, on which the Roll-
 er C is to roll up, touching the said Plane at the Point c; AM is the
 Wheel behind that Plane; another such Plane, and equally inclin'd, being
 also suppos'd behind the Wheel, to support the other End of the Roller.

The Lines of Direction of the Power and Weight being *aP* and *dW*;
 through the Point of Contact or Center of Motion *c*, draw *AD* parallel
 to the Horizon, and perpendicular to *aP* and *dW*; through the Center
 of the Engine *C*, draw *ad* parallel to *AD*. Suppose the Angle *BcA* of
 the Plane's Inclination to be 30°, the right Sine will then be equal to half
 the Radius; therefore dividing *C2* (the Radius of the Roller) into two
 equal parts at *k*, if you draw *kc* and *Cc*, the Angle *kcc* will be equal
 to *BcA*, and its Sine will be *Ck*. Now since it is evidently the same
 thing to make use of *ad* for a Lever, whose Center of Motion is at *k*,
 as of *AD* equal and parallel to it with its Center of Motion at *c*: it
 follows, that, in this Inclination of the Plane, the Distance of the Weight
dk is greater than *dC* (the Distance of the Weight in the common Use of
 this

Annotat. this Engine) by the Addition of this Quantity Ck , the Sine of the Angle
 Lect. IV. of Inclination; and ka , the Distance of the Power is less than Ca (the
 Distance of the Power in the common way) by the Subtraction of the
 said Quantity or Sine Ck : consequently that on an inclin'd Plane; the Pow-
 er is to the Weight $::$ as Dc : to cA . Q. E. D.

C O R O L L A R Y I.

Hence it follows, that the Radius of the Wheel, and the Radius of the Roller being given, the Loss of Power may be found in any Inclination of the Plane. Thus, as here the Power, which in the common way wou'd be but one 5th of the Weight, must be one 3d part of it: So if the Angle of the Plane's Inclination was but $11^{\circ} 32'$, the Power wou'd be one 4th of the Weight, &c.

C O R O L L A R Y II.

Hence follows also, that if the Plane BE be horizontal, no Force of the Power will be lost, because $cg :: CG : CF$.


S C H O L I U M.

As the Friction of the winding of the Ropes, such as Bc in the new Way, is greater than the Friction of the Pivot in the old Way (besides the Friction of the Collars of the Counterpoise to the Engine) so that Friction diminishes, as the Ropes bear less Weight, according to the Diminution of the Angle of the Plane; and when the Plane is horizontal, and without a Counterpoise, even then the winding up of the Ropes, and Pressure of the Roller against the Plane, is equal to the Friction in the common way.

N. B. *I made the Experiment with Pivots twelve times less in Diameter than the Roller, and fine pliable Silk instead of Ropes.*

6. [Page 243. — *Five Men are equal in Strength to one Horse, &c.*] Such *English* Authors as have compar'd together the Strength of Men and Horses at a Capstane have found their Forces to be in that proportion, as Sir *Jonas Moore* and others, &c. But the *French* Authors always make an Horse equal to seven Men; which I believe to be according to their Observations. Nay, and I have observ'd the Labourers in *Holland* (one with another) to work with a Force pretty near in that Proportion. So that we may say, — That five *English* Labourers are equal to an Horse, and only seven *French* Men or as many *Dutch* Men. But here we do not at all consider Skill, or Sleight, which make a Man of the same Strength do much more than another. But in *Turkey* the Porters will carry twice more than the strongest *English* Porters, as we shall more particularly consider in the next Note.

7. [Page 248. — *a prodigious Force may be exerted, &c.*] About 30 Years ago one *Joyce*, a *Kentish* Man, famous for his great Strength (tho' not quite so strong as the King of *Poland*, by the Accounts we have of that Prince) shew'd several Feats in *London* and the Country, which so much surpriz'd the Spectators, that he was by most People call'd the second *Sampson*; but tho' the Postures which he had learn'd to put his Body into, and found out by Practice without any mechanical Theory, were such as would make a Man of common Strength do such Feats as would appear surprizing to every body that did not know the Advantage of those Positions of the Body; yet no body then attempted to draw against Horses, or raise great Weights, or to do any other thing in Imitation of him; because, as he was very strong in the Arms, and grasp'd those that try'd his Strength that way so hard, that they were oblig'd immediately to desire him to desist, his other Feats (wherein his manner of acting was chiefly owing to the mechanical Advantage gain'd by the Position of his Body) were intirely attributed to his extraordinary Strength.

Annotat.
Lect. IV.


But when he had been gone out of *England*, or had ceased to shew his Performances, for eight or ten Years; Men of ordinary Strength found out the Way of making such Advantage of the same Postures as *Joyce* had put himself into, as to pass for Men of more than common Strength, by drawing against Horses, breaking Ropes, lifting vast Weights, &c. (tho' they cou'd in none of the Postures really perform so much as *Joyce*; yet they did enough to amaze and amuse and get a great deal of Money) so that every two or three Years we had a new *second Sampson*.

About 15 Years ago a *German* of middle Size, and but ordinary Strength, shew'd himself at the *Blue Posts* in the *Hay Market*, and by the Contrivances abovementioned, pass'd for a Man of uncommon Strength, and got considerable Sums of Money by the daily Concourse of Spectators. After having seen him once, I guess'd at his manner of imposing upon the Multitude; and being resolved to be fully satisfied in the Matter, I took four very curious Persons with me to see him again, viz. the Lord Marquis of *Tullibardin*, Dr. *Alexander Stuart*, Dr. *Pringle*, and a mechanical Workman, who us'd to assist me in my Courses of Experiments. We plac'd our selves in such manner round the Operator as to be able to observe nicely all that he did, and found it so practicable, that we perform'd several of his Feats that Evening by our selves, and afterwards I did most of the rest as soon as I had a Frame made to fit in to draw, and another to stand in and lift great Weights, together with a proper Girdle and Hooks. I likewise shew'd some of the Experiments before the *Royal Society*; and ever since, at my Experimental Lectures, I explain the Reason of such Performances, and take any Person of ordinary Strength that has a mind to try, who can easily do all that the *German* abovemention'd us'd to do, without any Danger or extraordinary straining, by making use of my Apparatus for that Purpose. I don't hear that any of these *Sampsons* have attempted since to impose upon People in the same manner in or near *London*.

But

Annotat. But now it will not be improper to mention what were the Feats of
 Lect. IV. Strength which the *German* us'd to perform (for I never saw *Joyce*) and to
 shew, from the Make of the human Body, how easily any one may do the
 same.

1st, The strong Man IHL on the Frame AGBEFC D sat upon an
 * Pl. 19. F. 1. horizontal Board, or rather inclining backwards as FG, * with his Feet
 against an upright immoveable Prop as DCF counterbrac'd at E, with a
 strong Girdle H round him a little below his Hips; to the Iron Rings of
 which Girdle was fasten'd a Rope by means of a Hook. The Rope went
 between his Legs and thro' a Slit of the Prop at L, and several Men at
 MN, or two Horses could not by their pulling, move him out of his
 Place.

N. B. His Hands at K seem'd to pull, but were of no Advantage to him:
 nay had he lifted the Rope ever so little with them, it would have been a
 Disadvantage: and tho' the Board on which he sat was (when I saw
 him) in the horizontal Position LP, it is much better and less dangerous
 to have it inclin'd as FG, and only an Hole at L for the Rope to run
 thro' instead of an opening from L up to D, as I shall shew in explaining
 this Operation.

† Pl. 19. F. 3. 2^{dly}, The same Man IHL † having fix'd the Rope abovementioned
 round a strong Post at R, and then pass'd it thro' a fix'd Iron Eye at L,
 fix'd it to his Girdle, and setting his Feet against the Post near the said Eye
 rais'd himself from the Ground by the said Rope, which he broke by
 suddenly stretching out his Legs and fell backwards on a Feather Bed at
 B laid on the Ground to catch him that his Fall might not hurt him.

* Pl. 19. F. 4. 3^{dly}, He lay down on the Ground in the Posture IHL * with an An-
 vil KH on his Breast at H, upon which another Man M hammer'd with
 all his Force the Iron K with a Sledge Hammer; and sometimes two
 Smiths cut a great cold Iron Bar in two with Chizzels. Sometimes a
 great Stone of which the half is seen at S was laid upon his Belly and
 broke with a Blow of the great Hammer. But he had the Stone broke
 upon his Belly in the Posture of Fig. 5. which is much less dangerous
 when nothing is under the Back, than when a Man lies upon solid Ground,
 as we shall shew.

4^{thly}, The pretended *Sampson* puts his Shoulders (not his Head as he
 us'd to give out) upon one Chair and his Heels upon another, and sup-
 ports one or two Men standing upon his Belly, raising them up and down
 as he breathes, making with his Back-bone and Thighs and Legs the Arch

† Pl. 19. F. 5. IHL, † whose Abutments are at I and L.

N. B. The Stone of one and a half Foot long, one Foot broad, and five or
 six Inches thick, is laid on at H when it is to be broken by a Blow of a
 Hammer.

* Pl. 19. F. 6. 5^{thly}, He lies down on the Ground in the Posture IHL *, and the
 Man M standing upon his Knees, he draws his Heels towards
 his Breech and so raises his Knees over C, and lifts up the Man gradually,
 till having brought perpendicularly under the Man as in Fig. 7. he raises
 his

his own Body up, and putting his Arms about the Man's Legs rises with Annotat. him and sets him down on some low Table or Eminence about the Height Le^{ft}. IV. of his Knees: and this he sometimes does with two Men; which is no difficult Performance.

6thly, He stands in the Frame ABCDEF, * and pretends to raise up, * Pl. 20. F. 2. but does really sustain, a Cannon C laid on the Scale SS, the Ropes of the Scale being fix'd to a Rope or Chain LH, hanging at his Girdle H, his Assistant knocking away the Rollers R, r, from under the Scale when once he has fix'd himself so as to have his Ropes tight and his Legs and Thighs quite streight.

N.B. *It is very near as easy to break the Rope with the Eye L fix'd into the Ground or Floor by means of the Girdle H, † Fig. 1. as in † Pl. 20. F. 1. the manner represented in the * third Fig. of Plate 19. But he never * Pl. 19. F. 3. tried it that way; because it is so obvious that several People would immediately have tried it too, and would find that there is no difficulty in breaking the Rope thus, as I have often done it; but by making the Fall backwards seem necessary in the Operation, few Persons care to try it his way.*

The † 4th Figure represents the Girdle made of strong double Horse Girt, † Pl. 20. F. 3, with strong Iron Loops at G and R. The Hook is seen at Fig. 5; and 4, and 5. the Position of the Iron Eye at Fig. 3, where you may observe that the Edge of the Eye, and not the open part, is towards the Post, so that the Rope does not easily slip thro' the Hole, but jams or stops in it, whereby the whole Strength of the Man's Effort acts upon one part of the Rope, and so it is easily broken.

The Man likewise us'd to take a flat Piece of Iron of the Figure mark'd 7, * and twist it into a Screw. But his manner of doing it made it very * Pl. 20. F. 7, easy; for he first bent the Iron to a right Angle, as at Fig. 8: then wrap- 8, and 9. ping his Handkerchief about the broad flat upper End of the Iron, he held that End in his left Hand, and with his right Hand applied to the other End twisted about the angular Point, as in Fig. 9.

N.B. *My Lord TULLIBARDIN took one of his Irons and did the same Thing before him; and indeed what was harder, for he untwisted one of the Irons that the Man had twisted.*

In order to explain how the abovemention'd Feats may be perform'd by Men of no extraordinary Strength, I have in the 6th Figure * drawn * Pl. 20. F. 6. the lower part of a Skeleton, containing so many of the Bones of the human Body as are concern'd in these Operations, making the Figure pretty large, to shew the better how the Girdle is to be applied.

The Bones mark'd, ISAPAI (a) which compose the Cavity call'd the Pelvis, contain a bony Circle or double Arch of such Strength, that it wou'd require an immense Force to break them by an external Pressure directed

L 1

rected

(a) *These Bones are thus distinguished by Anatomists. S, the Os Sacrum: II, the Ilium: AA, the Os Ischium; whose strongest part has on each Side an hemispherical Concave, in which the round Head of the Thigh Bone is receiv'd and turns round, being held by a strong Ligament in its middle: those Parts of the Bone that join together before betwixt AA and above P are call'd the Os Pubis or Ossa Pubis.*

Annotat. rected towards the Center of the Circle, or the middle of the *Pelvis*. It
 Lect. IV. is also to be observ'd, that those Parts of this bony Circumference, which
 receive the Heads of the Thigh-bone above, at, and below A, call'd the
Ischium or *Coxendix*, are the strongest of all, so that a very great Force may
 push the Heads of the Thigh-bones upwards (or, which is the same thing,
 the upper Parts of the *Coxendix* downwards) or towards each other in a
 lateral Direction from A to A, without doing any Hurt to the human Body.

Now if the Girdle above describ'd be put round the Body in the man-
 ner represented in the Figure, and be drawn downwards at G by a great
 Weight W, it will press on the *Os Sacrum* behind, and the *Ilium*; then it
 will by its Pressure on TT the great *Trochanters* of the Thigh-bones
 drive the round Heads the faster into their Sockets, so as to make them
 less liable to slip out and strain the Ligament by a Push directed upwards.
 So that the semicircular Part of the Girdle TC SCT presses together the
 bony Arch denoted by the same Letters, which, according to the nature of
 Arches, is the stronger for that Pressure. The Abutments of the Arch
 cannot come nearer together by reason of the Resistance of the strong Bones
 APA, neither can they fly outwards, because the Girdle keeps them to-
 gether. Then the Thighs and Legs TDB are two strong Columns, ca-
 pable of sustaining four or five thousand Pounds at least, provided they
 stand quite upright. The Muscles here are put to no Strain, being no
 farther concern'd than to balance each other; that is, the antagonist Mus-
 cles, *Extensors* and *Flexors* only keep the Bones in their Place, which
 makes them resist like one entire Bone form'd into an Arch.

* Pl. 20. F. 2. This shews how easily the Man of Fig. 2. * may sustain a Cannon of
 two or three thousand Pound Weight. The same Solution will also serve
 † Pl. 19. F. 1. for the Resistance of the Man of Fig. 1. Plate 19. † whom five Men (nay
 ten Men or two Horses) cannot pull out of his Situation when he sits so as to
 have his Legs and Thighs in the horizontal Line PF, or in a Line in-
 clining downwards towards A; for then, tho' there is a difference in the
 sitting Posture from the standing Posture before describ'd, yet by reason
 of the Mobility of the Heads of the Thigh-bones in the *Acetabula* or Ca-
 vities of the *Coxendix*, the Arch is the same and as strong as before, its
 Abutments being equally supported by the Legs and Thighs. It is only
 the bending of the Back-bone above the Girdle to bring up the Body
 which makes the difference of Position in the Man, tho' not sensibly
 in the resisting Parts. The Impossibility of overcoming the Resistance of
 the Man that sits with the Girdle about him, without crushing his Legs
 and Thighs end-wise into one another, depends upon what has been said
 in the * fifth Note of the third Lecture (Page 144) where we have shewn
 * Ann. L. III. p. 144. that a Power acts ineffectually upon a Leaver, when it draws it against the
 Center of Motion. This will be further explain'd by the 2d Fig. of Plate
 * Pl. 19. F. 2. 19, * where the Leaver HL, whose Center of Motion is at L, represents the
 strong Man's Legs and Thighs, the Power of the Men or Horses pulling at
 M, being applied at H, and drawing in the Direction HL. The same
 will happen when the Leaver is in the Position HL; but if the Man shou'd
 sit

fit with his Breech higher than his Feet, so as to have his Legs and Thighs in the Position of the Leaver bL , Mlb , the Line of Direction of the Power wou'd make with the Leaver the Angle lbL , whose Sine being lL wou'd reduce the Action of the Power to the same Thing as if the Weight of the Man was suspended at b the long *Brachium* of a bended Leaver bLl , and the Power shou'd draw at the Point l , by the short *Brachium* lL . Then if the Power was to the Weight in a *Ratio* something greater than that of bl to lL , the Man wou'd be pull'd upwards in an Arc whose Center is at L , his Resistance decreasing continually; because then only his Weight wou'd act against the Power, by the help of a Leaver which in its Motion wou'd continually encrease the *acting Distance* of the Power, and diminish that of the Weight. Now as this may happen to a Man sitting upon a horizontal Board, if his Girdle be a little too high, or he be pull'd on the sudden, before he is rightly fix'd and his Legs and Thighs are in their due Position; I wou'd always advise the Board to recline in the manner FG * to prevent such a Surprise, which can hardly happen then, because the Point H must rise quite above the Line LP before the Power can gain any Advantage. Nay, for greater Security, instead of the slit DL (or Ll † *Fig. 2.*) I use only an Hole at L , for the Rope to pass thro', and always be between the Legs and Feet. Annotat. Lect. IV.

I have observ'd the pretended *Strong Man* sometimes to have a short strong Stick of about a Foot long tied to the Rope at K , that in case of a Surprise, that Stick might stop against the Props D and C , so as to prevent his being drawn any farther forward in such a Case; and then he held the Stick in his Hands, pretending to pull with his Hands to make the Trick appear the more strange.

But in breaking the Rope the Muscles must act in extending the Legs; and that we may the better explain that Action, we must consider a Man breaking the Rope as represented in the first *Fig.* of *Plate 20.* * that way being more simple than when it is broken in manner of *Fig. 3.* *Plate 19.* † * Pl. 19. F. 1. † Pl. 19. F. 2.

The Rope being fasten'd to a Post at P , or any other fix'd Point, is brought thro' an Iron Eye L to the Hook of the Girdle H of the Man HI , and so fix'd to it by a Loop, or otherwise, as to be quite tight, whilst the Man's Knees are so bended as to want about an Inch of having his Legs and Thighs quite upright. Then if the Man on the sudden stretches his Legs and sets himself upright, he will with Ease break the very same Rope which held two Horses exerting their whole Strength when they draw against him; such as a Cart Rope, or a Rope of near three quarters of an Inch Diameter, which may be broken by a Man of middling Strength, by the Action of the ten Muscles (*a*) that extend the Legs, five belonging to each Leg.

L 1 2

If

(*a*) The five Muscles that extend each Leg are describ'd by the Anatomists. Their Names are, 1. *Membranofus* arising from the upper part of the Spine of the Os Ilium, and inserted a little below the Knee into the outer and the fore-side of the Tibia and Fibula. 2. The *Rectus* springing from the lower part of the Spine of Os Ilium, and inserted also a little below the Knee into the fore-side of the

Annotat. i. If the Rope is strong enough to bear 1800 lb, but will break by hanging a little more Weight to it; two Horses, or ten Men, cannot break it by fair pulling or drawing against the sitting Man of *Fig. 1. Plate 19.* *


* Pl. 19. F. 1. For as an Horse in common hard Work of six Hours a Day can only pull † Page 241. 240 lb, † he cannot draw more than double that Weight when he is whipp'd and exerts himself; so that two Horses, or ten Men, equivalent to them, cannot with a Jirk draw above 1000 lb, whereas the Rope has been suppos'd strong enough to sustain 1800 lb, and yet it may be broken

* Pl. 20. F. 1. by an ordinary Man in the Posture of *Fig. 1. Plate 20.* * Neither need we wonder that the Muscles *Extensors* of the Legs should exert so much Force, when we consider their Bigness and Length; especially if we compare them with the four Muscles that pull up the lower Jaw-bone (which tho' they all four do not weigh a Pound) yet enable some Men to crack an Apricock or a Peach-stone, which could not be broken without an immense Weight. † See the 6th Note of our third Lecture, from Page 153 to Page 159; or more fully to satisfy one's Curiosity, one may consult *Borelli's Book de Motu Animalium*, where he has shewn the particular Strength of the Muscles.

* Pl. 19. F. 3. The manner of breaking the Rope as represented in * *Fig. 3. of Pl. 19.* tho' more troublesome, is also more effectual for breaking the Rope, than that which we have been describing; for the same Man may break a Rope in this Position, which he cannot break in the other. To understand this we must observe, that the Man takes the Rope so short, that when he climbs up against the Post, if the Eye L (thro' which the Rope passes) be between his Toes, his Heels being lower at T, when his Knees are streight, the length of his Legs and Thighs TH is greater than the length of the Rope and Girdle from L to H; so that we may consider in the Man and Rope the Triangle Lh_t drawn below the Man in the Figure; the Side Lh, representing the Length of the Rope and Diameter of the Girdle; the Base Lt, the Man's Feet; and the longest Side th the extended Limbs or the Legs and Thighs when streight. Now in the Rotation of the long Sides of this Triangle, when the Side Lh comes to be horizontal at Lr (moving in the Arc hrr round the Center L, the Side th will be in the Situation tb (as it moves round the Center t in the Arc hbs) and consequently either the Rope must stretch from r to b, or the Point h, by the bending of the Knees must be brought nearer and come to r; or else the Rope must break, which is what will happen, especially when we consider that the lower the Body (with the Limbs stiff) comes down round the Heels for the Center of Motion, the greater will the Distance rb be, as we may see a little lower at rs; so that if the Rope did stretch a little at first, it must break at last, and the Man fall down upon the Feather-bed,

or

the Tibia. 3. The Vastus Externus, springing from the Root of the greater Rotator, and inserted a little below the Patella, near the same Place with the former. 4. The Vastus internus, which arises from the Root of the lesser Rotator, and likewise is inserted a little below the Patella. 5. The fifth is the Crureus springing from the forepart of the Thigh bone, between the two Rotators, and ending in the same Place with the former.

or other soft Body to receive him, at B. For if the Man finds that the Annotat. Action of the Muscles Extensors of the Legs does not break the Rope, Lect. IV. he can in this Position easily add the whole Weight of his Body with a  Swing and a Jerk as he throws himself backwards.

The Posture of Fig. 4. Plate 19. * (where the strong Man having an* Pl. 19. F. 4. Anvil on his Breast or Belly, suffers another Man or two to strike with a Sledge Hammer and forge a Piece of Iron, or cut a Bar cold with Chiz-zels) tho' it seems surprizing to some People, has nothing in it to be really wonder'd at; for sustaining the Anvil is the whole Matter, and the heavier the Anvil is, the less are the Blows felt; and if the Anvil was but two or three times heavier than the Hammer, the strong Man would be kill'd by a few Blows. This will be easily understood by calling to mind what we have said in the second Lecture; for the more Matter the Anvil has, the more *Inertia* and the less liable it is to be struck out of its Place; because when it has by the Blow receiv'd the whole *Momentum* of the Hammer, its Velocity will be so much less than that of the Hammer as it has more Matter than the Hammer. Neither are we in that Case to attribute to the Anvil a Velocity less than the Hammer in a reciprocal Proportion of their Masses or Quantities of Matter; for that would happen only if the Anvil was to hang freely, in the Air (for example) by a Rope, and it was struck horizontally by the Hammer; but the Resistance of the Ribs which make an Arch under it, will still diminish that Velocity: So that if the Hammer striking the Anvil when hung freely, could make it move an Inch out of its Place; by putting a Resistance behind it equal to the Weight of the Anvil it would move but half an Inch, and but a quarter of an Inch if that Resistance was double, &c. Thus is the Velocity given by the Hammer distributed to all the Parts of a great Stone, when it is laid upon the Man's Breast to be broken; but when the Blow is given, the Man feels less of the Weight of the Stone than he did before, because in the Reaction of the Stone, all the Parts of it round about the Stone, rise towards the Blow, and if the Tenacity of the Parts of the Stone, is not stronger than the Force with which it moves towards the Hammer, the Stone must break; which it does when the Blow is strong and struck upon the Center of Gravity of the Stone.

N.B. That the Parts of Bodies struck move towards the Blow, is a Consequence of a Law of Nature, which shall be explain'd in my next Lecture.

I should be too tedious to be as particular in my Explanation of the other Feats of Strength as I have been in these above-mentioned. Therefore I will only lightly consider the following ones; especially, since the Principles already explain'd in the Lectures past, and the Considerations just mention'd, will enable any one easily to discover the Reason of all such Performances.

In the 5th Fig. of Plate 19. † The Man IHL (the Chairs I, L, be-† Pl. 19. F. 5. ing made fast) makes so strong an Arch with his Back-bone and the Bones of his Legs and Thighs, as to be able not only to sustain one Man, but three

Annotat. three or four, if they had Room to stand; or, in their Stead, a great Lect. IV. Stone to be broken with one Blow.

Pl. 19. F. 6. In the 6th and 7th Figure of the same Plate * a Man or two are rais'd in the Direction CM by the Knees of the strong Man IHL lying upon his Back. Now we must observe that the five Muscles (a) which bend the Legs (tho' weaker than the Extensors, because they are not to carry the Body in our common Motions) act with their greatest Force at the Beginning of this Operation, as all Muscles do when from their full Extent they begin to contract; and to relieve them in their Action as the Heels come forward from the Point L they stop against the Ground and keep the Body M in the Place which it is risen to: So that the Action of these Flexors is reiterated, and they have time to be recruited with fresh Spirits (or whatever Fluid is the first Cause of their Inflation) and when they are so far contracted as to act more weakly, the Pressure of the Weight affects them less and less, the Bones supporting more of the Weight as they become more perpendicular, and consequently the Muscles have less occasion to act. See

* *Pl. 19. F. 7.* Fig. 7. * The rest of the Performance, viz. of setting the Man M upon a Table, is very easy and obvious, the strong Man having now only his own Body to lift up, which he helps by putting his Hands round the Mans Feet or Hams, and in raising himself up, rather pushes him off of his own Knees than lifts him up upon a Table plac'd at N at Arms ends, as he pretends. †

† *Pl. 19. F. 7.* In breaking the Rope one thing is to be observ'd, which will much facilitate the Performance; and that is to place the Iron Eye L, * thro' which the Rope goes, in such a Situation, that a Plane going thro' its Ring shall be parallel, or nearly parallel to the two Parts of the Rope; because then the Rope will in a manner be jamm'd in it, and not slipping through it, the whole Force of the Man's Action will be exerted on that part of the Rope which is in the Eye, which will make it break more easily than if more Parts of the Rope were acted upon. So that the Eye, tho' made round and smooth, may be said in some measure to cut the Rope. And it is after this manner that one may break a Whip-cord, nay, a small Jack-line with one's Hand without hurting it; only by bringing one part of the Rope to cut the other; that is, placing it so round one's left Hand, that by a sudden Jerk, the whole Force exerted shall act upon one Point of the Rope. See the 11th Figure of Plate 20 † where the Cord, to be broken at the Point L in the left Hand, is mark'd according to its Course,

(a) The five Muscles that bend the Legs are these. 1. The Longissimus, or Fascialis, arising from the inner Knob of the Os Ilium, and a little above the Knee ending in a Tendon; which is inserted under the Knee, into the fore and inner Side of the Tibia. 2. The Gracilis springing from the jointing of the Os Pubis, and inserted by a strong Tendon a little lower than the former, in the inner Side of the Tibia. 3. The Seminervosus arising from the Knob of the Ichium, turning into a round Tendon under the Ham and inserted also into the inner Side of the Tibia towards the back-side, running as far as its Middle. 4. The Semimembranosus proceeding from the same Knob, and ending by a broader Tendon than the third, in the hinder part of the Tibia. 5. The fifth is call'd a Biceps, likewise beginning at the Ichium, and at last inserted into the outer Side of the upper Appendix of the Fibula.

Course, by the Letters R T S L M N O P Q, folding once about the right Annotat. Hand, then going under the Thumb into the middle of the left Hand, Lect. IV. where crossing under another Part it is brought back under the Thumb again to M, then round the back of the Hand to N, so thro' the Loop at L to O, and three times round the little Finger at P and Q; which last is only that the Loop N O may not give way. Before the Hands are jerk'd from one another, the left Hand must be shut, but the Thumb must be held loose, left pressing against the Fore-finger it should hinder the Part T L of the Rope from carrying the Force fully to the Point L; but the little Finger and that next to it must be held hard, to keep the Loop N O firm in its Place.

The making use of the Muscles that extend the Legs for lifting great Weights is no uncommon Practice among some of our working Men, tho' it is not observ'd, because it is done without an *Apparatus*. We see Hackney Coach-men often get out of their Boxes and with their Rumps easily lift up the Coach behind to make way for one another, or to avoid some great Rub, or some Hole, or any other Inconveniency; and this they do with so much ease, that if they have four Persons in the Coach, and three or four Trunks behind, they never think it worth while to desire any Person to get out, or to take off any of the Weight. The Coal-Porters at the *Custom-house* Key (commonly call'd *Fellowship Porters*) carry one hundred and three quarters Weight of Coals, running all the way, tho' at every Turn they go up two Ladders, and often the length of St. *Dunstan's* Hill, which is a Street pretty steep and ill pav'd, and perhaps climb up a Stair-case or two before they shoot their Coals: and this most of them will do above sixty times a Day. But their manner of doing it, is to stoop so as to let the Sack bear chiefly on their Rumps, holding one Hand behind them to keep together the Mouth of the Sack, that they may with more Expedition shoot out their Coals, whilst their other Hand secures the Sack from slipping down from above; and this Posture very much eases the Action of the Muscles of the Loyns, the Extensors of the Legs being then chiefly concern'd.

Since I began to write this, I have been credibly inform'd that the Porters in *Turkey* carry seven or eight, nay sometimes nine hundred Weight upon the lower end of their Back, or rather their Rump, only resting on a Stick before them, whilst they receive their Loads, to support their Body and save the Muscles of their Loyns; but we may easily guess that other Persons must be very careful in putting on and taking off the Burthen.

I believe the Strength of the *Testudo* made by the *Roman* Soldiers when they stood close together with their Shields over their Heads, must be owing to some such sort of Posture of the Body: otherwise they could never have been able to bear the Weight of Chariots driving over them, as some Historians have inform'd us. In that Case every Man, except those of the first Row, (a) cover'd the Man who stood before with his Shield

at

(a) Those of the first Row held their Shields inclin'd before them as they stoop'd. Sometimes the first Row kneeled, and the second Row bore on the Shoulders of the first as they stoop'd, and cover'd them with their Shields, &c.

Annotat. at the same time bearing upon his Rump that stood before him: and when
 Lect. IV. they stood against any Shock, their Muscles had no other Labour but to
 keep their Knees stiff; the bony Arch already describ'd † being sufficient
 † Pl. 20. F. 6. to support a much greater Weight.

There are several Cases, wherein it would be of singular use to apply the Force of one or more Men, by means of the Girdle and Hook and Chain, in the manner abovementioned; as for example, when the Resistance is very great, but the Bodies that resist are to be remov'd but a little way: if we lift very heavy Goods a small Height to remove any thing from under them: if we would draw a Bolt or Staple, and find we can't do it even with an Iron Crow, the Hand pulling it upwards at the End, then the Hook of the Girdle being applied at the End of the Crow, the Force exerted by stretching the Legs would be tenfold of what the Hands were able to do, without more help at the same Place.

There may also be many Occasions on Board a Ship. I'll instance but
 * Pl. 21. F. 1. one. Let FG* be the Tackle for raising or lowering the Main-top-mast, part of which is represented by m_1 , m_2 ; the Block G is fix'd below, and as the Block F comes down it pulls along with it the top Ropes FBC, m_1 running over the Block B (fix'd at A) and round the Block C in the Heel of the Top-mast, so as to draw up the lower End m_1 of the said Main-top-mast, which when hoisted up to its due Height, is made fast by the Iron Pin or Fidd I which is thrust thro' it, and then its own Weight and the Hole D of the Cap will keep it in its Place. We'll suppose that the Force requir'd thus to raise the Mast must be that of six Men pulling upon Deck at the fall of the Tackle, that is, at the running Rope FGK at K on the other side of the Main-mast LI. Now in order to let down this Mast on the sudden, as in case of hard Weather it is necessary, the same Tackle and Power must be made use of, tho' it be but to lift it a very little Way, that a Man may be able to get out the Fidd I before the said Mast can be let down and slip to N on the side of the Main-mast. I say, that if the Hands are so employ'd otherwise, that instead of six Men there be only one Man at the Rope K; if he has a strong Girdle to which he fastens it (or makes a Bow in the Rope it self to fix it round the lower part of his Back, &c.) he may exert much more Force in the Direction GK than the six Men in the common way of pulling: and if he draws to him (sitting on the Ground and pushing his Feet against the first firm Obstacle that he finds, as against OP) only two Inches of the Rope KG, he will raise up the Main-top-mast the third part of an Inch, which will be sufficient for the Iron or Fidd I to be drawn out.

N. B. If more Force should be requir'd for this Operation, as in large Ships, several Men at once might make use of Ropes about their middles instead of Girdles, and fasten them all to different parts of the Fall of the Tackle; and for fix'd Points they might set their Feet against the wooden Steps of a Stern Ladder taken down and lying on the Deck, fasten'd at one end to one of the Ring Bolts: for tho' in this case each Man could not apply so much Force as the single Man before suppos'd,
 because

because as they all must sit a little on one side of the Rope GK, their Pull will be something oblique; yet five Men in this case, will very easily do the Work of fifteen.

Annotat.
Lect. IV

8. [Page 243.— The whole Force whereby a Man draws, &c. — I have given a Translation in my Notes, &c.]

To illustrate and confirm what I have said in the Lecture, I shall here give part of *Mons. De la Hire's* Memoire given in to the Royal Academy of Sciences, in the Year 1699, entitled, *An Examination of the Force of Men to move Weights, whether by lifting, carrying, or drawing; consider'd as well absolutely as when compar'd with that of other Animals which carry and draw, as Horses, &c.* In which all his Reasoning is just, tho' some of his *Data* being wrong, lead us also to a wrong Conclusion; but I shall set all right by my Observations upon it.

“ I suppose first that a Man of a middle Stature who is pretty strong, weighs 140 lb of our Weight. (*) I consider first, that such a Man as I have suppos'd, having both Knees on the Ground, can rise bearing only on his Toes, still keeping his Knees together; and as this Action is perform'd by means of the Muscles of the Legs and Thighs, it is evident by the Supposition I made of his Weight, that the Muscles of the Legs and Thighs have a sufficient Force to raise 140 lb. (a)

“ But a Man bending a little in the Hams can raise himself up though loaded with a Weight of 150 lb, together with the Weight of his Body, which he raises at the same time; (b) so that the Force of the Muscles of the Legs and Thighs can raise a Weight of 290 lb, that is 150 lb the Burthen carried, and 140 lb the Weight of the Body, when the Rise is but 2 or 3 Inches.

“ Such a Man, as we have suppos'd, and shall all along suppose him, can also lift from the Ground a Weight of 100 lb plac'd between his Legs, taking hold of it with his Hands as with two Hooks, and raising himself up. (c) Whence it follows that the Muscles of the Loyns alone have Force enough to raise 170 lb, namely, the 100 lb of the Weight, and 70 lb half the Weight of the Man; because he is to raise not only the 100 lb Weight, but also the whole upper part of his Body above the middle, because we have suppos'd him to stoop to take up the Weight.

“ As for the strength of the Arms for drawing or lifting a Weight, one may suppose it of 160 lb, which depends upon the Muscles of the Shoulders and Arms. For if a Man with both Hands takes hold of some fix'd Body plac'd over his Head, he may easily enough, by the strength of his Arms alone, draw up his Body, and even 20 lb more, just as if he was burthen'd with a Weight of 20 lb. It is easy to make the Experiment; for if a Weight of 160 lb be fasten'd to a Rope and thrown over a Pulley, and a Man who weighs only 140 lb pulls at the other end

M m

“ of

(a) The French lb is betwixt one 11th and one 12th part greater than our Pound Averdupois.

Annotat. " of the Rope, it is plain that he will never be able to raise the 160 lb
 Lect. IV. " Weight; the utmost he can do being only to hang with all the Weight
 ~~~~~ " of his Body by the Rope; for the Weight at the other end being greater  
 " er than the Weight of the Man, will keep him suspended; the Pulley  
 " being only a continued Balance with equal Arms: but if the Man has  
 " fasten'd to him a Weight of 20 lb, then he will make an *Equilibrium*  
 " with the Weight on the other Side; and if ever so little be added to the  
 " 20 lb Weight he will raise the opposite Weight, the Muscles of his  
 " Shoulders and Arms being of sufficient Force to sustain the whole aggregate  
 " Weight.

" Tho' the Muscles of each part of the Body can exert great Forces  
 " to raise Weights, the Force of a Man is not to be reckon'd as the Sum  
 " of the different Forces of all his Muscles taken together, even tho' the  
 " Spirits which swell the Muscles, and by shortening them draw the Tendons  
 " at their Ends cou'd as easily be distributed to all the Parts as to any one  
 " particular Part, because each Part commonly serves to support that which is  
 " immediately next to it. As for Example; the Muscles of the Arms and  
 " Shoulders by their Contraction can lift a Weight of 160 lb; but if the  
 " Body be inclin'd, the Arms will not be able to sustain that Weight, unless  
 " the Muscles of the Loyns are at the same time strong enough to sustain the  
 " upper part of the Body, together with the Weight which it is loaded  
 " with; and if the Hams were also bent, then the Muscles of the Legs  
 " and Thighs must still exert a greater Strength, because they must sustain  
 " the Weight of 160 lb and also at the same time the whole Weight of  
 " the Body. Whence it happens, that in this Disposition of the whole  
 " Body, the Force is distributed by the Distribution of the Spirits into  
 " all the Parts; for which reason a Man will not be able to raise 160 lb  
 " from the Ground.

" Not but there are Men, whose Spirits flow so abundantly and so swiftly  
 " ly into their Muscles, that they exert three or four times more Strength  
 " than others do; and this seems to me to be the natural reason of the  
 " surprizing Strength that we see in some Men who carry and raise Weights  
 " which two or three ordinary Men can hardly sustain, tho' these Men be  
 " sometimes but of a moderate Stature, and rather appear weak than  
 " strong. There was a Man in this Country a little while ago, who wou'd  
 " carry a very large Anvil, and of whom were reported several wonderful  
 " Feats of Strength; but I saw another at *Venice*, who was but a Lad,  
 " and did not seem able to carry above 40 or 50 lb with all possible Advan-  
 " tages: Yet this young Fellow standing upon a Table, rais'd from the  
 " Earth and sustain'd off of the Ground an Ass, by means of a broad Girt,  
 " which going under the Creature's Belly, was hung on to two Hooks that  
 " were fasten'd to a Plat of small Cords, coming down in Tresses from the  
 " Hair on each side the Lad's Head, which were in no great Quantity;  
 " and all this great Force depended only upon the Muscles of the Should-  
 " ers and those of the Loyns (*d*); for he stoop'd at first whilst the Hooks  
 " were fasten'd to the Girt, and then rais'd himself and lifted up the Ass  
 " from

“ from the Ground, bearing with his Hands upon his Knees. He rais’d Annotat.  
 “ also in the same manner other Weights that seem’d heavier, and us’d to Lect. IV.  
 “ say he did it with more ease, because the As kick’d and struggled when  
 “ first lifted off of the Ground. (e)

“ Now to examine the Force of a Man to carry a Burthen upon his Shoul-  
 “ ders: I say that Weight may be of 150 lb, and a Man may walk with  
 “ that Load easily enough upon an horizontal Plane, provided he does not  
 “ take great Steps; but he can by no means go up a Mountain or a Stair-  
 “ case with the same Weight. For the Action of walking, when we bear a  
 “ Burthen, must be consider’d as the circular Motion of C, the common Cen- Pl. 20. F.  
 “ ter of Gravity of the Body and the Weight together about the Foot F, 12.  
 “ that advances as the Center of the Arc of Motion, the Effort of the  
 “ Muscles of the other Leg which act against the fix’d Point D, serving  
 “ only to drive the Center of Gravity forward; and if the Arc CE,  
 “ which that Center describes, be small, the Effort of the hind Leg need  
 “ not be great to make the Center describe it; because then it pushes up  
 “ the Weight of the Body, and the Burthen only the Height of AB the  
 “ vers’d Sine of half the Arc, which is inconsiderable in this Case in re-  
 “ spect to the whole Arc, which is the length that the Burthen advances.

“ So that a Man carrying a Burthen, can walk so much the easier as his Steps  
 “ are shorter, because the vers’d Sine will be so much the less; for if he shou’d  
 “ take large Steps, the hind Leg must push up the Body and Burthen the  
 “ Height of the vers’d Sine of so much a greater Arc as the Steps are  
 “ greater; that is, the vers’d Sine of the Arc which measures half the Di-  
 “ stance that the Burthen advances forward.

“ It is also plain, that a Man thus loaded can by no means go up a Stair-  
 “ case, or a very steep Hill with that Burthen; because according to what  
 “ we have already explained, the Action of the Muscles of the Legs being  
 “ only able to raise a Weight of 150 lb to the Height of two or three Inches,  
 “ he wou’d not be able to raise it five Inches, which is the Height of  
 “ common Stairs or Steps, (f) nor go up a Mountain, unless he took such  
 “ short Steps as to rise only two or three Inches perpendicularly every time.

“ What remains now is to consider the Force of a Man for drawing or  
 “ thrusting horizontally. (g) But to make this the more clear and intelli-  
 “ ble, I shall consider that Force as apply’d to the Handle of a Roller or  
 “ Windlass whose Axel is horizontal, on which Axel the Rope which sus-  
 “ tains the Weight winds up, supposing the Distance from the Center of the  
 “ Roller to the Elbow of the Handle to be equal to the half Diameter  
 “ of the Roller, that the Force applied may be estimated without any  
 “ Augmentation on account of the Machine: neither do I consider here  
 “ the Friction of the Pivot or Difficulty of folding the Rope.

“ First then, it is plain, that if the Elbow of the Handle be in an hori-  
 “ zontal Situation, and about the Height of a Man’s Knees, a Man that  
 “ lifts it by drawing up, may exert a Force capable of raising 150 lb  
 “ hanging at the end of the Rope if he takes all possible Advantages, ac-  
 “ cording to what I have already explain’d. But if he is to depress the

Annotat. " Handle his Strength for that purpose cannot be more than equal to 140 lb,  
 Lect. IV. " which (as has been suppos'd) is the Weight of his Body; for he can  
 lay on no more Weight than that, except he was loaded with some  
 Weight, and then the Effect might be so much greater.

" *Secondly*, If the Elbow of the Handle be plac'd vertically, and it be at  
 the same Height as a Man's Shoulders, it is certain that a Man can ex-  
 ert no Force to cause it to turn, by pushing or drawing with his Hands,  
 if both Feet being set together the Body be upright as represented  
 \*Pl. 20. F. 13. " in \*Fig. 13. by the Line AP, the Line AM representing the Line of  
 the Arms being horizontal, and making a right Angle with AP; be-  
 cause in that Position, neither the Force of the whole Body, nor of any  
 part of it, nor its Weight, can exert any Force for drawing or pushing;  
 which is known by Mechanicks; for I look upon the Breadth taken up  
 by the Feet but as a Point at P. But if the Handle be higher or lower  
 than the Level of the Shoulders, then the Line AM, which goes from  
 the Shoulders to the Hands, and the Line AP going from the Shoulders  
 to the Feet, will make an obtuse or an acute Angle, and then a Man  
 may exert some Force for drawing or pushing the Handle; but that  
 Force depends only upon the Weight of the Body, as may be easily per-  
 ceiv'd and demonstrat'd; and that Weight or Force must be consider'd  
 as collect'd in one Point, which is the Center of Gravity of the Body,  
 and about the Height of the Navel: I say, we must have regard only  
 to the Weight of the Body, in order to determine the *Æquilibrium*; for  
 the Action of the Muscles of the Legs and Thighs serves only to pre-  
 serve that *Æquilibrium* as we walk.

\* Pl. 21. F. 1. " Let the Handle in Fig. 1. \* be plac'd at D the same height as the  
 Shoulders A, and the Center of Gravity of the Body be at C, the Body  
 being very much inclin'd towards the Handle; but the End of the Feet  
 must be at P. *First*, We must consider that Point P as the *Fulcrum* or  
 Center of Motion of a Leaver or strait Rod PCH, which going thro'  
 C the Center of Gravity of the whole Body meets the Line of the  
 Arms MA at the Point H. *Secondly*, We must consider the Point C of  
 the Leaver loaded with the whole Weight of the Body equal to 140 lb,  
 which Weight endeavours to descend perpendicularly according to its na-  
 tural Direction, the end H of the Leaver being sustain'd in the horizon-  
 tal Direction MAH; whence it will be easy to determine by mechan-  
 ical Principles, what Effect the Weight of the Body at C, acting in its  
 natural Direction, can have on the Handle according to the horizontal  
 Direction HD.

" For, first, let us suppose PH to be divided into 240 Parts, of which  
 PC contains 80. Now since the Weight of the whole Body acting at  
 C is but 140 lb, when apply'd at H it will act but with the Force of 80  
 lb; just as if a Weight of 80 lb was to be suspended at that Point and  
 act in its natural Direction, which, according to what we have suppos'd,  
 must be perpendicular to MA. If therefore from the Center of Mo-  
 tion P the Sine PF be drawn perpendicular to MAF, the Weight  
 " of

“ of 80 lb at H acting in its natural Direction: will be to the Action of Annotat.  
 “ the same Weight upon the Handle according to the horizontal Directi- Lect. IV.  
 “ on MAH :: in the Ratio of PF: to HF. And this does very much  
 “ diminish the Action of the 80 lb in a moderate Inclination of the Body  
 “ ACB. And if we suppose for Example, that the Line PCH makes an  
 “ Angle PHF of 70 Degrees with the Line MAF, the Line of the Bo-  
 “ dy ACB will then be inclin'd to the Horizon in an Angle of more than  
 “ 60 Degrees, which is the greatest Inclination of Body with which a  
 “ Man can walk; then

“ The Sine of 70 Degrees, which is PF:

“ Will be to the Sine of its Complement, which is HF ::

“ As 3:

“ To 1, nearly;

“ And consequently the Action of the 80 lb suppos'd to act at H accord-  
 “ ing to the natural Direction, will be to the Action exerted in the hori-  
 “ zontal Direction, but as 80 to its third, which is something less than  
 “ 27 lb.

“ Those who have not made the Experiment of pushing horizontally with  
 “ the Arms, or drawing a Rope horizontally as a Man walks with the Bo-  
 “ dy inclin'd forward, whether the Rope be fasten'd towards the Should-  
 “ ers or towards the middle of the Body (for the Effect will not differ pro-  
 “ vided the Inclination of the Body is the same; because the Sine of In-  
 “ clination and its Complement are always in the same Ratio) can hardly  
 “ be persuaded that the whole Force of a Man is reduc'd to so little in  
 “ pulling horizontally as not to exceed 27 lb.

“ Not but that a Man may lean or incline his Body so as to sustain a  
 “ much greater Weight than 27 lb; for if the Line PH should make an  
 “ Angle of 45 Degrees with HF, it is certain that the Weight of the  
 “ Body would sustain 70 lb, but as he must then have his Body in the  
 “ Position of the Line AB much more inclin'd to the Horizon than 45  
 “ Degrees, he would be so far from being able to walk, that he would not  
 “ be able to stand in such a Posture.

“ The same Demonstration may also serve to shew that a Man has much  
 “ more Force to draw if he goes backwards than in going forwards. For  
 “ in such a Situation of the Body, the Line PCH, \* Fig. 2. which goes \*. Pl. 21. F. 2.  
 “ from the Feet at P thro' the Center of Gravity C, and on which the  
 “ Encrease of Force depends, will always be much more inclin'd to the Ho-  
 “ rizon than the Line of the Body ACB, quite contrary to what it was  
 “ in the former Position. But then this way of drawing cannot be put in  
 “ practice, except for pulling a Rope, when a Man continues in the same  
 “ Place; otherwise we should naturally come into that Posture; for Nature  
 “ and Experience have taught us to take all possible Advantages in common  
 “ Operations.

“ For the same Reason Watermen and those that row at Sea always pull  
 “ their Cars backward; for that way they can exert more Force than in  
 “ pushing forwards like the Gondoliers at Venice, who work in that manner,

“ for

Annotat. "for no other Reason than to see well before them, which is more necessary to them than exerting a great Force, by reason of the many short Turns they must make in the Canals, and the Care they must take to avoid one another."

What remains is to compare the Force of Men with that of Horses for drawing, &c. but having consider'd that in my Lecture, I need not give the Translation of the rest of *Mons. de la Hire's* Paper; only proceed to make my Observations upon it.

(a) *The Muscles of the Legs and Thighs are much stronger than M. De la Hire supposes, as appears by what has been said concerning the Feats of Strength; and the Action of rising backwards from ones Knees is very short of the utmost that a Man can do in that Posture; for a Man may carry some considerable Weight and yet rise from his Knees, tho' the Arc describ'd by the Center of Gravity be then an Arc of a good many Degrees.*

(b) *It is usual in London for Men to raise themselves up with, and carry 250 lb, which is almost as much more as M. De la Hire has suppos'd; therefore all the Consequences drawn from this Supposition must fall short.*

(c) *Working Men generally lift 150 lb with their Hands, and some 200 lb; but here the Excess of the Force of the Muscles of the Loyns is not so much greater than M. De la Hire's Supposition, as the Excess of the Force of the Muscles of the Legs.*

(d) *What he attributes here to the Muscles of the Loyns was really perform'd by the Extensors of the Legs; for the young Man's stooping with his Hands upon his Knees was not with his Body forwards and his Knees stiff, but his Body upright and his Knees bent, so as to bring the two Cords with which he lifted to be in the same Plane with his Ankles and the Heads of his Thigh-Bones; by which means the Line of Direction of the Man and the whole Weight came between the strongest part of his two Feet, which are the Supports: then as he extended his Legs he rais'd himself without changing the Line of Direction. That this must have been the manner I am pretty well assur'd of, by not only observing those that perform such Feats, but having often try'd it myself. As for the Muscles of the Loyns, they are incapable of that Strain, being above six times weaker than the Extensors of the Legs, at least I found them so in myself.*

About the Year 1716, having the Honour of shewing a great many Experiments to his late Majesty King GEORGE the First, his Majesty was desirous to know whether there was any Fallacy in those Feats of Strength that had been shewn half a Year before by a Man, who seem'd by his Make to be no stronger than other Men: upon this I had a Frame of Wood made to stand in (such as is represented by the 2d \* Fig. of Plate 20.) and with a Girdle and Chain lifted an Iron Cylinder made use of to roll the Garden, sustaining it easily when once it was up. Some Noblemen and Gentlemen who were present, try'd the Experiment afterwards, and lifted the Roller; some with more Ease, and some with more Difficulty than I had done. This Roller weigh'd 1900 lb, as the Gardener told us. Afterwards I try'd to lift 300 lb with my Hands (viz. two

\* Pl. 20. F. 2.

two Pails with 150 lb of Quicksilver in each) which I did indeed raise from the Ground, but strain'd my Back so as to feel it three or four Days; which shews that, in the same Person, the Muscles of the Loyns (which exerted their Force in this last Experiment) are more than six times weaker than the Extensors of the Legs; for I felt no Inconveniency from raising the Iron Roller. Annotat. Lect. IV.

(e) The Reason why the A's kicking and moving made his Weight more inconvenient than an heavier Burthen, was, that by such a Motion the Line of Direction vacillated, and as it went forwards and backwards, the Muscles of the Loyns were forc'd to act to bring it into its Place again.

(f) That M. De la Hire has taken the Strength of the Muscles that extend the Leg and stretch the Foot, too little, appears from the Practice of all those that carry Grain in Sacks and Meal, who easily go up Stairs with 200, and often above 250 lb Weight; and the Men who carry Coals from Carts into Houses, go up Stairs with 250 lb Weight. They can't indeed go down Stairs with so great a Weight as they can go up.

(g) All that follows in this Dissertation concerning a Man's pushing or drawing may be depended upon; because it is mathematically deduc'd from the Weight suppos'd, and cannot but be true in a Man weighing 140 lb.

9. [Page 245. — *there must be a sufficient Velocity given, &c.*] If there should be no greater Velocity given to the Hand which is to turn the Winch, than I have suppos'd, there could not be Motion enough communicated from the advantageous to the disadvantageous part of the Revolution, as would enable a Man to raise 30 lb with the same Velocity as the Hand mov'd: therefore we must so contrive Matters as to encrease the Velocity of the Hand moving, at least one sixth part; but then we must lay only 25 lb upon the Hand moving so much the swifter, which yet will perform as much Work in a Day as if the other way had been practicable. Nay, in a great many Cases it will not be amiss to give the Hand one third more Velocity, and load it but with 20 lb, especially if a Fly be made use of; and the bigger the Circle is which the Fly describes, the better will the Force be distributed. This Work thus perform'd may be continued ten Hours a Day, and very little fatigue the Labourer.

10. [Page 246. — *Daily Labour of the London Porters, &c.*] At the Custom-house Key, and at several Wharfs one may observe what great Loads are carry'd by Porters employ'd to carry Goods to and from the Vessels on the Water: Such indeed are the Loads which some carry, that an Horse would soon be kill'd by carrying the same Weight. They that work for the Cheesemongers at so much *per Ton*, generally carry 300 Weight of Cheese at every Turn, and work all Day long.

11. [Page 247. — *the Horses strong, &c.*] If too great a Load be laid upon an Horse's Back, there is danger of breaking it; which is the Reason that People do not generally lay on very great Loads. The Felt-mongers and Skinners, are said to load their Horses more than any other People; but they

Annotat. they lay several Skins and Hides over the Horses Shoulders and Hips, that  
 Lect. VI. the Back may not be too much strain'd: I am inform'd that they put on  
 sometimes 4 or 500 Weight; but then the Horses go extremely slow.  
 The most that can be done with an Horse is to make him draw: and those  
 Horses that do most Work are such as draw great Loads in Carts on ve-  
 ry high Wheels (the Horses themselves being very tall) up St. *Dunstan's*  
 Hill in the *East*, where a Carter does sometimes lay on 2000 lb Weight,  
 and makes one Horse draw it up the Hill; but at every difficult Place, the  
 Man sets his Shoulder to the Cart in such a manner, as considerably to help  
 the Horse, who would be unable to draw up the Load without that Help:  
 and so sensible is the Horse of it, that he does not exert his whole Strength,  
 till he finds that his Master helps him.

I promis'd in the 15th *Note* to *Lect. III.* to give an Account of Mr. *Al-  
 len's* Carriages at *Bath*, and having just receiv'd the Account from a Friend  
 well skill'd in Mechanics and drawing, who took the Measures and Draughts  
 upon the Place, think proper to communicate it here, as having consider'd  
 the Nature of Carriages in this Lecture.

*A DESCRIPTION of the Carriages made use of by Ralph Allen, Esq; to  
 carry Stone from his Quarries, situated on the top of a Hill, to the Wa-  
 ter-side of the River Avon, near the City of Bath. By CHARLES de  
 LABELYE.*

These Quarries are at the Distance of a Mile and an half from the River,  
 and about 500 Feet above the Level of its Surface, which makes a Slope so  
 steep, that the small Price the Stone is sold at, would hardly defray the  
 Charges of bringing it down without some proper Contrivance, such as the  
 following, which is a great Improvement on some Carriages and *Waggon-ways*  
 made use of at the Coal Mines near *New-Castle*.

\*Pl. 21. F. 4. 1. *Fig. 4. Plate 21.* \* represents this Carriage in Perspective, as seen  
 from a Distance of 12 Feet from the left of the fore Wheels, the Height  
 of the Eye being about 6 Feet. The *Geometrical Plan* was laid down from  
 a Scale of 20 to an Inch; that is, every 20th of an Inch, answers to one  
 Inch in the Machine.

\*Pl. 21. F. 5. 2. *Fig. 5.* \* represents the Elevation of one of the Sides of this  
 Carriage, when both the Fore-wheel and Hind-wheel of that Side are lock'd,  
 from a Scale of 20 to an Inch.

†Pl. 21. F. 6. 3d. *Fig.* the 6th. † represents the Elevation of the hind part of the  
 Carriages, with all the Iron-work employ'd in the locking of the Wheels,  
 and it shews also the Profil of the Wheels, and Axel-tree, together with  
 the Sections of the Side-planks and of the Frame on which it moves, from  
 a Scale of 10 to an Inch.

4. From the Consideration of these three Figures, it will appear, that  
 this Carriage consists of a strong Floor of (Oaken) Planks, three and a  
 half

half Feet wide, and about 13 Feet long, strengthen'd above by several Annotat. Ribs to defend it from the Stones that lye upon it, and fix'd upon four Lect. IV. Beams of the same Wood, about four Inches square and 14 Feet long.

5. At six Inches from the Ends are fix'd the fore-side and back-side strongly fasten'd to the Beams, and to the Floor, by several Screws and Nuts.

See Fig. 4 and 6. †

† Pl. 21. F. 4.

6. To these two Ends, when Occasion requires, may be fasten'd two Sides made of Planks 13 Feet long, which fit into the Side of the outward Beams by means of *Hooks and Rings*, and are kept up by means of *Latches* to be seen in Fig. 4, 5, 6. † These Sides are also further strengthen'd by a Chain going across in the middle of the Carriage.

and 6.

† Pl. 21. F. 4, 5, and 6.

7. At right Angles under these Beams, at a proper Distance (see Fig. 5.) are fasten'd two strong Timbers, by means of large Screws and Nuts.

8. In these Timbers well strengthen'd and plated with Iron, where the greatest Stress lies, are placed two semicylindrical Pieces of Brass at each End, to serve as a *Collar* for the Axel-trees of the Wheels, which being well greased, turn with very little Friction.

9. There is likewise under these four Beams (already mention'd) another piece of Timber of about 6 Inches by 4 well fasten'd to it, at right Angles, and at such a Distance, as is seen in Fig. 5.

This Piece serves as a fix'd Point to place a Leaver, which locks (or keeps from turning) the Hind-wheel, by pressing upon it.

10. The Axel-tree is about three Inches Diameter. See † Plate 22. Fig. † Pl. 22. F. 1.

1. One of its Ends is square, the other round, and on these two Ends, the Wheels are plac'd in an alternate Position; that is, the right-handed Fore-wheel is on a Square, and the left on a round part of the Axel-tree, whilst the right-handed Hind-wheel is on a round End, and the left on a square End of the Axel-tree; thereby any one of the Wheels may be lock'd separately; for when the Wheel placed on the round End is lock'd, the other, together with the Axel-tree, revolts within it, and when the Wheel which is fasten'd on the square is lock'd; the other revolves notwithstanding as usual upon the Axel-tree, which is then unmoveable.

11. These Wheels are made of cast Iron about 20 Inches Diameter, and have a *Flanch* 6 Inches broad next to the Carriage which hinders them from running off from the Oaken Frame on which they move. Their Plan and Profil are seen in Fig. † 5. of Plate 21, and Fig. 1. of Plate 22. and their Sections either thro' their Spokes (or Radii) or between any two of them, as † Fig. 3d and 4th of Plate 22.

† Pl. 21. F. 3

& Pl. 22 F. 1.

† Pl. 22. F. 3

12th. The manner of locking and unlocking the of the Wheels is as follows: When either of the Hind-wheels is to be lock'd, a strong Leaver (which they call the Jigg Pole) is placed on that End of the Timber (described in Paragraph 9.) next to the Wheel to be lock'd, and after it is pass'd thro' the Iron Loop to secure it the better, a Chain coming from the Roller to be seen in † Fig. 5 and Fig. 1 of Plate 22. is clapt over the Extremity, and by means of a short iron Bar, and the *Rocket* and *Click*, seen in Fig. 5 and 6 Plate 22, \* one of the Drivers or Persons that attend the Carriage in a very little time stops the Wheel either partly or intirely.


and 4.

† Pl. 21. F. 5

& P. 22. F. 1.

\* Pl. 22. F. 5,

and 6.

Annotat. To unlock which, 'tis only lifting up the End of the *Click* or *Catch* made  
 Lect. IV long on purpose, for then the Leaver pressing no longer, the Chain is slack-  
 ken'd, taken off, and the Jigg Pole laid in the Cart, till another Occasion  
 serves. They have two Jigg Poles, one for each Hind-wheel.

13. The Fore-wheels are lock'd, by means of a thick square iron Bolt,  
 † Pl. 21. F. 5. seen in *Fig. 5* of *Plate 21*. † coming out in the Direction of the Axel-tree,  
 between the Spokes or Radii of the Fore-wheels: these Bolts, are pro-  
 truded forwards to lock the Wheels, and drawn back to unlock them  
 separately, by means of a Contrivance, part of which is seen in *Fig. 5*.

† Pl. 21. F. 5 of *Pl. 21*. and 1st of *Pl. 22*. † Towards the middle of the back-side are  
 Pl. 22. F. 1. two iron Rods, (see one in *Fig. 6. Pl. 22.* \*) turning separately on the square

\* Pl. 22. F. 6. of an Axel. When either of these is brought by the Hand from a verti-  
 cal Position, in which they are drawn, to an horizontal one, the iron Rod,  
 of which it takes hold by its lower End, is pushed forward four or five  
 † Pl. 23. F. 1. Inches, and by means of the Contrivance in † *Fig. 1, 2, 3, 4*, of *Pl. 23*.  
 2, 3, 4. shoots the square Bolt between the Spokes or Radii.

When the Wheel must be unlock'd, this Bolt is drawn back into its for-  
 mer Position, by bringing the iron Rod into its former vertical Situation, as  
 † Pl. 22. F. 1. in *Fig. 1.* of *Pl. 22*. † As these Carriages are loaded with a considerable  
 (often with upwards of four Tons) Weight of Stone, when they come  
 down the Hill, all the Contrivances explain'd above would be usefess, and  
 that great Weight would sink the Carriage too deep into the Ground, with-  
 out pieces of Oak laid all along the Way which these Carriages then pass o-  
 ver. These are sufficiently seen in *Fig. 4, 5*, of *Pl. 21*. \* and *Fig. 1.* of *Pl. 22*.  
 \* Pl. 21. F. 4, 5. and Pl. 22. F. 1. Altho' these Carriages are very heavy even when empty, yet by means  
 of the Frame on which they move and the little Friction the Axel-trees  
 feel revolving in the brass Collars, two Horses not only draw up them the Hill  
 very easily when empty, but draw them along on the Plain when loaded, at  
 a very good Rate. As soon as the Carriages come to the Brow of the Hill,  
 the Horses are taken off, and one or more of the Wheels lock'd, by the  
 Driver, who stands behind to moderate the Motion as he thinks proper.

When the Carriages are come to the Water-side, and have been unload-  
 ed, they change the Horses from End to End, so that the part of the Car-  
 riage which went before descending, becomes the hind-part in ascending  
 the Hill, which avoids the Trouble of turning with these Wheels.

These Carriages are loaded at the Quarries, and unloaded at the Water-  
 side, by means of a very good and curiously contriv'd Crane, fully describ'd  
 by Dr. J. T. Desaguliers, in his *Course of Experimental Philosophy*, Lect. III.  
 Ann. 15.

N. B. Mr. Allen, to whom I am oblig'd for a full and thorough Sur-  
 vey of these Carriages, told me, that one of them, when compleatly fi-  
 nished, and ready to be used, stands him in upwards of 30 Pounds:  
 which I thought proper to mention as a very reasonable Price consider-  
 ing the good and workmanlike manner in which every thing is per-  
 formed.

Tho'

Tho' Mr. *Labelye's* Description is very intelligible, and his Draughts Annotat. extremely well done, yet, to make every thing still plainer, I have added Letters to his Draughts, and the following References. Lect. IV.

*References to the Figure of Mr. Allen's Carriages and their several Parts.*

*Plate 21. Figure 4.*

ABCDIIIIHFEG. The Body and Bottom of the Carriage, without the Sides, which are put on upon occasion in the Place BDHF and fastened by means of the Hooks *ggg*, and the Latches *e f*.

IIII. The cross Pieces on the Bottom to strengthen it.

MN. Strong Pieces of Timber under the Bottom.

K. A cross Piece under the Bottom having an Iron Loop at Top to receive the End of a Leaver that presses on the hind Wheel *L 2*, to stop it from turning round when the Motion is too rapid.

L, *L 1*, *L 2*. Three of the four Wheels, the fourth being out of sight in this Position of the Carriage, whose Circumferences have a Flanch on the inside that the other part may bear on the Timbers or Waggon-way.

H. An Iron Roller for the Chain to wind on to hold down the stopping Leaver, as it presses on the hind Wheel.

O, *O 4*, *O 1*, &c. The Waggon-way, or parallel Timbers laid with a Descent for the Carriages to run down by their own Weight.

*Figure 5.*

FHDB. The right Side of one of the Carriages fix'd by means of the Hooks at *gggg*, and the Latches *e, f* shewn in the fourth Figure.

L. A Fore-wheel with a round Hole in the Nave to receive a round End of the Axis that goes thro' the Piece of Timber *P*, from another part of which the Bolt *p* is shot between the Spokes to stop the Wheel from turning round, when the Motion is to be retarded.

H *p*. An Iron Rod push'd forwards from behind to bolt or lock the Fore-wheel abovemention'd.

*L 2*. A Hind-wheel fix'd upon, and turning round with the Axis coming thro' the Piece of Timber *Q*, the end of which Axis is made square for that Purpose.

R K. A Leaver, whose End goes thro' an Iron Eye on the Timber *K*, having there its Center of Motion, with the Compass-piece *qq* to press on the upper part of the Wheel *L 2*, to stop it upon occasion, from turning with the Axis.

H. A Roller on which is wound the Chain HR which pulls down the End of the Leaver at R and keeps it in its place, to press hard upon the Wheel at *qq*.

*O 2*, *O 1*. The Waggon-way which supports the strong part of the Circumference of the Wheel, while the Flanch or larger Circumference of

Annotat. each Wheel falls on the inside of the Timber, that the Carriage may  
 Lect. IV. not jump or run out of the Way.

*Plate 22. Figure 1.*

This Figure drawn by a larger Scale (*viz.* of 10 Inches to an Inch) shews the Elevation of the hind-part of one of the Carriages, with the Profil of the Wheels, &c.

FHGE. The End of the Carriage behind bearing upon four Timbers, whose Ends are seen here.

*f g, e g.* The Ends of the Sides hook'd on at *g, g*, and latch'd at *f* and *e*.

*H b r i*, and *G b r i*. A Roller with its Winch, and Ratchet and Latch, to receive the Chain that draws down the Leaver or Jig-pole. The Chain goes on the Part *H*, or *G*, the Holes to turn the Axis of the Roller with an Iron Hand-spike are seen at *h h*, the Ratchet at *i i*, and its Catch at *r, r*.

*l k, l k*. Two perpendicular Iron Bars, whose lower Ends *k k* shoot forward each an horizontal Bar (not represented here, but shewn in the last describ'd Figure at *H p*) to bolt either of the Fore-wheels singly, or both at once.

*L 2, L 4*. The Sections of the two Hind-wheels, with their Flanches *m m, m m*, and bearing Parts *n n n, n n n*; the left Wheel receives the square End of the Axis *Q* in a square Hole so as to turn with it, and the right Wheel has a round Hole to receive the End of the Axle at *P* which is round also; so that this Wheel can turn round without turning the Axle *P S Q* along with it.

*O 3, O 4*. Shews the Section of the Timbers or Waggon-way; where may be seen the manner of the Wheels bearing on the Timbers at *n n, n n*, whilst the Flanches *m m, m m*, come down to keep the Carriage in its Place.

*Q 1, P 1*. The cross Timbers thro' the Bottoms of which the Iron Axes of the Carriages pass, which are fix'd up under the Carriages by Pins and Nuts here represented by prick'd Lines.

*Figure 2.*

Represents one of the Iron Axle-trees *S*, whose end *Q* at the right Hand is square, the other end at the left Hand being round, with an Hole for the Linch Pin at each End.

*Figure 3.*

Shews the Section of the Hind-wheel on the left Side, or the Fore-wheel on the right Side, with a square Hole *Q*, the Rim of Circumference of the Wheel *n n*, and the Flanch *m m*.

*Figure 4.*

Shews the right Wheel behind or left Wheel before, mark'd with the same Letters except *P*, which shews the round Hole to receive the round end of the Axis.

*Figure*

Figure 5.

Shews the Catch *r*, and the Ratchet *i*.


Annotat.  
Lect. IV.  


Figure 6.

Shews one of the perpendicular Rods of the sixth Figure, whose Handle is at *l* and the bottom or open End is to join to one of the horizontal Bars which shoots a Bolt in between the Spokes of one of the Fore-wheels.

Plate 23, Figure 1.

Pl. 23. F. 1.

This Figure represents the upper part of the Timber under the Carriage between the Fore-wheels, thro' the lower part of whose Ends the Axel-tree passes, and in the Substance of which is let in the Machinery for stopping the Fore-wheels, where the two Bolts AB and CD are seen, which may be separately shot out to the right or left Hand thro' a square Staple P or Q, and resting on one of the cross Pins *e f* EF or GH, *g h*, by means of the horizontal Bars IK already describ'd, one of which is to be seen in the 5th Figure of Plate 21, mark'd H *p*, and the Ends of both of them are seen in this Figure. The Bolt AB on one Side is represented as shot out between the Spokes of the Wheel, and the other is in its usual Place, where it does not touch the Wheel. N. B. Pulling back the Bar LI unbolts BA, by means of the Elbow LNB, and pushing forwards the Bar KM bolts CD, by means of the Elbow MNC.

Figure 2.

Pl. 23. F. 2.

The second Figure shews an End of the Timber or the Section of the Machinery to move the Bolts as cut across one of the Bolts and at right Angles to the Axis at A, where is seen the Hole P over the Axis, and the prick'd Lines EF shews the Pin or Shaft of the Screw on which the Bolt slides as it comes out of its Staple.

Figure 3.

Pl. 23. F. 3.

The third Figure represents the rectangular Elbow-Piece, such as BLN or CNM of the last Figure, where is seen the Center of Motion round the Pin N, the End L receiving the protruding End of the horizontal Bar, and the End B receives the End of the Bolt to throw it forward as at B Fig. 2, or draw it back as at C in the same Figure.

Figure 4.

Pl. 23. F. 4.

The fourth Figure represents the Iron-work about the End of the Timber at III with the Holes for the cross Pins, the Bolt BA and Staple P.

*Before I begin my next Lecture, it may not be improper to give an Account of a Man of very great natural Strength who lives now here in London, and shews several surprizing Effects of his Strength. I should indeed have given an Account of him in the 7th Annotation; but I was unwilling to do it upon common Report, till I had seen him my self, which I only did since the last Sheet was printed off.*

Thomas

Annotat. *Thomas Topham*, born in *London*, and now about 31 Years of Age, five  
 Lect. IV. Foot ten Inches high, with Muscles very hard and prominent, was brought  
 up a *Carpenter*, which Trade he practised till within these six or seven  
 Years that he has shew'd Feats of Strength; but he is intirely ignorant of  
 any Art to make his Strength appear more surprizing: Nay, sometimes he  
 does Things which become more difficult by his disadvantageous Situation;  
 attempting, and often doing, what he hears other strong Men have done,  
 without making use of the same Advantages.

About six Years ago he pull'd against an Horse, sitting upon the Ground  
 with his Feet against two Stumps driven into the Ground, but without the  
 Advantages represented by the *first Figure, Plate 19*; for the Horse pull-  
 ing against him drew upwards at a considerable Angle, such is represented  
 in the 2d Figure of that Plate, when  $hN$  is the Line of Traction, which  
 makes the Angle of the Traction to be  $NhL$ : and in this Case his  
 Strength was no farther employ'd than to keep his Legs and Thighs  
 strait, so as to make them act like the long Arm of a bended Leaver, re-  
 \* Pl. 19. F. 2. presented by  $Lh^*$ , on whose End  $h$  the Trunk of his Body rested as a  
 Weight, against which the Horse drew, applying his Power at right Angles  
 to the End  $l$  of the short Arm of the said Leaver, the Center of Motion  
 being at  $L$  at the Bottom of the Stumps  $l, o$  (for to draw obliquely by a  
 Rope fasten'd at  $h$  is the same as to draw by an Arm of a Leaver at  $lL$ ,  
 because  $lL$  is a Line drawn perpendicularly from the Center of Motion  
 † Ann. 5. to to the Line of Direction †  $hN$ ) and the Horse not being strong enough to  
 Lect. III. p. raise the Man's Weight with such disadvantage, he thought he was in the  
 142. right Posture for drawing against an Horse; but when in the same Posture  
 he attempted to draw against two Horses, he was pull'd out of his Place by  
 being lifted up, and had one of his Knees struck against the Stumps, which  
 shatter'd it so, that even to this Day, the *Patella* or Knee-pan is so loose,  
 that the Ligaments of it seem either to be broken or quite relax'd, which  
 has taken away most of the Strength of that Leg.

But if he had sat upon such a Frame as is represented in the first Fi-  
 \* Pl. 19. F. 1. gure \* and describ'd in Page 256, 258 and 259, he might (considering his  
 Strength) have kept his Situation against the pulling of four strong Horses  
 without the least Inconvenience.

*The Feats, which I saw him perform a few Days ago, were the following.*

1. By the Strength of his Fingers (only rubb'd in Coal-ashes to keep them from slipping) he roll'd up a very strong and large Pewter-dish.
2. He broke seven or eight short and strong pieces of Tobacco-pipe with the Force of his middle Finger, having laid them on the first and third Finger.
3. Having thrust in under his Garter the Bowl of a strong Tobacco-pipe, his Legs being bent, he broke it to pieces by the Tendons of his Hams, without altering the bending of his Leg.
4. He broke such another Bowl between his first and second Finger, by pressing his Fingers together side-ways.
5. He

5. He lifted a Table six Foot long, which had half a hundred Weight Annotat. hanging at the End of it, with his Teeth, and held it in an horizontal Po-Lect. IV. sition for a considerable time. *It is true the Feet of the Table rested against his Knees; but as the Length of the Table was much greater than its Height, that Performance requir'd a great Strength to be exerted by the Muscles of his Loyns, those of his Neck, the Masseter and Temporal (Muscles of the Jaws) besides a good set of Teeth.*

6. He took an Iron Kitchin-poker, about a Yard long, and three Inches in Circumference, and holding it in his right Hand he struck upon his bare left Arm, between the Elbow and the Wrist till he bent the Poker nearly to a right Angle.

7. He took such another Poker, and holding the Ends of it in his Hands, and the Middle against the Back of his Neck, he brought both Ends of it together before him; and, what was yet more difficult, he pull'd it almost streight again; because the Muscles which separate the Arms horizontally from each other are not so strong as those that bring them together.

8. He broke a Rope of about two Inches in Circumference which was in part wound about a Cylinder of four Inches Diameter, having fasten'd the other End of it to Straps that went over his Shoulders; but he exerted more Force to do this than any other of his Feats, from his awkwardness in going about it; for the Rope yielded and stretch'd as he stood upon the Cylinder so, that when the Extensors of the Legs and Thighs had done their Office in bringing his Legs and Thighs strait, he was forc'd to raise his Heels from their Bearing, and use other Muscles that are weaker. But if the Rope had been so fix'd that the Part to be broken had been short (in the Manner explain'd in the 7th Annotation of this Lecture) it would have been broken with four times less Difficulty.

9. I have seen him lift a rolling Stone of about 800 lb with his Hands, only standing in a Frame above it, and taking hold of a Chain that was fasten'd to it. By this, I reckon he may be almost as strong again as those who are generally reckon'd the strongest Men, they generally lifting no more than 400 lb in that manner. The weakest Men, who are in Health and not too fat, lift about 125 lb, having about half the Strength of the strongest. N. B. This sort of Comparison is chiefly in relation to the Muscles of the Loyns; because in doing this one must stoop forwards a little. We must also add the Weight of the Body to the Weight lifted. So that if the weakest Man's Body weighs 150 lb, that added to 125 lb makes the whole Weight lifted by him to be 275 lb: Then if the stronger Man's Body weighs also 150 lb, the whole Weight lifted by him will be 550 lb, that is, 400 lb, and the 150 lb which his Body weighs. Topham weighs about 200 lb which added to the 800 lb that he lifts, makes 1000 lb; but he ought to lift 900 lb, besides the Weight of his Body, to be as strong again as a Man of 150 lb Weight who can lift 400 lb.

Now as all Men are not proportionably strong in every Part, but some are strongest in the Arms, some in the Legs, and others in the Back, according to the Work and Exercise which they use, we can't judge of a Man's

Annotat. Man's Strength by lifting only; but a Method may be found to compare together the Strength of different Men in the same Parts, and that too without straining the Persons who try the Experiment.

*Here follows the manner of doing it, which was communicated to me by Richard Graham, Esq; F. R. S. to which I made some small Additions.*

*Plate 23, Figure 5.*

ABCD is a strong Frame of Wood with an Hole thro' the upright Piece DC; at D big enough to receive an Iron cylindrick Bar of an Inch Diameter or something bigger, a strong Iron Plate being fix'd on each Side, that the Iron may not gull the Hole. This Bar has a Square upon it, whose Side is about an Inch and one eighth to receive the two separate and unequal Arms of a bended Leaver DF and DE; and then a strong Nut *d* is screw'd over them at D to keep all tight. The Arm DE, which as a Stilyard is to carry a great Weight W, is kept from falling below an horizontal Position by an Iron Pin at K, which stops the short Arm DE from inclining towards G; but both Arms are moveable round the Axis D towards *e* or N. The Arm DF has a round cross Bar at top about six Inches long, as may be seen in its separate Representation *d f*. To the upright Timber AB is made fast the Iron LN, with a Cross also at Top (See *n l*) and Holes for Iron Pins to fasten it in its Place. There is also another strong piece of Iron HGI fasten'd to the Timber that carries the Leaver by a strong Wood Screw at I, and the Pin K going thro' its Wings and the Timber. See its separate Figure at *h g I*. S is a Collar to put on when you don't use the upright Arm of the Leaver. M is the Center of Gravity of the Stilyard DE.

1. To try one's Strength by means of the Machine; with the left Hand take hold of the round part of the Cross at N, and of the round part of the Cross at F with the right Hand, and bring your right Hand towards the left in the Direction of FN which will move DE and raise the Weight W. When you can just raise it up so as to make FD quit the Pin at K, the Force of the Arms will thus be found. Multiply the Weight W (suppose an half hundred, or 56 lb) by its Distance from the Center WD (which we will suppose here 15 Inches) which will give 840 for the *Momentum* of W on the Stilyard: Then add to it the *Momentum* of the Stilyard it self, which you will have by finding what Weight can draw up the Stilyard by its Center of Gravity, namely the Weight W (*Fig. 6.*) which draws it up by a String going over the Pulley C; and multiplying that Weight by MD its Distance from the Center (which we will here suppose 10 Inches) you will have 60, which being added to 840 make the Sum 900, and that Sum, divided by FD the Distance of the Power, will give 90 lb for the Force of the Man's Arms who applies his Hands at F and N. If another Man raises double the Weight at W + so much Weight as will answer to double the Weight of the Stilyard at M, he is twice as strong

strong. The removal of the Weight *W* toward *E* will also serve to shew Annotat.  
how much the Force is greater, instead of adding Weight at *W*. † Lect. IV.

The same way may be found what Force the Arms can exert in pulling from each other, by applying one Hand at *F* and the other at *H*. † Pl. 23. F. 5.  
And to try the Force with which *Topham* bent the Poker bearing behind his Neck; a Man must put a Strap round his Neck, which must be fasten'd to *F*; then the Head being plac'd on the Side *N* of *NL*, with both his Hands he must take hold of the Cross at *N* and push forwards with his Hands, all the while pulling backwards with his Neck, to bring *F* towards *N*.

2. The sixth Figure is another wooden Frame with the upright Timbers *AB, CO* a little farther asunder. \* So that a Man may stand upon the strong † Pl. 23. F. 6.  
Plank *FG*, and with the Girdle and Chain thro' the Hole *H* pull up the Stil-yard *DE* by the Hook *I*, which Stil-yard in this Case has not the upright Piece *FD*, but instead of a bended Leaver becomes a Leaver of a third kind, where *DI* is the Distance of the Power, and *DW* the Distance of the Weight; and therefore  $\frac{W \times WD + w \times MD}{DI}$  will be the absolute

Weight that is lifted, or Force of the Muscles extending the Legs. But here the Weight of the Trunk of the Body must be added, and consider'd according as some Men are heavier than others. N. B. *The Notch K kinders the Leaver from falling below the horizontal Situation; and w drawing over the Pulley C shews how the Leaver acts at its Center of Gravity: So that when w is hung on, the Leaver is to be look'd upon as an Instrument without Weight.*

† Pl. 23. F. 7.

3. The seventh Figure † has the Stil-yard at *DE* with a wooden Cylinder of an Inch and an half, or two Inches in Diameter, upon its Axis continued behind at *DF*, which Cylinder is drawn separately at *df*, and its Iron Axis at *gh*, and the Nut at *i*.

† Pl. 23. F. 8.

4. The eighth Figure † represents a Machine to try the Strength of the Fingers; in which, when you thrust the Fore-finger under *G*, and the third Finger under *H*, which are fix'd Points, the middle Finger may push upon *N* to lift a Weight at *W*, in the same manner that *Topham* breaks the pieces of Tobacco-pipes. If the Hand being held with the Palm upwards, the two first and two last Fingers be thrust in under *G* and *H*, and the Thumb presses upon *N*, that will shew in what manner the tops of Pewter Pots and Silver Tumblers have been squeez'd together by Men very strong in the Fingers; and thus will be shewn the Strength that any Man can exert for such an Action, &c.



## LECTURE V.

*Concerning Sir Isaac Newton's three Laws of Motion.*

I. **T**HERE are several Organs or Instruments, which may be call'd Mechanical, or enumerated among the Instruments, commonly, but erroneously, call'd *mechanical Powers*; which I purposely omitted (or only mention'd without explaining their manner of acting) in the third Lecture; because the Knowledge of the Laws of Motion is necessary for understanding upon what Principle they act: And such are the *battering Ram*, the *Hammer* or *Mall*, the *Fly*, the *circular Pendulum*, the *Sling*, and the *Bow* or *Spring*.

I SHALL therefore in the first place consider those Laws, and draw several Corollaries or Consequences from them, which I shall illustrate by Experiments, and apply them among other Things to explain the Use of those Instruments.

### LAW I.

*Every Body perseveres in a State of Rest, or of uniform Motion in a right Line, unless it be compell'd to change that State by Forces impress'd thereon.*

2. THERE is in all Matter (whatever kind of Body it be shap'd into) an Inactivity whereby it resists any Force that endeavours to make it change its State, in proportion to its Quantity of Matter; and this is call'd the *Vis Inertiae* or Force of Inactivity. For it is as impossible for a Body to go into Motion of it self from Rest, as to change its Shape from one Figure to another. This is evident to Sense, and I believe no body doubts of it; but the second part of this Law does not appear so evident without a little Attention.

WE

WE see plainly that there must be some extrinfecal Agent or Power not essential or belonging of necessity to the Body, to put it into Motion: but we don't so readily perceive that a Body in Motion would continue to move for ever without the action of an extrinfecal Agent; because we see Bodies here on Earth gradually lose their Motions, and for want of attending to all the Causes that destroy the Motion of Bodies, we often imagine that Motion languishes and at last quite perishes of it self. But if we consider what external Causes retard and destroy Motion, we shall soon perceive that if those Causes were remov'd, a Body once put into Motion in any Direction wou'd continue in that Motion and Direction for ever.

Lea. V.  
~

A STONE thrown forwards with the Hand goes on with the Motion that it has received from the Hand, and wou'd continue in that State of Motion for ever, if there was neither Air nor Gravity. When we consider the Resistance of the Air, it is evident that the Body in going forwards must remove the Parts of the Air to make it self way; and as it must communicate Motion to those Parts which it removes; so much Motion as it communicates, so much it must lose; so that after some time and having gone through a certain Space it must hang in the Air immoveable, that is, if nothing but the Air acted upon it. But besides the Air, Gravity (which is a Force pushing downwards) alters its Direction and brings it gradually to the Ground. This Endeavour which every body has to continue in its State of Motion does not seem to some to be properly call'd a Force of Inactivity (*Vis inertiae*) but when we consider that the Body is purely passive, neither augmenting nor throwing off any of its Motion of it self, we shall find it to be entirely inactive even in that State. Thus when we stand by a River Side and observe a live Fish carried down the Stream, he is wholly inactive in that Motion, and continues in that Motion all the while he is inactive: it must be an Action exerted (equal to the Force of the Water) that will make him come to Rest and appear so in respect of the Spectator on the River Side.

SUPPOSE a Man exerts a certain Force to roll a Bowl on a Bowling-green neither mow'd nor roll'd, and can only throw it 20 Yards with that Force; when the Green is mow'd, he shall with the same Force throw it farther, 30 Yards for Example; then if the Green be roll'd as well as mow'd, the same Force may throw the Bowl (for Example) 40 Yards; and still if more of the Obstacles are taken away,

~~~~~  
 Lect. V. the Bowl will yet go farther; whence one may easily conclude, that if the Plane on which the Bowl rolls could be made perfectly smooth and mathematically true, the Bowl truly spherical, and the Resistance of the Air wholly taken away, the Bowl would for ever roll (or rather slide) on that Plane if it was infinitely extended.

2. WE have shewn by a Quotation from Sir *Isaac Newton* (L. 3. N^o. 85.) that if a Body be acted upon by two Forces whose Directions make any Angle, the Body will move in the Diagonal of a Parallelogram, two contiguous Sides of which represent (by their Lengths) the respective Quantity of those Forces, and (by their Inclination) their Directions: and also that the Body will go thro' the whole Diagonal by the Action of the said two Forces, in the same Time that it wou'd have gone thro' either of the contiguous Sides, if only one of the Forces had acted upon it.

THIS is not so readily conceiv'd by those who are not accusom'd to mathematical Reasoning; because, while they observe the Action of one of the Forces, they do not attend to the other: but it will appear very evident, by considering such Cases wherein the Space in which the Body moves is carried in a different Direction from that which the Body appears more immediately to be mov'd in. As for Example, let us suppose TS (* Pl. 23. Fig. 10.) to be a *Tragschuit* or *Dutch* travelling Boat (I mean the Plan of it) Hh g G, the Canal in which the Boat goes in the Direction TS, and A and B two Persons sitting over against one another in the Boat. Now let us suppose the Person at A tossing any Body, as for Example, a Ball, to the Person at B, and so reciprocally: all the People in the Boat will consider the Ball so toss'd as only moving in the Line AB, whether the Boat stands still or goes on along the Canal, tho' there be but one Force acting upon the Ball in the Direction AB when the Boat stands still, whereby it really moves in that Line; but when the Boat goes along, another Force in the Direction Aa does also at the same Time act upon the Body, which by that compound Action is really carried in the Line AB, tho' the Persons sitting by fancy the Ball to go still in the Line Ab, because they are carried along too, and forget that the Force which draws the whole Boat does also carry the Ball along, which is a part of it, as well as themselves. Now this will be evident to the Sight of a Man that stands upon the Bank at C; for when the Boat stands still he looking towards D on the opposite Bank, sees the Ball move (as it really does) in the Line
 AB.

A B. But when the Boat moves and goes from the Position T S Lect. V. to the Position ts in the same time that the Ball moves across from the Person at A to the Person at B, it is plain (and will be seen by a Man on the Bank at C) that the Ball moves in the Diagonal Line A b , because the Person who was at B does not receive it till he is carried to b by the Motion of the Boat; which his Opposite (being at the same time carried to a) does not attend to, tho' the Thing be so visible to the Man at C. Yet if the Man at C stood upon a Plank at A C fasten'd to the Boat, being then carried along from C to c , he would not distinguish between the Line A B and the Line A b , but (when he came to c) suppose that the Body had mov'd in the Line ab , which he would mistake for A B; unless looking over the Canal, he shou'd consider that he no longer saw the Point D but the Point d . Thus in a Ship, if a Man standing on the Deck near the Mast tosses up a Stone to another who stands on the round Top, or receives a Stone dropp'd from thence, the Stone will move parallel to the Mast (supposing the Mast upright) whether the Ship be at Anchor or under Sail; tho' in the first Case the Stone moves perpendicular to the Horizon, and in the last Case oblique to it as it rises or falls in the Diagonal of a Parallelogram, whose contiguous Sides represent the two Forces acting upon the Body, the one perpendicular, and the other parallel to the Horizon; just as in the last Instance, the two Forces A B and A a acted upon the Ball horizontally projected in the Boat.


N. B. *Whether the Boat goes faster or slower (in which Case A b becomes longer or shorter) the Ball will perform its diagonal Motion in the same Time; for since the Force A B continues the same, all that is done by the lateral Force A a is only to make the Ball come to a different Point of the Line B n as m or b or n instead of B, while the Force A B in the same Time carries it cross the Boat, or from some part of the Line A a to some part of the Line B b .*

I HAVE been the more prolix in this Explication, because I have met with a great many Persons of good Sense, who for want of Attention have not been able to conceive this compound Motion, on which all Mechanicks are founded. For this Reason I have subjoin'd variety of Experiments to the same Purposes, because if one don't, another may, convince.

EXPERIMENT I. *Pl. 23. Fig. II.*

Pl. 23. F. II. THE Machine of Plate 23. Fig. II. consists of a brass Plate ABCD, upon which another Plate IKLM slides forwards and backwards in the Direction TS or ST between the Rulers AD and BC under the Edge of two other Rulers EF and GH, of which EF is a little Rack with Teeth to receive the Teeth of the Wheel N, which is fasten'd to the Plate IMKL by the Cock NO, and turns round its Axis N as the said Plate is push'd forwards and backwards. The Wheel in its Motion along EF carries another Rack PQ at right Angles to the former either to the right or to the left, according as the Axis of the Wheel at N is push'd in the Direction TS or ST. This last Rack has an Arm SR with a little Socket at R to receive a black Lead Pencil, which according to the Motion of the End R will draw a black Line upon the Paper on which the Machine is to be laid.

THERE must be drawn upon the Paper a Square $efgh$, one of whose Sides is equal to the Distance between the Edges of the Rulers AD and BC. Then having push'd up the Wheel N as far as it will go, lay the Machine upon the Paper, so that its End AB may be parallel to gh one side of the Square abovemention'd, at such a Distance as to have the Point of the Pencil fall upon the Point e , and the Edge of the Rack PQ over the Line ab : then with one Hand preffing down the Plate ABCD of the Machine at T, to keep it immoveable upon the Paper, pull down the Cock NO in the Direction ST a Length equal to eg ; and the Pencil, instead of describing the Line eg , will describe the Diagonal Line eb , because as the Wheel in its Descent is turn'd round by the Rack EF, it throws the Rack PQ to the right, just the length of ef , by means of the Teeth in the said Rack, and the Slit in it which lets it go laterally along the Pins IM by the Action of the Wheel, while the said Pins bring it perpendicularly down from the Line ab to the Line de . And to shew that the Diagonal is describ'd in the same time that the Side eg wou'd have been describ'd if there had been no lateral Motion; take off the Wheel, and then the Pencil's Point will describe the Line eg , while the Cock is drawn down the same Length as before, or push'd up again. This shews that it is the same thing for a Body at e to be acted upon by two Forces in the Directions ef and eg , as to be carried from the Line eg to the Line fb ,

fb, at the same time that it is brought down from the Line *ef* to Lect. V. the Line *gb*. 

EXPERIMENT II. *Pl. 23. Fig. 12.*

THE 12th Figure of Plate 23 is a square Frame of Wood, of *Pl. 23. F. 12.* which the part *CDBE* is to be set up on its Side *BE*, whilst the other part of it *cdeb* is drawn out of its Groove into the Position *dd ee*; by which means the Ball *A* is made to describe the Diagonal *Aa* ascending, and the same Diagonal descending is describ'd when the sliding part is push'd back into the Frame in the Direction *sd*. There is a Wire *Ac* fix'd to the sliding part for the Ball to move up and down on, and a String *bc* which goes thro' an Hole in the Pieces *dc* and *DC*, so as to have its End fasten'd at *C*; by which means the Ball must rise when *dd ee* is drawn out laterally; and it falls back again by its own Weight when *dd ee* is push'd in again: and when *dd ee* is left in its containing Frame, the Ball will rise and fall by pulling or letting go the String *Ccb*. If the Wire *AB* be consider'd as a Rope parallel to the Mast of a Ship, what we have said of a Body let fall from the round Top of the Mast and moving parallel to the Mast (whether the Ship be at Anchor or under Sail) by one, or two Forces acting upon it, will be illustrated.

BUT as I have heard some object, that this, and the last mention'd Machine, did indeed, by the Contrivance of them, make a Body move in a Diagonal; but they did not prove that Nature acted that way: I have made the Thing evident to Sight by the following

EXPERIMENT III. *Pl. 23. Fig. 13.*

THE round Table *GEFD* stands upon a Pivot *P*, and has an *Pl. 23. F. 13.* Hole on each Side of its Center half way from the Circumference, at *A* and *B*. At right Angles to the Line that goes thro' *ACB* is drawn a Diameter *DE*, and from several Points of that Diameter equally distant from the Center on each Side, are drawn the Rhombs or equilateral Parallelograms, *DAEB*, *dAeB*, and *saB*. The Ball *D* has two Strings fasten'd to it at the same Point, and those Strings being carried thro' the Hole *AB*, have hanging at their End two equal Weights *W* and *X*, which by the joint Force of

Lect. V. of their Descent will bring the Ball D to rest upon C the Center of the Table, whilst the parts of the Strings on the Table are in the Line ACB. If the Ball D be pull'd back to δ, d , or D (whereby the Weights W and X will be rais'd up, and the Strings will lie upon and have their Directions in the contiguous Sides of one of the Parallelograms) as soon as you let it go again, it will run in the Line DE which is the common Diagonal of all the Parallelograms; as may be best perceiv'd by holding one's Finger or a Wire at the Center C, which the Ball will strike in its Motion. *Here it is plain that the natural Gravity of the Bodies W and X is the only Force (divided into two equal Parts) which acts upon the Ball D, and that whether the Directions of those Forces be at acute, right, or obtuse Angles, the Ball is still carried in the Diagonal of a Parallelogram of equal Sides, representing equal Forces. But if one of the Weights as W, be taken as heavy again as the other, the Ball from D, d, or δ , will run in a Direction between A and C; and not run over the Center till you let it go from F, in which Case the two Strings will make two contiguous Sides of the Parallelogram FAGB, whose Proportion to each other is also as Two and One; the Diagonal being FCG, which passes thro' the Center.*


It was objected to Galileo when he asserted the Motion of the Earth, that if the Earth did turn round from *West* to *East*, a Cannon Ball shot upright, would not fall down in, or near the Place from which it was shot (as we see by Experience it does) but move to the *West* according to the space of Earth that had been carried towards the *East*, during the Rise and Fall of the Ball; which he answer'd, by saying, that the Ball is acted upon by two Forces, One, that of the Powder which throws it up, and the Other, that of the Earth which carries it to the *East*, making it in its Rise describe a Diagonal Line towards the *East*, and from the upper End of that Diagonal, another Diagonal Line also towards the *East* in its Fall. He alledg'd the Case of the Ship that we have already mention'd. But nothing will make the Thing so plain as the following,

EXPERIMENT IV. *Pl. 24. Fig. 2.*

To an horizontal Piece of Wood GH, fix'd about 10 Foot above the Floor of my Room, I screw'd the two Hooks S s, and by means of four Strings hung on the said Hooks, the brass Plate or flat Pendulum ABCD, which (by reason of the Distance of the Strings at their Points of Suspension S s) perform'd its Vibrations in a circular Arc over the graduated upright wooden Plane EF which stood under it, the Center of the Pendulum never moving out of the Plane of the Wood as it vibrated in the Direction E, or back again, at the Distance of about nine Foot from the Piece GH. Over the two Pullies I, K, screw'd into the upper Side of the Piece GH, ran the String LKI W, whereby the Weight W might be let down upon the Pendulum ABCD, or lifted off from it at pleasure; there being an Hole thro' GH, just under the Circumference of the Groove of the Pulley I.


Let the Pendulum (the Weight W being upon it) be drawn out of the Perpendicular to E, and then, when it is loos'd, it will vibrate towards F, and so back again towards E, &c. for a considerable Time. During that Motion, if by pulling the String at L the Weight W be lifted off of the swinging Plate, and then let down again, it will always fall upon the same Part of ABCD, tho' ABCD be mov'd a great way out of its Place during the Time that the Weight W is off of it; and that, at whatever Part of the Vibration W is lifted off, and at whatever Part of it, it is receiv'd again. For tho' the Force of the Hand pulling the String is the Cause of the Rise of W, and Gravity the Cause of its Descent, yet it has also another Force acting upon it, which moves it in the Arc of Vibration, as Part of the Pendulum compounded of ABCD and W. Just so as any Part of the Earth, on which a Cannon is set upright, is carried from *West* to *East*, by the Motion of the Earth round its Axis; the Cannon Ball shot up, besides the Motion receiv'd from the Explosion of the Powder, has another Force impress'd upon it by the Earth's Motion, as it is a Part of the Earth, and is carried from *West* to *East* in the Air, as well as the Cannon is upon the Earth, so as to fall down again into, or near, the Mouth of the Cannon.

A CANNON Ball or any other Projectile never describes a right Line in its Motion, but when it is thrown perpendicularly up or
P p down,


 Lect. V. down, which right Line of Ascent or Descent is the Diagonal of a Parallelogram, as we have already shewn; but *when the Body projected is thrown horizontally or obliquely, it describes a Curve call'd a parabolick Line.* Now this Curve is made up of an infinite Number of small Diagonal Lines, that continually change as the Directions of the Forces alter in the Body's compound Motion, which is easily explain'd from what has been said of the Action of two Forces, and the Consideration of the Acceleration of Motion in the Fall of Bodies; which (as it is a Fact and Matter of Observation) must be now taken for granted; tho' we shan't explain the Cause of it, but in the Explication of the second Law of Nature.

*Pl. 22. F. 4. LET a Body at A (*Pl. 24. Fig. 2.) as for Example, a Cannon Ball, be shot forward in an horizontal Direction A F, so as to move with a determinate Velocity; for Example 4 Rods in a Second of Time, or from A to B. It is evident by the First Law, that it must the next Second go on in the same Direction thro' the same Space, *viz.* from B to C, and so to D, then to E, then to F, &c. It is also plain, that if there be another Force whose Direction is at right Angles to the Line A F, as in the Line A L, which is perpendicular to the Surface of the Earth, (or to take the thing more generally, a Force pressing from the Line A F towards the Line L f,) the Action of this last Force upon the Body setting out at A will neither accelerate nor retard its Motion from the Line A L to the Line B M, nor from the Line B M to the Line C N, nor from the Line C N to the Line D O, nor from the Line D O to the Line E P, nor from the Line E P to the Line F f; but only cause the Body to go thro' different Points of those Lines instead of the Points B, C, D, E, F. Now I say, that *b, c, d, e* and *f*, will be the Points of the Lines B M, C N, D O, E P and F f, which the Body will go thro'. The Force pressing downwards in the Direction A L is the Force of Gravity, by whose Action it is observ'd that Bodies fall with an accelerated Motion, in such Manner that in the first Second of Time a Body falls 1 Rod (or 16½ Feet,) the 2d Second, 3 Rods; the 3d Second, 5 Rods; the 4th, 9 Rods, &c. which Spaces are here express'd by A G, G H, H I, I K and K L. And if the falling Body, during the Time of its Fall, be acted upon by another Force in an horizontal Direction, as in the Direction A F, (or rather from the Line A L towards the Line F f) that Force will not at all disturb the uniform Acceleration of the Motion of the

the

the descending Body, but only make it fall to some other Point of Lect. V. the Line Lf (or of the Surface of the Earth) than the Point L , as  for Example to the Point f .

The several Points of the Curve in which the Body moves will be thus found. The Body at A is acted upon by the two Forces AB and AG ; the Lines Gb and Bb , respectively equal and parallel to AB and AG , compleat the Parallelogram Ab : consequently the Body must move in the Diagonal Ab . The Body at b is acted upon horizontally by the Force bk equal to AB (or brought down from B C) and at the same time by the Force bb increas'd by Acceleration from 1 to 3, and transfer'd from GH to bb ; for as without the horizontal Force, it would in the 2d Second of Time have carried the Body from G to H (as it is not alter'd in the Quantity of its Effect by the horizontal Force) it will therefore be able from b to carry down the Body to b ; but the horizontal Force bk acting at the same Time, does in the same Time carry the Body from the Line bb to the Line kc , as the Force bb carries it from the Line bk to the Line bc , and consequently at the End of the 2d Second of Time the Body will be at c where the two above-mentioned Lines meet, having describ'd the Diagonal bc by the joint Action of the Forces bk and bb . The Body at c is acted upon by the horizontal Force cd , ($= CD$) and the perpendicular Force ci (so increas'd as to be $= HI$) at the same Time; therefore the Body will go in the Diagonal cd . For the same Reason, when the Body is at d it will go in the Diagonal de of the Parallelogram $deez$, by the joint Action of the Forces $d\epsilon$ and dx ; and so from e it will go to f in a new Diagonal, *viz.* that of the Parallelogram $e\phi fl$, by the Action of the two Forces $e\phi$ and el , &c.

S C H O L I U M.

IF we had divided the Time of the whole Motion into more, and therefore less, parts; we should have had more Diagonals from A , the Beginning of the Fall of the projected Body, to f the End of it, which would have made its way $Abcdef$ more curve by having more changes of Direction in it. Now as the Motion of Bodies downwards by the Force of Gravity is continually increasing (or the Line AG continually becoming longer) and not increasing by Fits, as we have suppos'd it to make: the Action of the two Forces the better

Lect. V. conceiv'd, there is a continual Change of Direction in the Motion of the Body forwards and downwards (or the Diagonals become infinitely small) so that it describes a Parabola in going from A to f.

THIS may be illustrated by the following

EXPERIMENT V.

Pl. 24. F. 3, 4, 5. THE Machine represented by the 3d, 4th, and 5th Figure of Pl. 24. was contriv'd by Dr. *Gravesande* (Professor of Mathematics and Astronomy at *Leyden*) to make this parabolick Motion of Projectiles evident to sight. A B b G F is a solid Piece of Wood 15 Inches high and 2 Inches thick, standing up upon its End F G, and set upon the Board or flat Piece F E, which has a little shallow Pit at E to be fill'd with soft Clay, so that when the Ball B falling from B along the curve Surface B b (which is made very smooth, or fa'cd with Brass or planish'd Tin) and describing in the Air the Parabolick Line b x δ E it may make a Mark to shew exactly the Place where it fell. The Piece G (*Fig. 4.*) is sometimes to be put on upon G E of *Fig. 3.* where the prick'd Lines represent it at G δ , having also a Pit to fill with Clay to receive the Mark of the falling Ball at δ . The Piece K (*Fig. 5.*) is to be put on over it upon Occasion, where it is represented by the prick'd Lines 1 3 x, to intercept the falling Ball, and shew by the Clay in its Pit where the Point x, of the Line b x falls. There is also an upright Board B D E to be taken off and on upon Occasion, on which are fasten'd the 3 Rings r, r, r, thro' which the Ball will move when it describes a Parabola from b to E, the Pieces of *Fig. 4.* and *Fig. 5.* being then remov'd.

The Height B C, or the Distance between the Line A B D and the Line C b c d e, is of six Inches; and the Height b G is of nine Inches, being divided into three unequal Parts, of which the uppermost contains 1 Inch, the next 2 Inches, and the lowest 5 Inches. Now if the Time, in which the Ball falls from b to G, be divided into three equal Parts (which we will here call *Instants*) the Ball will in the first of them fall the first Space mark'd 1, that is, one Inch; and in the next Instant the Space 3, or 2 Inches. and in the 3d Instant the space 5, or 5 Inches. N. B. *This happens on account of the accelerated Motion, the Cause of which we shall explain in considering the second Law of Motion.* When we let the Ball fall from B, and

B, and it rolls down along the curve Surface B *b*, it has acquir'd such a Velocity at *b* as is able to throw it forwards (double the Distance B C or Height from which it fell) by an uniform Motion, *viz.* from *b* to *e*, in the horizontal Line *b c d e*, if it was not forc'd out of it downwards by the Action of Gravity; but Gravity neither accelerates nor retards its Motion forwards, only makes it come to the Point E instead of the Point *e*, in the same perpendicular Line, and in the same Time. And if we consider Two other Points of the Parabola *b x s E*, *viz.* *x* and *s*, we shall find, that the Ball which in the first Instant of Time must have gone to *e* by the horizontal Motion or projectile Force, is brought down by Gravity to *x*; and instead of being carried to *d* the second Instant, it is brought to *s* in the Parabola, which Point is directly under *d*, and in the same horizontal Line as the End of the 4th Inch or 2d Space in the Line of Fall *b G*. To prove it experimentally; put on the two Pieces G and K of *Fig. 4* and 5, then letting the Ball fall from B it will run down the Curve B *b*, and then in the Air go from *b* to *x*, where it will make an Impression on the Clay: take off the Piece of *Fig. 5*. and let the Ball roll down again from B, and it will fall on the Point *s* as will also be seen by its Impression on the Clay at *s*, then take off the Piece of *Fig. 4*. and the Ball will go to E: and lastly, when you put on the Board B D E with its Rings *r, r, r*, the Ball falling from B will describe the Parabola *b E* passing thro' all the Rings, tho' there were as many more, provided the Parabola pass thro' their Centers. N. B. *The Reason why the Ball falling in the Curve B b acquires a Velocity to carry it forwards uniformly twice the length or Height B C, in a Time equal to the Time of the Fall, will be also shewn in explaining the next Law.*

If a Cannon Ball or any other Body be shot or projected obliquely upwards, it will likewise by its Rise and Fall describe a Parabola.

CONSTRUCTION. *Pl. 24. Fig. 6.*

DRAW a Line A *i* for the Amplitude of a Parabola, and upon its middle Point 3 raise the Perpendicular 3 E, upon which take the Height 3 *e* for the Axis of the Parabola. From *e* to E upon the Axis continued set off *e E = e 3*, and draw the Line A E, which will be a Tangent to the intended Parabola in the Point A. Divide the Line A E into any Number of equal Parts; as for Example, four,

Lect. V. four, which will be mark'd by the Points B, C, D, E: continue on the Line A E beyond F, so as to set on upon it four more equal Parts as E F, &c. not express'd in this Figure for Want of Room, but suppos'd to be over the Points G, H, I. Then from the Points B, C, D, F, &c. let fall the Perpendiculars B *k*, C *m*, D *o*, F *s*, G *u*, H *x*, and lastly I *i*, which will fall upon the Point *i* of the Line A *i* (or Amplitude of the Parabola.) Divide the Axis *e z* into 16 equal Parts, and mark the Points *q*, *z*, *2*, at the End of the first, fourth, and ninth Divisions; so that the four unequal Parts. *e q*, *q z*, *z 2*, and *2 3* may respectively contain one, three, five, and seven of the equal Parts. Thro' the Points *q*, *z*, and *2*, draw the Lines *d f*, *c g*, and *b b* parallel to the Line A *i*, which Lines will meet the Perpendiculars in the Points *b*, *c*, *d*, *f*, *g*, and *b*, and be Ordinates to the Axis *e z*. Upon the Line *e E* or Axis continued mark *e p* = *e q*, and *e Z* = *e z*, and likewise *e K* = *e 2*: from the Points K, Z, and *p* draw K *b*, Z *c*, *p d*, and on the other side *p f*, Z *g*, and K *b*. If a curve Line be drawn thro' the Points A, *b*, *c*, *d*, *e*, *f*, *g*, *b*, and *i*, it will be a Parabola, or the Line which a Projectile shot from the Point A in the Direction A B will describe; provided it has a Velocity given to it able to carry it the Length A B, in the same Time that by the Force of Gravity it would fall the Length B *b*. Produce, to, or beyond *t*, *w*, and *y*, the Lines *p f*, Z *g*, and K *b*, which (as well as *p d*, Z *c*, and K *b*) are Tangents to the Parabola in the Points *b*, *c*, *d*, *e*, *f*, *g*, *b*, *i*. Then thro' those Points draw the Lines *b a c*, *k d m*, *e o*, *f q*, *g s*, *b u*, and *i x*, which will be terminated by the Perpendiculars at the Points *a*, *k*, *m*, *o*, *q*, *s*, *u*, and *x*, and compleat the Parallelograms A B *b a*, *b l c k*, *c n d m*, *d p e o*, *e r f q*, *f t g s*, *g w b u*, *b y i x*. A *a*, B *b*, *b k*, *l c*, *c m*, *n d*, *d o*, *p e*, *e q*, *r f*, *f s*, *t g*, *g u*, *w b*, *b k*, *y i*, are all equal; which may be easily demonstrated from the Construction, and the Nature of the Parabola; but we rather chuse to demonstrate it from Gravity, in the following Explication.

DEMONSTRATION.

Pl. 24. F. 6. LET a Ball be projected in the Direction A B C D, &c. (Pl. 24. F. 6.) By the First Law it would go thro' the equal Spaces A B, B C, C D, D E, E F &c. in equal Times or Instants, and go on continually in the Line A F, &c. if Gravity did not act upon it to bring it down; but for the Reasons before alledged during the first Instant, Gravity will have brought it down to *b* (or, which is the same thing, it will move in the Line A *b* Diagonal of the Parallelogram

lelogram $A B b a$, two contiguous Sides of which $B A$ and $A a$ represent the two Forces that act upon the Body;) and at the End of the second Instant, instead of going forward according to the new Direction $A b$ in the Line $b l$, it will be brought down to c , moving in the Line $b c$ Diagonal of the Parallelogram $b l c k$, where $b l$ represents the *now* projectile Force, and $b k = B b = l c$ the Force of Gravity, which always pressing downwards with the same Force must be represented by the equal Perpendiculars $A a, b k, c m, d o, e q, f s$, &c; whilst the projectile Force in its different Directions is represented by the unequal and decreasing Lines $A B, b l, c n$, and $d p$, in the Ascent of the Body; but in its Descent, by the unequal increasing Lines $e r, f t, g w$, and $h y$, all Tangents to the Parabola. To go on, the projectile Body will successively describe the Diagonals $c d, d e, e f, f g, g h$, and $h i$, of the Parallelograms, two contiguous Sides of which represent the Quantities and Directions of the Forces $c n$ and $c m, d p$ and $d o, e r$ and $e q, f t$ and $f s, g w$ and $g u, h y$ and $h x$.

N. B. *We have call'd Diagonals the Curves intercepted between the Perpendiculars at the Points A, b, c, d, e, f, &c. or, which is the same thing, made the Diagonals curved; because those 'Diagonals, are really such, as we have hinted in the Explication of the Motion of a Body, projected horizontally.*


S C H O L I U M.

THE Projected Body moves with a Velocity uniformly diminish'd till it comes to the Vertex of the Parabola at e , then goes down the other half of it with a Motion uniformly accelerated. For since the Spaces $A B, B C, C D, D E, E F$, &c. have been taken equal in the Line of the first Direction of the Projectile; by the first Law, the Projectile must go thro' them in equal Times, if none but one Force (*viz.* the projectile Force) acted upon the Projectile. But as Gravity acts upon the Body at the same Time, tho' it cannot hinder it from successively reaching the Perpendiculars (which are equidistant and parallel to each other) in the same Space of Time that it would have done if there had been no Gravity, it will bring the Body, at the End of every Instant, to other Points of the Perpendiculars, which are still nearer and nearer to each other, till the Body comes to e : as for Example, the Body at the End of the first Instant instead of being at B is brought down by Gravity, to b falling one Space equal to $e q$; at the End of the second Instant, instead of

Lect. V. of being at C, it is brought down to c , having fallen from the first Direction or Line of the projectile Force thro' a Space equal to $e z$, or four times $e q = B b$, the Space fallen thro' the first Instant; at the End of the third Instant, instead of being at D, it will be brought down to d thro' a Space equal to $e 2$, containing 9 times $e q$, or $B b$, the Space fallen thro' the first Instant: at the End of the fourth Instant, instead of being at E, it will be brought down to e , thro' a Space equal to $e 3$, containing sixteen of the Spaces of the first Fall $B b$ or $e q$. Now since the Spaces which the Body goes thro' in its Ascent, in equal Times (that is, the Lines $A b$, $b c$, $c d$, and $d e$) are not only less than the Spaces $A B$, $B C$, $C D$, $D E$, but also less than each other, because their Change of Direction makes them continually cut the parallel Perpendiculars at less oblique Angles, it is evident that *the projected Body continually diminishes its Velocity till it comes to its utmost Height at e*, the Vertex of the Parabola: and this will also appear, if we consider the Motion of the Projectile here, as we did that of the horizontal Projectile; viz, by examining how several Parts of the Line describ'd by the Projectile are like so many Diagonals describ'd by the Action of two Forces, one of which changes its Quantity and Direction by extremely small Intervals, tho' here we take great Intervals to make the Thing more plain.

WHEN therefore the Body projected sets out at A in the Direction A B, A B represents the projectile Force, and A a the Force of Gravity; as the Lines B b and $b a$ compleat the Parallelogram by Construction, the Body will go in the Diagonal A b †. The Body being at b will, by the first Law, endeavour to continue to move in the Line $b l$, being the Diagonal continued, that is in the new Direction which it has now acquir'd, and with the same Velocity which it now has (which is less than what it had at A, because the Diagonal A b is shorter than the side A B); therefore $b l$ now represents the projectile Force, and $b k$ the Force or Gravity, which we take equal to A a ; because we have no Regard to the Body's being in Motion, but consider it as setting out from the Point b , by the Action of the Two Forces $b l$ and $b k$: the Body thus acted upon will move in the Diagonal $b c$ of the compleated Parallelogram $b l c k$, with a Velocity as much less than it would have had in the Line $b l$, as the Diagonal $b c$ is shorter than $b l$. The Body at C will, by the first Law, endeavour to go on in the Line $c n$ with the Velocity which it now has, but the Action of Gravity represented

† Pl. 24. F.6.

represented by cm ($= bk = Aa$) bringing it down out of that Lect. V.
Line cn will make it go in the Diagonal cd with a Velocity lessen'd 
in Proportion as cd is shorter than cn . Lastly the Body at d acted
upon by the Forces dp and do will go in the Diagonal de with
chang'd Direction and diminish'd Velocity, as we have shewn be-
fore.

As the Body comes down from its utmost Height at e , it continually increases its Velocity. For, first, if we consider it when at e ; the projectile Force is er , Gravity eq , and ef (the Diagonal of the Parallelogram $erfq$) the Line describ'd by the Body. Now as the Line er representing the Direction and Velocity of the Body at e is shorter than the Tangent ft representing its Direction and Velocity when at f (because er is perpendicular and ft oblique to equidistant Parallels) the Diagonal fg must be longer than the Diagonal ef ; therefore the Body's Velocity will be so much the greater. Thus will the Diagonal gb of the Parallelogram $gwbu$, the Space gone thro' by the Body in the next Instant be greater than fg , and consequently its Velocity encreas'd in that Proportion. And lastly, bi the last Space being the Diagonal of the Parallelogram $byix$ will be still greater than the last, and consequently the Velocity also greater.

C O R O L L A R Y I.

HENCE may be known the different Velocity of a Projectile in any Point of the Parabola which it describes, whether in its Ascent or Descent: and *it will always be to the Velocity in any other Point, directly as the Lengths of the Parts of the Tangents to the Parabola at those Points which are intercepted between the same or equidistant Parallels.* As for Example, the Velocity at A : is to the Velocity at b :: as BC : to bl ; for since (by Construction) BC is equal to AB , and AB represents the Velocity of the Body at A , as bl represents the Velocity of the Body at b ; the Velocities at the said Points A and b will be respectively as BC to bl . Thus if we compare the Velocities at A and at d , they will be to one another as DE to dp , for the same Reason. So likewise in the Descent the same will hold good; as for Example, the Velocity at e : is to the Velocity at f :: as ri to: to ft ; or er : ft , 107: t 5, or 78: 56. So the Velocity at f : is to the Velocity at b :: as 56: to by ; or ft , or t 5: to by .

Q q

C O R O L-



C O R O L L A R Y II.

HENCE follows also, that at equal Heights above the horizontal Line or Amplitude A *i*, the Body will have the same Velocity; because it will then be in correspondent Points of the ascending and descending Parabola, where the Tangents, having equal Obliquities to the Parallels to the Axis, will have equal Parts cut off by those Parallels wherever they are equidistant.

C O R O L L A R Y III.

It also follows; that however unequal the Spaces be that a Projectile describes in equal Times, the horizontal Distances (that is, its Advances forwards) will all be equal in equal Times; but we have already prov'd this another Way. N. B. *I don't mean that the same Quantity of projectile Force will make a Body go forward equally fast; for that will vary according to the Angle that its Direction makes with the Horizon; but that if the Amplitude of the Parabola (or the whole horizontal Distance which the Body goes thro', whilst it describes any one Parabola) be divided into equal Parts by Perpendiculars; the Body will go from one Perpendicular to another in the same Time.*


4. In considering the Motion of a Projectile, we have made the Line A *i* a straight Line, which would be strictly so if the Earth was flat; and indeed so much of it as a projected Body goes over must be taken for such: but if the Force of the Powder, or whatever throws the Body forward, was much greater; or the Force of Gravity (that is, the Attraction of the Earth) much less; then we must consider the Line A *i* as a Curve or part of the Earth's Circumference.

* Pl. 24. F. 7. As for Example; Let A B C D E * represent the Surface of the Earth, and A F an high Mountain, such as *the Peek of Tennessee*. Now if a Cannon was fir'd in the Direction F L, the Ball would go forwards in a Curve, for the Reasons before alledg'd, perhaps as far as B, where it would come to the Ground. If the Force of the Powder was proportionably greater, it would go as far as C before it came to the Ground. Suppose the Force of the Powder yet greater, and the Ball would go to D: suppose it still greater, and the

the Ball would go to E, not falling to the Ground till it had gone over Lect. V.
two thirds of the Circumference of the Earth. Lastly, one might
suppose the Force of the Powder great enough for the Ball not to
come to the Ground at all, but to come to the Point E from whence
it set out at first, and (the Cannon being remov'd out of the way)
to go continually round the Earth, at the Distance of about 3 Miles
from the Surface of the Earth (that is, at the Height of the *Peek*
of Tennerif) Gravity only keeping it from going off from the Earth
in a Tangent Line. Were the Force of the Powder greater than in
the last Supposition, the Ball would go farther and farther from
the Earth in a spiral Line.

If the Point F was remov'd sixty times farther from the Center
of the Earth, or to the Height of 240,000 Miles, that is, to the Di-
stance of the Moon; then the Force of Gravity (or the attracting
Force, call'd in that case *the accelerating Force*, of the Earth)
would be 3600 times weaker than at the Surface of the Earth;
because, as we recede from the Center of the Earth, Gravity de-
creases as the Squares of the Distances increase; and the Moon being
60 times farther from the Center of the Earth, than the Earth's Sur-
face, the Square of that Distance 60 is 3600. In such a Case there
would be no Occasion for a greater projectile Force than that of our
common Powder to make a Cannon Ball circulate round the Earth, at
the Distance of 24000 Miles. The Moon it self may be confi-
der'd as such a Projectile; for having once receiv'd an Impulse in a
Line parallel to a Tangent of the Earth, while it is endeavouring
(by the First Law) to keep its first (rectineal) Direction, Gravity
is continually impelling it towards the Center of the Earth, and by
that Impulse turns it out of the right-lin'd Direction of the projectile
Force, and carries it continually round the Earth. Neither can
Gravity bring the Moon to the Earth, because the projectile Force
still subsisting always endeavours to throw off the Moon along the
Tangent of its Orbit; and Gravity only alters the Direction con-
tinually, whereby the Moon endeavours every Moment to fly off
along a new Tangent. Thus in every Point of the Orbit, these
two Forces balance each other.

5. That Endeavour of the Moon, or any other Body that
moves in a Circle or any other curv'd Orbit, to fly farther
from the Center of its Motion, is call'd a *Centrifugal Force*;
and the Force acting against it to keep the moving Body in its
Orbit (whether it be the Force of Gravity or Attraction,

Lect. V. or a String, driving, impelling, or drawing the Body towards the Center of its Orbit) is call'd a *Centripetal Force*. Thus when a Stone is whirl'd in a Sling, the Stone, which endeavours to (and when one of the Strings of the Sling is loos'd, actually does) fly off in a Tangent to the Curve which it describ'd before, is hindred by the Strings to fly off of the Sling from so doing; that Endeavour is the centrifugal Force; and the Force of the Strings holding the Stone, is the centripetal Force. By the Force which endeavours to, but cannot really, carry the Stone in the Tangent, the String is stretch'd in a Direction from the Center of the Revolution to the Circumference. Let us, for Example, suppose, that the Body F  is whirl'd round in the Circle F I N: whilst the Body, not held by the String, would have mov'd the Length FH in the Direction FG by the first Law, being retain'd by the String it moves in the Arc FI, being brought down from H to I; the Line HI therefore may represent the Quantity of the Centripetal Force directed towards the Point M; and the same Line IM directed from the Point M towards H may also represent the centrifugal Force, both those Forces being equal, otherwise the revolving Body would not be kept in Orbit. For tho' in Orbits that are not circular, these Forces may encrease and decrease (as shall be hereafter shewn) yet they both equally encrease and decrease, always balancing one another. N. B. *The swifter a Body moves in its Orbit, the more is the String stretched, that is, the greater is the Centrifugal and Centripetal Force. If, for Example, FG represents the Velocity of the Body instead of FH, the Body will describe the Arc FN instead of FI; in which case the Point N (where the Body is) being farther remov'd from the Tangent (at G, where the Body would go to by the projectile Force alone) the Centripetal and Centrifugal Force must be represented by the Line GN, which is greater than HI.*

ALL these will be further prov'd and illustrated by the following Experiments,

EXPERIMENT VI. *Pl. 24. Fig. 8.*

6. ABCD is a round Table which may be swiftly turn'd upon a Pivot, as at F (the same that is represented by the 1th Fig. of *Pl. 23.*) There is a little Brass Pipe screw'd in at the Center C, into the Top of which the String of the leaden Bullet B is thrust so as as to go out at an Hole in the Side of the said short Brass Pipe; thence

thence it is carried under the Table thro' the Hole *b*, and so fasten'd to a Pin in the Side of the Table at A. When the Bullet is laid at B, if the Table be turn'd swiftly round, it leaves the Bullet behind at first, which thereby appears to have a Motion contrary to that of the Table, till by the roughness of the Surface of the Table, it goes round at last along with the Table on the same Part of the Table; then if the Table be stopp'd on the sudden, the Bullet goes on several turns, till having communicated all its Motion to the rough Surface of the Table, it comes to Rest at last. This illustrates the first Law of Nature; for as the Part of the Table under the Bullet leaves it behind for a while, because it endeavours to continue in its State of Rest, it would for ever leave it where it was at first, if the Table was perfectly smooth: and when the Bullet is once in Motion it would for ever go round on the Table, if (besides the smoothness of the Table) the String that holds the Ball had no Friction at the Center C. It is also to be observ'd, that the String, which is slack at C B, is always stretched as at C *b* by the Motion of the Bullet; and this shews the Centrifugal Force.

If we screw a forked Prop toward the Edge of the Table as at D, and put the String of the Bullet into its slit so as to let the Bullet hang down as at 1, the Force of the Bullet's Gravity may be so overcome by the Centrifugal Force, which the whirling of the Table produces, that the Bullet shall rise to 2, the String 3 *d* becoming horizontal: as the Table turns slower and slower, the Ball comes down to 2 and so to 1 at last, Gravity becoming sensible as the Centrifugal Force diminishes.

EXPERIMENT VII. *Pl. 24. Fig. 9.*

If a String be tied round the Brim of a Pot full of Water, and the Pot be whirl'd round swiftly about the Hand or Center K in a Circle or Curve of which A C B is an Arc, the Water acquiring a Centrifugal Force greater than that of Gravity, will not be spill'd when the Mouth of the Pot is downwards. If instead of the Pot, the Glass W C † (*Fig. 10.*) containing Liquors of different specifick Gravities be whirl'd round the Center K, after they have been confounded together by shaking) they will all recover their Places and Transparency, even sooner than if the Glass containing them had been hung up and at rest. The Reason is, that as the different Substances in the Glass have the same Velocity given them by the Centrifugal Force, their *Momentum* will be as their specifick

Left. V.
~~~~~

Pl. 24. F. 9.

† Pl. 24. F. 10.

Lect. V. sick Gravities, that is, their *Momentum* will be made up of the different Quantities of Matter, which they contain under equal   
 † L. 2. N<sup>o</sup>. 3. Bulks multiplied by the common Velocity † which the Centrifugal Force gives them in the Line K C from the Center of the Motion towards the Circumference. Therefore the Glafs Beads among the Liquors weighing more than the Drops of any of the Liquors, will have the greatest *Momentum*, and consequently go to the Part G most remote from the Center of Motion K. Then the Drops of Oil of Tartar (which is the heaviest of the Liquors contained in the Glafs) having for the same Reason more *Momentum* than the Drops of the other Liquors (tho' less than the Glafs Beads) will take up the Space T next to the Beads, and also fill their Interstices. The next Liquor which is Oil of Peter, will fill up the space P. And lastly, the Spirit of Wine, whose Drops are the lightest, will (notwithstanding its own Centrifugal Force) be brought nearer to the Center of Motion and occupy the space W; because the Beads and all the other Liquors having more *Momentum*, drive it from the End C, to which it has a Tendency all the while. N. B. *The Tube is hermetically seal'd at both Ends.*


The Glafs Beads, and different Liquors, settle in their proper Places when the Tube is hung up; because, as all Bodies tend downwards with the same Velocity,\* the *Momentum* of Particles of equal Bulk must be as their respective Quantities of Matter in their Descent: And that the Liquors will not be so soon settled in this Case as when the Tube is whirl'd round, is because we can give as great a Velocity as we please in the Direction K C, by a Centrifugal Force; whereas that which is owing to Gravity is always the same.

### C O R O L L A R Y.

HENCE it follows, that a Bottle of any Liquor (which after having been muddy is by length of Time become fine, and is again made foul by shaking) may sooner be brought to be fine by a Centrifugal Force, than by being set upright at Rest.

### EXPERIMENT VIII. Pl. 24. Fig. 11.

\* Pl. 24. F. 11. JOYN by a String to the two Balls T and M,\* whose Weights are to one another as 4 to 2 (here we use a two-Ounce and a four-

four-Ounce-Ball) and pass the String thro' the opposite Side-Holes of Lect. V. the little Pipe C; let the Length of the String, measuring from  Center to Center of the Balls, be 18 Inches, and the Distances of the Balls from the Center of the Table be reciprocally as their Masses; that is, the two-Ounce Ball M must be at the Distance of 12 Inches from C, and the four-Ounce Ball T at 6 Inches from C. Let the two little Squares, or rectangular Pieces S s and V v be fix'd on the Table at the Distance of about 1 Inch or 2 behind the Balls to stop them from flying off of the Table, and the long Sides of those Pieces so fix'd alternately, that when the Table is made to turn in the Direction mark'd by the Dart, the Balls may not be left behind, but immediately put into Motion. Now let the Table be whirl'd round with any Velocity, and the Balls will remain at the Points T and M, and describe round their common Center of Gravity unequal Circles \* in a reciprocal Proportion of the Masses, \* L. 2. N<sup>o</sup>. the *Momenta* given the Bodies by the centrifugal Force being equal, <sup>32.</sup> and (because of their contrary Directions) destroying one another. But if either of the Balls be remov'd farther from C than in the reciprocal Proportion above-mention'd, that Ball will gradually recede from the Center of Motion and draw the other along with it, till it be stopp'd by the End of the Piece V v or S s. So the Earth and Moon turn round one another and round their common Center of Gravity as has been already observ'd †.

† L. 2. N<sup>o</sup>.  
32.

EXPERIMENT IX. Pl. 24. Fig. 12.

LET the Table be put into an uniform whirling Motion, and that Motion continu'd (which may be done by a Wheel and Pulley hereafter to be describ'd) and at the same Time let a Piece of Chalk be drawn pressing upon the Table in the right-Line C P, and there will by those two Motions be described the spiral Line C D E F P: then if the Table be mov'd again in the same manner and the same Way (as mark'd by the Dart) and at the same time the Chalk be press'd on the Table in the same Line P C, but with a contrary Direction, *viz.* from P to C, there will be describ'd another Spiral like the former, but directed the contrary Way, as is shewn by the prick'd Line.

## C O R O L L A R Y.


HENCE it follows, that if a Body that has a Centrifugal Force whereby it recedes from, or a Centripetal Force, whereby it accedes to, the Center of its Motion, at the same time be carried round by a Force that gives it a circular Motion; it will fly from the Center in a spiral Line in the first Case, and run to the Center in a spiral Line in the second Case.

EXPERIMENT X. *Pl. 24. Fig. 13.*

\**Pl. 24. F. 13.* ON a Piece of Board A E K\*, which has a Piece under it across to raise up its broadest End A E to an Angle of 15 or 20 Degrees above the horizontal Position, are fasten'd 3 Tubes A K, C K and E K, shut up at both Ends. In the first there is a small Cylinder of Cork, which can easily slide up and down the Tube: in the Tube C K there is a little Cylinder of Lead moveable in the same Manner: and in the Tube E K, there is an Inch or two of Quicksilver shut up. This Board has a Screw under it, which going thro' one of the Holes in the Table (such as are mark'd A and B †*Pl. 23. F. 13.* in *Plate 22. Fig. 13*)† is fasten'd by a Nut so as to joyn the Board of Tubes firmly to the Table. Then when the Cork, Lead, and Quicksilver are in that Part of the Tube next to the Center of Motion K; let the Table be whirl'd round, and those Bodies will after a few Turns be carried to the Ends of the Tubes which are farthest from the Center, tho' 3 or 4 Inches higher than the Ends at K. Put on the Tubes B K and D K, the first of which being fill'd with Water has a Cork Cylinder moveable in it; and the other has in it an Inch or two of Oil, the other Part of it being full of Water. At first (when the Table is at Rest) the Cork and the Oil will be at B and D the higher Ends of the Tubes and farthest from the Center; but when the Table is whirl'd round, the Cork and the Oil will go towards the Center to K, because the greater Centrifugal Force of the Water (being greater than either that of the Cork, or that of the Oil) must give the Cork and the Oil a Centripetal Direction, as has been explain'd in the Experiment of the different Liquors in the Glass of *Fig. 10* \*.

*Pl. 24. F. 10.*

7. *Monf. Des Cartes* endeavour'd to explain the Motion of the Planets round the Sun, by a *Vortex* or Whirlpool of Celestial Matter

ter ; supposing that the Sun by by turning about his Axis, gave Lect. V.  
circular Motion to all the Celestial Matter about it as far as   
*Saturn* and beyond ; and that this Whirlpool of Matter (in *French*,  
*Tourbillon*) being without Vacuity (for he asserted *Plenum*) carried the Planets along with it, and so was the Cause of their Motion round the Sun. People that did not give themselves the Trouble to examine Things thoroughly, believ'd the *Vortices* to be the Cause of the Motion of the Planets ; because, where there are Whirlpools in Rivers and Brooks, we see that Straws, little Sticks, Saw-duft and other light Bodies are carried round as they float ; and therefore it was thought very probable, that the Planets might in the same manner be carried round in the Heavens. But, if we apply Experiments and Observations to the Doctrine of the *Vortices*, we shall find it insufficient for explaining the Motions of the Planets.

IN the first place, we'll suppose the Planets to be denser than the Matter of the *Vortex* ; and then the same Thing must happen to them, as happen'd to the Cork, or the Lead, or the Quicksilver contain'd in the Glass Tubes, and whirl'd round on the Table. For the Tubes full of Air being carried round, contain'd part of a *Vortex* of Air, where the Bodies carried in it were of different specific Gravities, but all specifically heavier than the Parts of the *Vortex* ; and those Bodies mov'd round and at the same time agitated by a centrifugal Force did continually recede from the Center of the *Vortex*, going off from it in a spiral Line. Planets therefore specifically heavier than the Parts of the *Vortex* would continually recede from the Sun moving off in a spiral Line. Since this won't do, let us suppose the Planets all rarer than the Matter of the *Vortex* ; and the Consequence will be---- That as the Parts of the *Vortex* are denser than the Planets, their centrifugal Force must give the Planets a centripetal Direction, and so make them go continually towards the Sun in a spiral Line, till they fall into it ; after the Manner of the Cork and the Oil in their respective Tubes of Water, where it appear'd by the Experiment that the centrifugal Force of the Water gave the Cork and Oil a contrary Tendency, whereby they continually approach'd towards the Center in a spiral Line.

THERE remains only now for the Support of the *Cartesian Hypothesis*, to suppose that the Planets are of the same Density as the

Lect. V. Parts of the *Vortex* (and indeed, if there was a *Plenum* as *Monf. Des Cartes* supposes, all Bodies must be equally full, and there wou'd be no such thing as different specifick Gravities ; for all Bodies  
 Pl. 24. F. 2. of the same Bulk, having the same Quantity of Matter, wou'd weigh equally) and in that Supposition it will follow that the Planets making up Part of the *Vortex* itself will move along with the Parts of the *Vortex* next to them ; and tho' at first those Parts of the  
 \* Ann. 2. *Vortex* which are nearest to the Sun will move fastest ; \* yet at last, the whole *Vortex* will move round like a solid Body or Sphere, the Parts most remote from the Center making their Revolutions in the same Time as those which are nearest ; which must be the Consequence of a *Plenum*. So that all the Planets wou'd at last have their periodical Times equal, which is contrary to Observation ; for the periodical Times of the Planets Revolutions are all different. *Mercury*, the nearest to the Sun, performs his Revolution almost 120 times sooner than *Saturn*, the most distant. In a word, this is the Proportion of their Distances and Revolutions, viz. *The Squares of the Periodical Times of the Primary Planets round the Sun, and of the Satellites round Jupiter and Saturn, are as the Cubes of their Distances from their respective central Bodies.* †

8. SINCE then the *Cartesian Hypothesis* is insufficient for explaining the Cause of the Motion of the heavenly Bodies ; we must shew what is the real Cause, and that not from Conjectures, but from Observations and Experiments. That Gravity or an Attraction towards the Sun keeps the Planet (or even the Comet) in its Orbit about the Sun (and likewise the Satellites about their Primaries, our Moon also being the Satellite of the Earth) whilst the projectile Force is continually endeavouring to throw it off along the Tangent, will evidently appear from what we have already said, and from Consequences of the Second Law of Nature, which we are going to explain, having first shewn several Laws relating to central Forces by the following Experiments.


But we must first describe the Machine for the making the Experiments.

Plate 25. Fig. 1.

Pl. 25. F. 1. THE Machine for central Forces consists of a strong wooden Frame CABDHGKEF triangular at Top and Bottom. On the horizontal

zontal Piece *G a* at Top is a Wheel *G*, which (by means of the *Lect. V.*  
String *G K H G*) when turn'd round, gives circular Motion to the *Pl. 25. F. 1.*  
Pullics and Spindles *K L* and *H I*, so as to move them either with  
equal Velocities, or with Velocities that are as 2 to 1, as 3 to 1,  
or as 3 to 2; because in the Pulley *K* there are Two Grooves, One  
of 3, and One of 2 Inches Diameter; and in the Pulley *H* there are  
also Two, One of 6, and One of 3 Inches Diameter. There are  
Two Pieces *M N*, *m n* (which we may call Planet-bearing Pieces)  
of about 30 Inches long, to be screw'd upon the Pullics *K, H*, so as  
to turn round with them. These Pieces have each of them an open  
square Tower *S, s*, with a little Pulley at Top and Bottom to con-  
duct a String from the Weights *S* and *s*, to the Brass Balls *P* and *p*,  
(which we must here call Planets) so that when *P* and *p* go to-  
wards *N* or *n*, the Weights are drawn up from their Bases, which  
is about an Inch above the Bottom of the Towers, and rise within  
the Towers till the Weight-carrying Piece strikes the Top of the  
Tower; each Ball having Two little Wings with Holes in them to  
slide easily along Iron Wires that go from one End to the other of  
the Planet-carrying Piece, passing thro' the Two perpendicular Brass  
Planes *M* and *N*, also thro' the Towers at about the Distance of  $\frac{1}{4}$  of  
an Inch from the Surface of the Planet-carrying Piece. N. B. *Only*  
*one of the Wings and one of the Wires is here drawn upon each*  
*Piece to avoid Confusion.* There are also Brass Collars at *H* and  
*K*, in which the Necks of the Spindles (which are of Steel) turn;  
and Iron Screws headed with Brass at *L* and *I*, with little Holes to  
receive the Bottoms of the Spindles.

THE Second Figure \* represents something more than half of one \* *Pl. 25. F. 2.*  
of the Planet-bearing Pieces divided into Inches both Ways from the  
Center. *B b* is the perpendicular Brass Plane at one End, thro'  
which the horizontal Wires *W w*, *W w* pass, to carry the Planet  
*P* by its perforated Wings *LL*, whilst the String that goes thro'  
the middle of the Planet is made fast, by thrusting in the little Pin  
*p* to give the Planet any certain Distance from *T* the Center of its  
Motion before it is mov'd round by turning the Wheel *G* (in *Fig.*  
*1.*) *S s* represents the Section of the Brass Tower fasten'd to the  
Wood by a cross Pin, whose Head is seen at *s*. *T* is the Base or  
Plate which is to support the Weight-carrying Piece, which is re-  
presented at (*Fig. 4.*) † and consists of a circular Plate and hollow † *Pl. 25. F. 4.*  
Stem of Two Ounces Weight, and on which may be slipp'd feve-  
ral

Lect. V.  ral leaden Weights like X *Fig. 5.* \* At T also (*Fig. 2.*) may be seen the little Pulley under which the String goes.

\* Pl. 25. F. 5.

† Pl. 25. F. 3. THE 2d Figure † is a vertical Section of one of the Square Towers S s, with the Weight X and Weight-carrying Piece X x in it, and Part of the Planet-bearing Piece, Pulley and Spindle under it, mark'd M N. One little Pulley is fasten'd to the Wood at T under the Plate on which the Weights stand; another is sustain'd by an Iron Arm V S over an Hole S in the Top of the Tower. So that the String coming from the Planet P goes first under the Pulley T, then thro' the hollow Stem of the Weight-carrying Piece, and so thro' the Hole in the Top of the Tower, then over the Pulley S, so down again to the Top of the Stem of X where it is fasten'd. By observing this Figure, one may easily see that if the Planet P be mov'd ever so little in the Direction P Q, the Weight bearing Piece X will be rais'd up towards S.

\* Pl. 23. F.

13. and Pl.

24. F. 8, 11,

12, 13.

THE whirling Table which we have already mention'd \* for making several Experiments is best turn'd round by screwing on to the Top of either of the Spindles L K or I H, instead of the Planet-bearing Piece M N or m n.

\* Pl. 25. F. 6.

THE sixth Figure \* represents the Section of the Wheel and Part of the horizontal Piece, the upper Part of the Frame which carries the Wheel, and the upper End of the Piece that supports it mark'd L L L, the Wheel's Axis and square-sliding Collar g, which is moveable on the square horizontal Iron H I fasten'd to the Wood by a Nut and Pin at I, and two Wood-Screws H b. N. B. *There is a Screw in the sliding Piece g to fix the Center of the Wheel, when it is brought forwards or backwards.*

*THO' it be of no Consequence of what Bigness the Machine above describ'd is made, provided its Parts are proportional; yet for the sake of those who would have such a Machine made, I give here the Measures of the principal Parts of mine in English Inches.*

Pl. 25. Fig. 1.

Pl. 25. F. 1. THE Thickness of the Wood every-where about 1 Inch, except the Feet at A and D, where it is 2 Inches thick.

M N

M N =  $mn$  = 30 Inches.

K H 33 $\frac{1}{2}$  Inches.

K L = I H = 8 $\frac{1}{2}$  Inches.

A D = 34 $\frac{1}{2}$  Inches.

A C = C D = 27 $\frac{1}{2}$  Inches.

B C = 24 $\frac{1}{2}$  Inches.

Diameter of the Groove of the Wheel G = 14 Inches.

Breadth of the Planet-bearing Piece M N or  $mn$  = 2 $\frac{1}{2}$  Inches.

Grooves of the Pulley K, the one 2 and the other 3 Inches.

Grooves of H, the one 6 and the other 3 Inches.

The Height A K = H D = 13 Inches.

Height of the Towers S or s above the Board M N,  $mn$  5 $\frac{1}{2}$  Inches.

Breadth of the Towers = 2 $\frac{1}{2}$  Inches.

There are 4 Brafs Planets made use of, Two of which weigh each 2 Ounces Troy, and the Two other 4 Ounces Troy each.

The Weight-carrying Plate and Stem weighs 2 Ounces, and each leaden Weight (as represented by *Fig. 5.*) weighs 2 Ounces also.

*What is to be consider'd in the Use of the Machine.*

THE Weights in each Tower are to represent the Sun, whose Attraction is shewn by the Force with which the Weight resists the Action of the Ball P or  $p$  (representing a Planet) that endeavours to raise it by the String P T S  $\propto$  † (*Fig. 3.*) when it receives a † Pl. 25. F. 3. centrifugal Force by turning the Wheel G. So that by putting equal or unequal Weights in the Towers; by using equal or unequal Planets as P, or  $p$ ; and by having their Distances equal or unequal in different proportions; and the Periodical Times equal or unequal (as the Wheel-String goes round equal or unequal Pul- lies) we may by Experiments shew those Laws of central Forces, which Sir *Isaac Newton* has mathematically demonstrated in his *Principia*.

IN considering the central Forces of Bodies (for Example, of the Primary Planets in respect of the Sun, and of the Moons in respect to their Primary Planets) which move round other Bodies that have an Influence upon them; we are to observe Three Things. 1<sup>st</sup>, The Periodical Times, or Times in which the Bodies perform their Revolution. 2<sup>dly</sup>, The Quantity of Matter in  
the

Lect. V. the revolving Bodies. 3<sup>dly</sup>, The Distance of the Bodies from the Center of the Revolution.

### EXPERIMENT XI.

FIRST, make the Periodical Times equal, by putting the Wheel String into the 2 Inch Groove of each Pulley, crossing it before each Pulley to give it the more Force to move the Pullies, but so that the Pullies may both turn the same Way that the Planet-bearing Pieces may not unscrew. Then put only the Weight-carrying Piece in each Tower: and lastly, fasten to their Strings a 2 Ounce <sup>† Pl. 25. F. 1.</sup> Brass Ball, as P and p, † at the Distance of 12 Inches from the Center on each Planet-bearing Piece. So you will have *the Periodical Times, the Quantities of Matter, and Distances from the Center equal.* Give circular Motion to the Wheel G, and the Planets by their centrifugal Force will raise the Weights S and s at the very same Instant of Time; which shews, that *in this Case the centrifugal Forces are equal.* N. B. *If upon each Weight-carrying Piece you put on one or two, or more equal Weights (such as are express'd by Fig. 5.) the Planets will always raise them at the same Instant, provided the Wheel be turn'd proportionally faster as there is more Weight.*

### EXPERIMENT XII.

SECONDLY, Instead of p put on a 4 Ounce Ball, and double the Weight in the Tower s q; then turn the Wheel, and both Weights will rise at once. This shews, that *when the Quantities of Matter in Planets are unequal, but the Distances and Periodical Times still remain equal, the centrifugal Force is proportional to the Quantity of Matter.*

### EXPERIMENT XIII.

THIRDLY, Take off the 4 Ounce Ball, and make use of p again, but put it only at 6 Inches from the Center. Take the additional Weight from s and add it to S; that is, let the Weight S, which has P at 12 Inches Distance, be = 4 Ounces; and the Weight s, which has p but at 6 Inches Distance, be = 2 Ounces: then upon turning the Wheel, they will both rise at the same Time. Hence it is plain, that *if the Periodical Times, and the Quantities of Matter*

*Matter continue the same, but the Distances are different, the* Lect. V.  
*centrifugal Forces will be as the Distances.*

Pl. 25. F. 1.

#### EXPERIMENT XIV.

FOURTHLY, at the Distance of 6 Inches, where  $p$  was last, change  $p$  for a 4 Ounce Ball, and put equal Weights in the Two Towers; then when you turn the Wheel, both Weights will rise at once; which shews, that *when the Periodical Times are equal, and the Distances from the Center reciprocally as the Quantities of Matter in the Planets, the Centrifugal Forces are equal.*

#### EXPERIMENT XV.


LASTLY, Change the String on the Pulley H, putting it into the Groove of 6 Inches Diameter, so that the Periodical Time of the Planet  $p$  may be double that of the Planet P, which last will then move twice as fast, if its Distance be the same from the Center, which it must be in this Experiment. Put 8 Ounces in the Tower S Q, and only 2 in the Tower  $s q$ , the equal Planets P and  $p$  being then each at 12 Inches Distance from the Center. Turn the Wheel, and both Weights will rise at once. This shews, *That Planets, that have an equal Quantity of Matter and the same Distance from the Center, but different Periodical Times, have their centrifugal Forces reciprocally as the Square of their Periodical Times; that is, directly as the Square of their Velocities.*

#### COROLLARY.

HENCE follows, that if the same Planet changes in Velocity in the same Orbit, its centrifugal Force will encrease or decrease according to the Square of the Velocity which the Planet has in that Part of its Orbit.

#### SCHOLIUM.

WHEN we compare the last Experiment with the 12th Experiment, and find that the Planet ( $p$  going round in a Circle of 12 Inches Radius at the same Time that P went round in a Circle of 6 Inches Radius,) rais'd twice the Weight because it had twice the Velocity; it will appear strange, that in the last Experiment, where  $p$  (going


 Lect. V.  $p$  (going twice thro' a Circle of 12 Inches Radius, whilst  $P$  goes once thro' such a Circle) has but double the Velocity of  $P$ , it should now raise four times more Weight. But this Proportion (which is of very great use in explaining the Motions of the heavenly Bodies) will be very clearly deduc'd from a Consideration of the First Law and what we have said upon it.

† Pl. 24. F. 7. ROUND the Body supposed at  $M$  † (*Plate 24. Fig. 7.*) or the Center  $M$ , let a Planet revolve in the Circle  $FIN$ ; it is plain, that when it describes the Arc  $FI$ , its centrifugal Force is represented by the Line  $HI$ : then if the same Planet be supposed to describe the Circle  $fin$  of double the Radius, in the same time, it is evident that it will have a double Velocity, and consequently will describe the Arc  $fn$  in this Circle in the same Time that it described  $FI$  in the little Circle; so that now  $gn$ , which is double of  $HI$  or  $bi$  (because the Triangles  $fgM$  and  $FHM$  as well as the Arcs  $fn$  and  $FI$  are similar) will represent the centrifugal Force to be double of what it was in the little Circle. But if, instead of going in the great Circle, the Planet doubles its Velocity by going twice round the Circle  $FIN$  in the same Time that it went once round before, it will describe the Arc  $FN$ , of twice the Number of Degrees, instead of  $FI$ ; and as the Arc  $FN$  (tho' of the same Length) is twice more curve than  $fn$ , the Planet at  $N$  will recede from the Tangent twice as much as if it was at  $n$ , consequently four times as much as when at  $I$  with half the Velocity. Therefore the centrifugal Force at  $N$ : will be to the centrifugal Force at  $I$ :: as  $GN$ : to  $HI$ ; or as 4: to 1.

N. B. *This holds good only in small Arcs, as  $FI$  and  $FN$  are suppos'd; but then it is only to be consider'd in such.*

THERE are many more Experiments relating to central Forces to be made with this Machine; but these are sufficient for our present Purpose. However, for the sake of the Curious we shall mention a few more in the Notes. †

† Ann. 3.

*The Second LAW of Motion.*

10. *The Change of Motion is always proportionable to the moving Force impress'd; and is made in the right Line in which that Force is impress'd.*

If any Force generates a Motion, a double Force will generate double the Motion, a triple Force triple the Motion, &c. whether that Force be impress'd all together and at once, or gradually and successively. And this Motion (being always directed the same way with the generating Force) if the Body mov'd before, is added to or subtracted from the former Motion, according as they directly conspire with, or are directly contrary to each other; or obliquely joyn'd when they are oblique, so as to produce a new Motion compounded from the Determination of both.

11. LET the Body A \* receive an Impulse in the Direction A L, \* Pl. 25. F. 7. so as to go thro' any determinate Space in a determinate Time; as for Example, the Space A B or one Rod ( $16\frac{1}{2}$  Feet) in a Second Minute of Time. The Body will, according to the first Law, by Virtue of the Force impress'd, go on uniformly thro' the Spaces A B, B C, C D, D E, E F, F G, G H, H I, I K, K L, &c. so as to describe each of them (supposing them all equal) in a Second, and so on *in infinitum*. Now when the Body is in Motion, if the same Force that acted upon it at first, or one equal to it, should act upon it again in the same Direction, when it is at B for Example; then would the Body be carried thro' a double Space, *viz.* the Space B D, in a second, and so thro' D F, F H, H K, &c. in every Second, that is, with double Velocity; because, whilst it was going on at the Rate of one Rod in a Second, it receiv'd an Addition of Force capable of making it go also a Rod in a Second, and consequently that Addition of those two Forces that conspire give the Body a double Velocity. If the Body in Motion, when at B, had received the Second Impulse by a Force double of the first Force, it would after that Impulse go on in each Second thro' the Spaces B E, E H, H L, &c. that is, with a triple Velocity. If after receiving the Second Impulse with a Force equal to the first, while it is going on with double Velocity (for Example, thro' the Space B D or D F, &c. in a Second) it should receive a third Impulse

Lect. V. pulse (still in the same Direction) equal to the first, then it would go from D to G, and so on from G to K, &c. every Second of Time, that is, with a triple Velocity, after three equal Impulses, just as it would do after two unequal Impulses in the Proportion already explain'd : as it would also do if it had receiv'd but one Impulse at first, but three times greater than we then suppos'd.

Pl. 25. F. 7.

12. IF, whilst the Body by the Force impress'd is (by the first Law) going on uniformly thro' the Spaces A B, B C, &c. a Force equal to the Force impress'd should act upon it in the Direction L A (that is, in a contrary Direction) the Body would lose all its Motion. But if this last Force should only act upon it after it had receiv'd several Impulses according to the first Direction, it would only destroy so much of the Motion as its own Quantity of Force would be capable of producing : for Example, if the Body, after three Impulses, was going on thro' the Spaces D G, G K, &c. that is, 3 Rods in a Second, and then it should receive a fourth Impulse, but in the contrary Direction, it would only lose so much Motion as to go thro' but two Rods in a Second, and continue to go on uniformly with that Velocity, just as if it had receiv'd but two Impulses. Again, if when it is going 3 Rods in a Second it should receive two Impulses at once equal to the first, but in contrary Directions, one in the former and the other in the opposite Direction; those Forces would destroy each other, and the Body would go on uniformly with the same Velocity that it had before those Impulses, viz. 3 Rods in a Second.

13. Now, let us once more suppose the Body to be going on uniformly at the Rate of a Rod in a Second or thro' the Spaces A B, B C, &c. Suppose when it is at E, a new Force equal to the first (or equal to the Force which the Body has then) should act upon it in the Direction E e, or at right Angles to its present Direction A L; the Body would change its Direction, and move (as has been shewn) in the Diagonal E f of the compleated Parallelogram E e f F. † If the new Force had been double, then the Body would have gone in the Diagonal E f of the double Parallelogram E e f F. But if the new Force had been only equal to half of the first Force impress'd, then the Body would only have gone in the Diagonal E φ of the Parallelogram E e φ F.

† L. 3. N<sup>o</sup>.  
85. L. 5.  
N<sup>o</sup>. 3.

C O R O L L A R Y I.

HENCE it follows, that let the Quantity of the new Force be <sup>Pl. 25. F. 7.</sup> what it will, if it acts at right Angles, the Velocity will be greater than if the Body had continued in its first rectilinear Direction; because (as has been shewn †) the Body by the Action of the <sup>† L. 3. N<sup>o</sup>. 85. and Page 286, 287.</sup> two Forces describes the Diagonal in the same Time as it would have described either Side of the Parallelogram by the single Action of one of the Forces; and in the Parallelograms  $E\phi$ ,  $E\psi$ ,  $E\chi$ , the Diagonals are longer than the Sides, as they subtend the right (that is, the biggest Angles) of the rectangular Triangles  $E\phi\psi$ ,  $E\psi\chi$ , and  $E\phi\chi$ . \* N. B. *When we make use of these Words the Force impressed, or the innate Force, we mean the Force which the Body has when it is in Motion, without considering what first gave the Motion: That is, we mean the same thing that Sir Isaac Newton calls Vis insita.* <sup>By 19. 1. Eucl.</sup>

C O R O L L A R Y II.

HENCE follows also, that if the new Force acts at acute Angles, its Effect will be more considerable the acuter the Angle is; but it will never encrease the Velocity of the Body so much as if it acted at no Angle, that is in the same Direction. For Example, supposing the new Force equal to *the Force impress'd*, and its Direction to be in the Line  $Fg$ ,  $GFg$  will be the acute Angle made by the two Directions. Now it is evident, that the less the Angle  $GFg$  is, the greater will be the Angle  $FGg$ , and consequently the Diagonal  $Fg$  (expressive of the Body's Velocity) which subtends that Angle will also be greater: but whilst there is any acute Angle at  $F$ , there will be an obtuse Angle at  $G$ , and  $GFg$  will be a Triangle in which one side  $Fg$  will always be less than the Two other Sides  $FG$  and  $Gg$ , † which are equal to the Two  $FG$  and  $GH$  thro' <sup>† By 20. 1. Eucl.</sup> which the Body goes, when the Two Forces act in the same Direction, in the same Time.


C O R O L L A R Y III.

IT follows likewise, that the new Force may act upon the Body in such a Direction, that the Body will not be accelerated thereby; nay sometimes retarded. For if the new Force neither conspires

Leſt. V. with the Force impreſſed, by acting at acute Angles with its Directions, nor acts at right Angles with it; the Body may keep its Velocity tho' in a new Direction, nay even go ſlower when the new Force acts at obtuſe Angles. Let us, for Example, ſuppoſe the new Force, equal to the Force impreſſ'd, to act when the Body in Motion is at K, in the Direction K k, ſo that the obtuſe Angle L K k may be of 120 Degrees: then will the Diagonal K l of the Parallelogram K k l L be equal to the Side KL, becauſe K L l and K k l are two equilateral Triangles, and therefore the Body will have the ſame Velocity as before. But if (every thing elſe remaining as we have ſuppos'd) the new Force ſhould be leſs than the Force impreſſ'd, as for Example, only equal to half of it, the Velocity of the Body would be diminifh'd: for then K k would repreſent the new Force and K l would be the Diagonal of the Parallelogram (which would now be K L l k;) and K l falling from K one Angle of an equilateral Triangle upon the Middle of its oppoſite Side would be ſhorter than any of the Sides; *therefore*, &c. If the new Force was more or leſs than half in any Proportion, the Velocity of the Body would indeed be leſſen'd, but not ſo much; for then the Point l would fall between L l, or between l L, and in any of thoſe Caſes the Diagonal would be longer than K l becauſe K l is perpendicular to L l, and the ſaid Diagonal would be oblique to it, therefore longer, tho' ſtill ſhorter than the Side K L.

If the new Force had been greater than the Force impreſſed, for Example double, the Body then (ſuppoſing the Angle of Application the ſame) would encrease its Velocity by the Action of it, for the new Force being repreſented by K 1, the Diagonal or Velocity would be K 2 longer than K L. But then if the Direction of this greater Force ſhould be more oppoſite to the Direction of the Force impreſſ'd, that is, if the Angle L K 1 ſhould be encreaſed, for Example 30 Degrees more, ſo as to become L K 2, then the Body would not change its Velocity with its new Direction, becauſe it wou'd go in the Diagonal K k, which is equal to K L. Nay, if the Angle of Application ſhould be greater than L K 2, or of more than 150 Degrees, the Velocity of the Body would be diminifh'd, the new Diagonal becoming ſhorter than K k.

If the new Force was only equal to the Force impreſſ'd and its Direction K 4, that is, if the Angle of Application ſhould be of 150 Degrees, the Velocity of the Body would be diminifh'd by half, the Diagonal K 5 being then but the half of K 1 (= K L) becauſe  
it

it subtends the Angle  $KL$  5, which is but half of the Angle  $KLl$ . *Lect. V.*  
*N. B. Knowing the Angle of Application and Quantity of the*  *new Force, we may always know what will be the Velocity of the* *11. 25. F. 7.*  
*Body after the Action of the new Force; because the Diagonal*  
*that shews it, is always known; as it is one Side of a Triangle,*  
*in which Two Sides representing the two Forces are given, and*  
*the Angle between them, and consequently the 3d Side, † which is* *† By 15. and*  
*the said Diagonal. The Angle known does always contain the* *17. 1 Euclid.*  
*Number of Degrees that the Angle of Application wants of*  
*180. Thus in the Triangle  $KkL$ , where  $KL$  and  $Lk$  ( $= K_2$ )*  
*represent the Forces; (because of the Parallels  $K_3$  and  $Lk$ )*  
*\* the Angle  $KLk$  is equal to  $IK_3$ , which  $LK_3$  wants of 180°* *\* By 29. 1.*  
*Degrees; and consequently the Diagonal  $Kk$ , which subtends that* *Euclid.*  
*Angle, is known.*

14. It has been observ'd that when a Body falls during one Second of Time, it goes thro' a Space equal to  $16 \frac{1}{2}$  English Feet, or one Rod, as we have already mentioned: \*\* therefore the Force impressed by Gravity at the Beginning of its Fall is capable of making the Body go downwards at the Rate of one Rod in a Second, † tho' † By the first Law, Page 286, 287.  
 it should act no longer upon it than during the first Second; that is, tho' the Body should for ever after cease to be heavy. For example, if the Body  $A$  \* falls thro' the Space  $AB$  during the first \* Pl. 25. F. 8.  
 Second of its Fall; if then it should cease to be heavy, yet it would go thro' the Spaces equal to  $AB$  during all the succeeding Seconds of Time, viz. thro' the Spaces  $Bc$ ,  $cD$ ,  $De$ ,  $ef$ ,  $fG$ ,  $Gh$ ,  $hi$ ,  $ik$ ,  $kL$ , &c. But as the Body does not cease to be heavy, we must consider the Action of Gravity as an Impulse given by a new Force equal to the first, acting downwards when the Body is got to  $B$  at the Beginning of the 2d Second, and the Body during the 2d Second will go thro' the Space  $BD$  double the Space  $AB$ , or equal to the Two Spaces  $Bc$  and  $cD$ . Then if the Body should cease to be heavy, it would go uniformly thro' the double Spaces  $Df$ ,  $fh$ , &c. every Second; but Gravity acting upon it also at the Beginning of the 3d Second, when it is at  $D$ , superadds a Force able to make it go thro' a Space equal to the first  $AB$  in a Second: consequently it will go thro' 3 Spaces or Rods (or thro' a Space  $DG$  equal to 3 Rods) the 2d Second. At the Beginning of the fourth Second, Gravity acting by a fourth Impulse superadds a Force equal to the former, whereby it will go the Length  $GL$ ,  
 that

\*\* Bodies in reality only fall 16 English Feet and one Tenth of a Foot in a Second; but we call that Space here 16 and a half Feet; because one Rod (which is a Measure of 16 and a half Feet) gives us such a Number as to avoid Fractions in the Examples of the Calculations that we give.

Lect. V. that is, four Rods or Spaces the fourth Second : and so on ; † and  
 this will be a Motion uniformly accelerated.

† N<sup>o</sup>. 11.

THIS Motion of a Body thus accelerated in its Descent would be the real Motion of falling Bodies, if Gravity acted by Intervals, as we have only suppos'd, it helps Conception : But as Gravity never ceases to act, we must fill up the Intervals between the Beginning and End of every Second or small Part of Time.

\* Pl. 25. F. 9. Thus, if we consider the Body falling thro' the Space A B \* (Fig. 9.) in the first Second, we must not only consider Gravity superadding a Force capable of making the Body fall one Rod farther at the Beginning of every Second ; but also during the Time of every Second. For Example, at B at the Beginning of the 2d Second, the Body not only receives an additional Impulse to carry it to d instead of c, but also during the Time of the 2d Second another Impulse which makes it go to E instead of d ; so that the Body will fall 3 Rods during the 2d Second. Likewise at the Beginning of the 3d Second when the Body is at E, it will again receive from Gravity one Impulse at the Beginning of that Second, and another during the Time of it ; so that it will go thro' the Five Spaces E f, f g, g h, h i and i K during the Time of that 3d Second. So in the Time of the 4th Second the Body will (for the same Reason) go thro' seven Spaces or Rods from K to R, and so on, the Number of Spaces describ'd encreasing by two every Second ; that is, according to the Series of the odd Numbers 1, 3, 5, 7, 9, &c.

THIS is the true Manner of Bodies falling with an accelerated Motion by the Force of Gravity, abstracting from the Resistance of Air, which we shall hereafter take into Consideration. N. B. *All Bodies fall equally swift where there is no Air, as has been prov'd by the Experiment of a Piece of Gold and a Feather falling in the same Time from the same Height in an exhausted Re-*

ceiver †.

† L. 1. N<sup>o</sup>. 8.

15. WHAT we have shewn (Step by Step) of the Effect of Gravity on falling Bodies, may be demonstrated another way after *Galileo's* Method, thus. As Bodies (abstracting from the Air's Resistance, as we have said) encrease their Velocity in falling according to the Time during which they fall, *Galileo* represents the Times and Velocities by equal right Lines, which he joyns at right Angles ; and then joyning their other Ends by a third right Line, makes a rectangular Triangle of which this last Line is the Hypotenuse

Hypotenuse to represent the Space fallen thro' in a certain Time ; *Lect. V.*  
*A B*, for Example,  $\dagger$  represents the Time ; *B V* represents the Velocity ; and the Triangle *A B V* the Space gone thro' during that Time ; as for Example, one Rod, if the Time be one Second.  $\sim$   
 $\dagger$  Pl. 25. F. 10.  
 Now if the Time be double, *A C* represents it ; and then the Velocity being double also, the Line *C U* must represent it. Draw *U A*, and the large Triangle *A U C* will represent the Space fallen thro' in that Time ; which greater Space being divided into Triangles equal to the First *A B V*, will appear to contain four of them. Therefore, since a Body falling one Second of Time goes thro' the Space of one Rod, the same Body falling two Seconds will go thro' the Space of four Rods. Had the Time been *A D* ( $= 3$  Seconds) the Velocity wou'd have been *D v*  $= A D$ , and the Space *A v D*  $= 9$  Rods, or 9 of the First Space. Thus if the Time be *A E*  $= 4$  Seconds, the Velocity must be *E u* quadruple of *B V*, and the whole Space *A u E* containing 16 times the First Space *A V B*.

### C O R O L L A R Y I.

HENCE it follows, that if you multiply the Times by the Velocities, or if you square the Times, or square the Velocities, you will have the Number of Spaces (equal to the First) which the Body has describ'd during its whole Fall. As here, the Times being 4, and the Velocities 4, the Spaces gone thro' are 16.

### C O R O L L A R Y II.

HENCE follows also, that if you mark the Times (or Number of Seconds) as the Numbers, 1, 2, 3, 4, in the Scheme shew ; you will have, against any of the particular Numbers, as many little Triangles as the Body has gone thro' Spaces during the Part of Time express'd by that Number. For Example, there is on the left Hand One Triangle over against the Number 1, Three over against 2, Five over against 3, and Seven over against 4. And to know how many Rods a Body falls in any determinate Second or Part of Time, as for Example, in the 10th Second, without drawing the Scheme ; we must first find how many Rods the Body falls in the whole Time, as here in 10 Seconds ; then how many Rods the Body falls in 9 Seconds, and subtract the last from the first.  $10 \times 10$  is 100 the Number of Spaces in 10 Seconds :  $9 \times 9$  is 81 the Number of Spaces in 9 Seconds ; 81 from 100 leaves 19, the Spaces or Rods fallen thro' in the 10th Second.

C O R O L L A R Y



## C O R O L L A R Y III.

† Pl. 25. F.  
10.

It appears likewise, as a Consequence from what has been said, that when a Body has fallen thro' a certain Space by a Motion uniformly accelerated, it has acquir'd such a Velocity as is able to carry it thro' a double Space in the same Time, if Gravity should cease to act, or have its Effect any way destroy'd, or if the Body's Direction should be chang'd from a vertical to an horizontal one. For Example, when the Body in the Time  $A B$  †, falling thro' a Space represented by the Triangle  $A V B$ , has acquir'd the Velocity  $B V$ ; if Gravity ceasing to act, it receives no Addition to its Velocity, tho' it continues to fall during the Time  $B C = A C$ , so that the whole Time must be represented by the whole Line  $A C$ ; yet the Velocity will not be represented by the equal Line  $C U$ , but by the Line  $C k$  equal to  $B V$  the Velocity the Body had when it ceas'd to accelerate its Motion; therefore to know how many Spaces the Body has gone thro' in the Time  $B C = A C$  with an uniform Motion, we must multiply  $B C$  by  $C k = B V$  the unchang'd Velocity of the Body, and we shall have the Rectangle  $B V C K$  containing two Triangles equal each to  $A V B$ : that is, two Spaces will be run thro' by the Body moving with an equal Motion in the Time  $B C$  equal to the Time  $A B$  in which the Body describ'd only one Space by an accelerated Motion. So likewise, if the Body having fallen thro' 4 Spaces during the Time  $A C$ , and having at the End of that Time acquir'd the Velocity  $C U$ ; the Velocity be no more encreas'd and the Body continues to fall during the Time  $C E = A C$ , it will fall thro' 8 Spaces instead of 4 that it wou'd have fallen if the Velocity had continued to encrease so as to become  $E u$  at the End of the Time  $C E$ ; but as it is then only  $E e$ , equal to  $V C$  its Velocity at the Beginning of the Time  $C E$ , we must multiply  $C E$  the Time by  $U C$  or  $E e$  the uniform Velocity, and then we shall only have the Rectangle  $V C E e$  containing 8 Spaces instead of the Trapezium  $C U u E$  containing 12.

THIS will become visible by changing the Direction of the moving Body from a vertical to an horizontal one, for Gravity acting only perpendicularly will neither accelerate nor retard the Body in its horizontal Motion. N.B. *The manner of doing this we shall hereafter shew.*

C O R O L L A R Y. IV.

WE may also further gather from what has been said, that Gravity does not act by Intervals (tho' we explained it that way at first to facilitate Conception) for if it did, after every Impulse the Body acted upon wou'd go on with an uniform Velocity, tho' it would have a greater Velocity after every Impulse. Neither after it had fallen a certain Space would it have acquir'd a Velocity capable of carrying it thro' double that Space in the same Time, should Gravity cease to act, but only thro' a Space equal to the first.


FOR Example; after *Galileo's* Method, the Triangle A C B Pl. 25. F. 11. (*Pl. 25. Fig. 11.*) represents the Space fallen thro' in any Part of Time, suppose one Second, in the Time A B with a Velocity *nascens* (or just beginning) at A the Beginning of the Second, and equal to C B at the End of the Second: whereas if a Body should fall one Rod in a Second by an Impulse as a Blow, the Velocity at first would be equal to A D and continue so during the whole Time, so that at the End of the Second it would only be B V equal to A D and half of B C, whereby without a fresh Impulse it would only carry the Body down one Rod more in another Second; for then the Time multiplied into the Velocity would only produce the Rectangle A D V B equal to the Triangle A C B, because it is of the same Height and has half its Base. †

† By 41. 1. Euclid.

16. IF the Direction of a falling Body be so chang'd as to make it go directly upwards beginning that Motion with all the Velocity it had at the End of its Fall, it will go up (by a Motion uniformly retarded) exactly to an Height equal to that from which it fell; and the Spaces it goes thro' in each Part of the Time are to be seen in the Scheme over-against the Numbers representing those Parts of Time. If, for Example, the Body has fallen 4 Seconds, it will have gone thro' 16 Rods in its Fall, and have acquir'd the Velocity E u † capable of carrying it 32 Rods in the same Time by an uniform Motion; but as Gravity acts against it in its Rise it will † Pl. 25. F. 10. destroy all its Motion by the Time that it has risen only 16 Rods; for it is the same thing to make a Body from Rest move 16 Rods downwards in a certain Time, as to destroy half of the Force which was able to carry a Body upwards 32 Rods in the same Time. The same Thing will also be plain if you consider the whole Time,


T t

as


 Lect. V. as divided into small Parts, and observe the uniform Diminution of the Body's Motion. For Example; let the Body be thrown upwards, and let the Spaces which it goes thro' in every Second be consider'd. If the Velocity given to the Body projected upwards be such, that in the Time  $ED$  (which will now be the first Second of Time) it goes thro' 7 Rods, the next Second it will go but 5, because Gravity gives it an Impulse downwards at the Beginning of, and during, the 2d Second, able to carry it downwards 2 Spaces or Rods; which is (in other Words) to take away 2 of the 7 Spaces, which the Body would have gone thro' in that Time. Thus likewise, instead of going 5 Spaces the 3d Second, Gravity taking away 2, it will go but 3 Spaces; and the 4th Second (the last of its Rise) instead of going 3 Spaces, it will for the same Reason go but one; and then the Body will for an indivisible Moment of Time be at Rest. From the Point of Rest the Body will come down again, falling with an accelerated Motion already describ'd, and coming down just in the same Space of Time that it went up.

*S C H O L I U M. Pl. 25. Fig. 12.*

Pl. 25. F. 12. If a Body, instead of going upwards in the Line  $AB$ , should rise to the same Height in the inclin'd streight Lines  $AC$  or  $AE$ , or in the Curves  $AaC$  or  $AeE$ , it would in the same Time come down in the streight Lines  $CD$  or  $EF$ , or in the Curves  $CcD$  or  $EfF$ . For we have already shewn, that when a Body is acted upon by a Force to carry it in the Line  $AB$  in a certain time, if another Force  $Ag$  acts upon it too, the Body will go in the Diagonal  $AC$  of the Parallelogram  $ABCG$  in the same Time: And we have also shewn, that it would go in a Parabolic Curve  $AeC$  in the same Time from  $A$  to  $C$ . Now since the Force of Gravity in the Direction  $BA$ , which destroys the Motion of the rising Body, is equal to the Force of Gravity acting in the Direction  $Cg$  to make the Body fall from its Point of Rest  $C$ , the Forces  $CE$  and  $Cg$  are equal to  $Ag$  and  $AB$ , and therefore the Diagonal right Line  $CD$ , or curv'd Line  $CcD$ , will be gone thro' in the same Time as the Lines  $AC$  and  $AaC$ . The same might be said of the Lines  $AE$  and  $AeE$  compar'd with  $EF$  and  $EeF$ . *N. B. When we throw a Body directly upwards and it falls upon the same Place of the Earth we threw it from, then it really describes Two such Lines as  $AC$  and  $CD$  moving from  $A$  to  $D$  by the Motion which the Earth gives it from East to West. But if we shoot it obliquely*

*liquely upwards it describes the two Curves A C, C D, or the Pa- Lect. V. rabola A C D; or any other Parabola, as A E F, according to the Angle of the Direction it sets out with.* 

C O R O L L A R Y I.

HENCE it follows, that one may at any time know to what Height a Projectile, as a Bomb, or Cannon Ball, &c. (whether shot up directly or obliquely) has risen. For if you take the Time between the firing the Mortar and the falling of the Bomb, the half of it will be the Time of the Fall of the Bomb. Square the number of Seconds in that Time, and you will have the Rods or Spaces fallen thro' in perpendicular Height. Thus if 20 Seconds are elaps'd from the firing of the Mortar to the Fall of the Bomb, half that will be 10, the Square of 10 is 100, which hundred Rods multiplied by 16 and a half will give the utmost Height of the Bomb in Feet, *viz.* 1650.

N. B. *We still abstract from the Resistance of the Air, which we shall hereafter consider and shew how to allow for it, as likewise we take 16  $\frac{1}{2}$  Feet instead 16, 1 for the Reason above given.*

C O R O L L A R Y II.

HENCE also follows, that knowing the Weight of a Body and the Height from which it falls, one may know what Stroke it will give; that is, what *Momentum* it has at the End of its Fall; for the Square Root of the Spaces will always give the Velocity, which being multiplied into the Mass and Weight of the Body give us its *Momentum*. \*

\* L. 2. N. 3.

SOME People have imagin'd that a falling Body has a *Momentum*, and strikes a Blow, proportionable to the Height from which it falls; for Example, that a Pound falling from an Height of 4 Foot has four times the *Momentum* that it would have falling from one Foot. But their Error lies in not taking the Time into consideration; for a Body spends twice the Time in falling 4 Foot that it does in falling one Foot, so that the Velocity is only double in the first Case. Indeed if a Body could fall 4 Spaces, or in any Direction whatever go thro' 4 Spaces, while another Body of equal Mass went thro' but one Space; it would then have four times the *Momentum*, and consequently be capable of a quadruple Effect.

Lect. V.

Others again allow that the Velocity is as the Square Root of the Spaces; but alledge that the *Momentum* is not as the Product of the Mass by the Velocity of the moving Body, but as the Mass multiplied into the Square of the Velocity; which Opinion they endeavour to support by several Experiments, and various Reasonings, many of which I shall consider in the Notes.\* I shall only take notice here, that tho' I do not think the Experiments of hard Bodies falling upon soft Substances (to be hereafter more fully describ'd) conclusive in Respect of the *Momenta* of those Bodies; yet they have very well prov'd that yielding Substances or soft unelastick Bodies yield to a Blow of the same percutient Body in proportion to the Square of its Velocity; and thence may be drawn very useful Consequences applicable to the Practice of Mechanicks.

\* Ann. 4

\* L. I. An. II.

IN explaining the Action of Gravity upon falling Bodies we have consider'd it as acting upon Projectiles always with the same Force, tho' it is certainly weaker in its Effect the higher the Body acted upon is above the Earth, as we have shewn in the 11<sup>th</sup> Annotation to Lect. I.\* For Gravity (that is, its accelerating Force) decreases, as the Squares of the Distances from the Center of the Earth encrease. But as the greatest Height to which we are able (even with Gun-powder) to project Bodies, bears no Proportion to the Distance from the Center of the Earth, (scarce one 2000<sup>th</sup> Part) we could make no Allowance for that Decrease of the Force of Gravity, without making too much, since the Difference is insensible.

\* L. I. Ann. II.

17. IF the Force of Gravity was greater or less than it is here, Bodies wou'd be accelerated by it in their Fall in the same manner that we have explain'd; only the Spaces which the falling Body goes thro' in the same Time, wou'd be greater or less in Proportion. If the Force of Gravity was 4 times greater, a Body would fall 4 Rods in the 1<sup>st</sup> Second of its Fall; and if the Force of Gravity was four times less (as it wou'd be if we were carried up to the Height of 4000 Miles, or remov'd farther from the Center of the Earth than we are, the Distance of one Semidiameter of the Earth) a Body would fall but a quarter of a Rod in the first Second of its Fall.\* And if we observe, that in any Place on the Surface of the Earth a Body falls not so great a Space as one Rod in a Second; we may be sure that the Force of Gravity is less there

there than in our Country where Bodies fall one Rod in a Second. *Lect. V.*  
Now this Observation has been made very near the *Æquator*,  
where from Experiments made on Pendulums it appears that Bodies do not fall a Rod in a Second; and it follows from thence that the Force of Gravity is less there than in greater Latitudes: which happens because the Surface of the Earth is higher (that is, farther remov'd from the Center there than at the Poles) about 21 Miles. But I shall speak more fully of this in another Place. In the mean time, I refer the curious to Sir *Isaac Newton's Principia*, Book 3. Prop. 20. and the *Philosophical Transactions*, No. 386, 387, and 388.

18. WHEN a Body runs down an inclin'd Plane, it cannot descend with its whole Gravity, because Part of it is supported by the Plane, and that in Proportion of the Length of the Plane to its Height (or of the *Radius* to the *Sine* of the Angle of Inclination) as has been shewn in the 7th Annotation of *Lect. 3.* But then that Part of the Weight which is not supported by the Plane, or the *relative* Gravity being always of the same Quantity, because every Part of the Plane has the same Inclination, will cause the Body that rolls down to accelerate its Motion in the same manner as a Body that falls freely down, but not so fast; or in the same manner that Bodies would fall freely if the Force of Gravity was so much less. *L. 3. Ann. 7.*

### CONSTRUCTION.

UPON any straight horizontal Line as C B, \* raise a Perpendicular at C, and make A B the Hypotenuse of the Triangle A B C (which is to represent an inclin'd Plane) in Length equal to twice the Height or Perpendicular A C. Divide A B into 4 equal Parts mark'd by the Points D, F, G, and likewise A C into 4 equal Parts mark'd by the Points E, H, I. Draw E D and C D. \* Pl. 25. FF.

IF A E, E H, H I, and I C be the 4 Spaces (or Rods) which a Body falls thro' in 2 Seconds, A E will be one Space which the Body goes thro' in 1 Second. † Now if the Force of Gravity was but half of what it is, the Body, instead of going down from A to C in two Seconds, would only go from A to E. Let the Body be laid upon the Plane A B, and then it will be supported


**Lect. V.** supported by the Plane as to descend towards B with only half its Weight, that is, it will lose half its Weight and advance towards the Center of the Earth no faster than it would do in the Line A C if it fell freely down and the Force of Gravity was but half of what it is. The Body therefore rolling on the inclin'd Plane will come down but to D, which is upon the same Level (or as near the Center of the Earth) as E, in 2 Seconds. Whereas, if it had not been supported by the Plane, it would have fallen 4 Spaces or down to C. It has then at D the same Velocity which a Body falling freely from A has at E, but is (in this Case) twice as long in acquiring that Velocity. If the Body, when it is come down to D, continues to move along the Inclined Plane, it will for the Reasons already alledg'd go thro' the next 3 equal Spaces D F, F G, G B, in the next 2 Seconds of Time; just as a Body falling freely in the Line A C would do if the Force of Gravity was but half; or as a Body will do now with the whole Force of Gravity in half the Time: and the Velocity of the rolling Body when it is at B will be as great as the Velocity of the Body at C; but it will be so much longer acquiring it as the Line A B is longer than A C.

#### C O R O L L A R Y I.

HENCE follows, that whatever the Inclination of the Plane is, a Body will have acquir'd the same Velocity when it comes down to the Bottom of it as if it had fallen along the Perpendicular; and if any two Points (as E and D, H and F, I and G) be taken in the same horizontal Line, one upon the Inclined Plane and the other upon the Perpendicular, the Body will have the same Velocity in both, tho' acquir'd in different Times. For if the Plane be less inclin'd, the relative Gravity which carries it down will be greater, and accelerate the Body so much quicker as the Plane is shorter; whereas the more inclin'd or the longer the Plane is, the more slow its Acceleration will be.

#### C O R O L L A R Y II.

HENCE follows also, that if a Perpendicular be drawn from any Point of the Line in which a Body falls freely to the inclin'd Plane, it will shew how far upon the inclin'd Plane another Body would roll in the same Time, if the Fall of both began from the same Point.

**Point.** That is, in this Case  $C D$ , which from the Bottom of the Lect. V. vertical Line  $A C$  comes upon the inclin'd Plane at  $D$  (to which  Point we have shewn that a Body would fall along the Plane, in the same Time that it wou'd fall freely as low as  $C$ ) is perpendicular to the Plane  $A B$ . For since by Construction  $A B$  is the double of  $A C$ , and  $A C$  the double of  $A D$ , and the Angle  $C A D$  common, the Triangles  $C A B$  and  $C A D$  will be similar  $\dagger$  and consequently  $C D$  is perpendicular to  $A D$ . Pl. 25. F. 13.  $\dagger$  By 6. 6. Euclid.

### C O R O L L A R Y III.

HENCE also may be shewn, that if a Body shou'd fall down from the same Height along several Planes differently inclin'd one after another, as for Example the Planes  $A D$ ,  $D E$ ,  $E B$ ,  $\dagger$  it would have  $\dagger$  Pl. 25. F. 14. acquir'd the same Velocity as if it had fallen along only one Plane as  $A D$ , or the Perpendicular  $A C$ .

### C O R O L L A R Y IV.

IT is likewise evident that if a Body falls in an Arc of a Circle, it will have the same Velocity at Bottom, as if it had fallen in the Perpendicular; because an Arc of a Circle may be consider'd as an infinite Number of Planes differently inclin'd.

18 THE Doctrine of Pendulums is naturally deduc'd from what we have explain'd about the Fall of Bodies; but as there are some Things in it that require the understanding the 2d Law of Motion, we must defer it till that Law is explain'd. In the mean time we will give an account of those Mechanical Organs or Instruments which were omitted in the 2d Lecture (when we describ'd the simple Machines commonly, but erroneously, call'd *Mechanical Powers*): because the Principles explain'd in that Lecture were not sufficient for it, without the knowledge of the 1st and 2d Law of Nature.

19. THE seven simple Machines or Mechanical Organs which I describ'd in my Second Lecture are only Instruments to perform the same thing different Ways,—namely to convey the Powers exerted or spent by some Body acting, so as to be applied for the moving of other Bodies, or to transmit or regulate the Power from one Body to another. Now the Instruments I am about to describe will also serve

Lect. V. serve to perform this, therefore they may very well be call'd Mechanical Organs; but the difference in the acting is this: In the Machines or Organs already describ'd, a great Part of the Strength of the Men or other Animals, &c. (that is, the Intensity of the Power) is lost in the rubbing, sticking, stretching, wearing, and yielding of the Parts of the Machine, and the Force of the Men, &c. employ'd can only be exerted by degrees, with the loss aforesaid. But in the Machines I am about to describe, the Power is collected and convey'd from Body to Body with little or no Loss by Accumulation continuing in the same Line.

20. If a Man weighing 140<sup>lb</sup> presses with his whole Weight, or exerts a Force equal to his Weight, upon a Leaver or Balance-Beam equally divided by its Center of Motion, it is impossible for him to overcome a Resistance greater than 140<sup>lb</sup>, tho' he should act upon the Beam a whole Day with all his Force, the Man's Power or Force being destroy'd as fast as he exerts it; but if he can communicate his Force by Degrees to a Body that shall keep it all and exert in a Moment the Sum of all the Impulsions given by the Man at several times, that whole Force of the Body so collected, and as it were condens'd, will perform at once what the Man could never have done by an Engine in the common Way. Such a Body was the *battering Ram* of the Ancients, which was a very large Piece of Timber (one of which we have describ'd in the 3<sup>d</sup> Annotation of the 2<sup>d</sup> Lecture\*) headed at one End with Metal, either Brass or Iron, which was contriv'd various Ways to be supported, and by the conjoyn'd Strength of many Men made to be mov'd with the Metal-headed End forward; till having receiv'd and kept all the successive Impressions of the Force of the Men (which was wholly employ'd to move the Ram forward; because its Weight was suspended by Ropes or Chains from a Distance above it to allow it to swing freely it had by little and little acquir'd a certain Degree of Velocity; and with that Degree of Velocity it met with or struck the Walls or Fortifications of Cities or Castles, and thereby remov'd or beat them down.\* This Machine is by the Men

\* L. 2. Ann.  
3.

\* No doubt but the first Invention of the battering Ram was copied from Nature, rather than mathematical Reasoning; that is, from what a Ram is observ'd to do by Instinct. If that Creature, having in vain push'd with his Head against an Obstacle by help of the Muscles that extend his Legs, his hind Feet being fix'd hard against the Ground, makes a second Attempt; and, by a sudden Stroke of his Head, removes the Obstacle which resist'd too much before; in such a case afterwards, he retires

Men accelerated in an horizontal Direction in the Manner that Lect. V. falling Bodies are accelerated by Gravity in a vertical one. The Man therefore that should be unable to overcome a Resistance greater than 140<sup>lb</sup> by the common mechanical Organs, might do it by a heavy Body swinging horizontally in such a Manner that he could give it a Motion accelerated till it reach'd the Obstacle; that is, he could do it by means of a small battering Ram. And if it was required to have a Force impress'd downwards, he might overcome the Resistance by the *Rammer* or heavy Body made use of to drive Piles into the Earth; where applying his Force successively against Gravity (that is, raising the Body to a certain Height) the Rammer would be put into such a Situation as to receive an accelerated Motion from Gravity before it acted upon the Resistance, which at last it would do with a great Force, because the same Quantity of Matter would move with a great Velocity; so much the greater, as it had spent more Time in falling, by reason of the Height it had been raised to. N. B. *It would be wrong to alledge here, that a Man, who could not raise a Weight, or overcome a Resistance something greater than 140<sup>lb</sup> with the Force 140 by means of a Leaver or Beam whose Brachia are of equal Length, might do it by applying his Power farther from the Center of Motion; because we are to suppose the Weight or Intensity of the Resistance to be always in a Proportion to the Power something greater than the reciprocal Proportion of Distances: that is, we are to suppose in this Comparison, a Case impossible in the common Use of mechanical Organs.*

20. THE Mall or Hammer, which is a Body or Mass of Wood or Metal directed or mov'd circularly or in a curve Line by the Handle, so as by degrees to receive a certain Velocity and thereby to have a Power of moving, or making Impressions on, other Bodies, partakes of the two above mention'd Organs, as it receives an accumulated Force from the Arm that moves it, and at the same Time from Gravity, when it is made use of to strike downwards. Tho' it is mov'd circularly, yet its Stroke is made in a straight Line, namely in the Tangent to the Curve in which it moves, just at the Point of the Curve in which the Blow is made;

*retires so far as to be able to accelerate his Motion to the utmost that his Muscles can exert their Action in running, and thereby gives a prodigious strong Blow by the accumulated Force with his arm'd Head, to the Thing or Animal which he runs against; his Horns being so plac'd and fix'd on his Head, that he feels no Pain from the Shock.*

Lect. V. because all Bodies mov'd in a Curve endeavour to fly off in a Tangent to that Curve as we have already shewn.\*

\* No. 5.

21. THE Fly (some of whose Uses I have already consider'd in the 4<sup>th</sup> \* Lecture) is an Organ whereby the Body mov'd is caus'd to circulate or move round about a Center or Axis, and thereby is capable of accumulating the Powers impress'd upon it by degrees, so long as shall be requir'd. This differs from the Hammer and the Mall in this Particular, as the Capstane differs from the Leaver; for whereas the Leaver can only lift or remove the Body for a very short Space, the Capstane can perform it for a given or requir'd Space. So that the Hammer or Mall can only receive so much Power as can be given it in the Part of an Arc or Circle, or in a streight Line of a short Length; but the Fly can acquire its Power by Accumulation in many Revolutions, and so is made receptive of a given Power and Velocity, and can exert the same on any other Body in a determinate Manner.

22. THIS mechanical Organ joyn'd to the Screw composes that powerful Machine whereby they impress the Stamp or Image upon Coins and Medals, which requires a prodigious as well as regular Force and Power. Now in this Engine there are accumulated Powers, by three mechanical Organs; first, By the Fly, whereby the Strength of the Man moving it is accumulated into the Weights at the End thereof: secondly, That Power accumulated is condens'd and impress'd on the Cylinder of the Screw by means of the *Radii* of the Fly, which are two Leavers that serve to condense the same in a given Proportion: thirdly, That Power so communicated to the Cylinder accumulated and condens'd, is again condens'd by the Slope of the Screw in a given Proportion; and so the whole Power (or all the successive Powers) exerted by the Man in moving the Weights of the Fly, is accumulated and condens'd in the last Impulsion, which is made upon the Medal.

23. ANALOGOUS to the Fly may be accounted the *circular Pendulum*, which is a mechanical Organ, whereby Motion may be accumulated into the Body or Weight thereof; which is suspended by a String from a Center above, that keeps it at due Distance from the said Center.

By this Organ, the Weight to be mov'd by the impress'd *Leſt. V.* Force, is more free to receive and retain the accumulated Power, and receives none of the first Impediment of the Fly, namely from the rubbing and wearing of the Pivots or Gudgeons, and is only subject to the latter, namely the Impediment of the Air or of the Medium thro' which it is mov'd, which is also less, by reason that the String by which the Weight is suspended is less than the Arms or Branches of the Fly; and hence it is, that this Organ preserves the impress'd Force much longer than the other, namely the Fly, and will continue in Motion much longer. When set a going it will often continue to move many Hours together; and, if the Length of the spiral Space in which it moves be computed, it will amount to several Miles in Length. *But not having yet given any account of the common or simple Pendulum, I must defer saying any more of the circular one till I come to explain the simple one.*

24. THE last of these kind of Organs that I mean to describe now is the *Sling*; for the Bow in its Action depends upon the *3d Law of Motion*, and therefore cannot properly be treated of till we have explain'd that Law. The *Sling* then is an Instrument or Organ which serves, by the means of Strings to convey gradually the Strength of an Hand moving in a small Circle to a Body detain'd by these Strings, to move in a greater Circle about the same Center, till it has fully accumulated in it all the Power of the Hand, or the design'd Quantity of it, at which Time it discharges it with a design'd Direction and Determination. This Instrument partakes of the Fly and the circular Pendulum; for like the Fly it can receive an Accumulation of Power for many Revolutions, and like the circular Pendulum it is free from the Friction of Pivots and Gudgeons; and none of the Force apply'd to it is lost, but just what is sufficient to overcome the Resistance of the Air.

It wou'd be too long to describe the various Ways that this Organ has been contriv'd and made Use of by the ancient as well as the modern Engineers. It is sufficient to observe that they are all reducible to this (the common Sling) which is the most simple; and the Power and Effect thereof is as easy and reducible to Geometrical Computation and Calculation as any of the others already mention'd. There being only one way applicable to all; and it is to

Lect. V. find what the Velocity of the Body is just at the Instant that the Percussion is made, † or the Machine apply'd to Use. For it being always suppos'd that we know the Weight of the Body or Instrument, we have nothing to do but to multiply it by the Velocity and it will give us the *Momentum* of our moving Body, † and thereby shew the Effect which it is able to produce in acting upon another Body, in stopping, driving, breaking or striking it, so as to shake, move, or remove the whole or any part of it.

† Ann. 4.

† L. 2. N<sup>o</sup>. 2, 3.

25. Now all the Difficulty consists in finding what is the real Velocity of the percutient Body just at the Moment of Percussion; and it may be done in the following Manner, which we will first shew with respect to the Rammer or heavy Weight to drive Piles, it being most easily considered in that Instrument, and all Cases deduced from that.

\* Pl. 25. F. 15. THE 15th Figure of Plate 25 \* represents an Engine to drive Piles, consisting of the Cill K I and Frame F L, on which are fix'd the upright Pieces L H and L G, supported by the Side Braces C, C, and the hind-Brace F E (which has Pins on it to make it serve as a Ladder) and held together by the square Collar E D. The Rammer A being a very heavy Piece of hard Wood or Iron slides up and down between the Cheeks or upright Pieces L H, L G, and is drawn up by means of its Hook B with two Ropes H O, G O, having each 5 smaller Ropes with Handles at N, N, for 10 Men to pull up the Rammer to a certain Height (the great Ropes running over two Pullies or Rollers on the Iron Pin H G) and then let it fall again all at once upon the Head of the Pile at M to drive it into the Ground. Now suppose the Rammer A weighs 500 lb and falls the Height of one Foot, it will fall that Height in a quarter of a Second, and consequently have a Velocity able to carry it uniformly 2 Foot in the same Time, † (that is, at the Rate of 8 Foot in a Second) at the very Instant that it strikes the Pile M. Therefore multiplying the Mass by the Velocity, viz 500 × 8, we shall have 4000 for the *Momentum* of the Rammer, with such a Fall. But if the Rammer be rais'd up to the Height of 4 Foot, it will fall that Height in half a Second, and have at the Time of Percussion a Velocity to carry it 8 Foot in half a Second, without any farther Help from Gravity; so that we must now multiply 16 Foot (the present Velocity, since it goes at the Rate of 16 Foot in a Second) by 500 the Mass of the Rammer, which will give us a double

† No. 15. Cor. 3. P. 324.

double *Momentum* wherewith it will strike the Pile in this last Lect. V. Case; for  $500 \times 16 = 8000$ . If we consider any other Height from which the Rammer falls (for one may employ a Capstane, Windlafs, or Pullies, to raise it to a very great Height, and then by an easy Contrivance loosen it at once from its Hook) the *Momentum*, with which it strikes the Pile, will always be as the square Root of the Height from which the Rammer fell; that is, as the Velocity which the descending Body has at the End of its Fall. N. B. *I cannot say but that the Pile may enter into the Earth sometimes farther than in that Proportion; but I shall shew the Reason of it in the Notes.\**

\* Ann. 4.

26. IF a Pile is to be driven obliquely; the Engine must be set so that the Cheeks may have the same Obliquity, and the Blow will still be perpendicular to the Head of the Pile; but then the Force of the Blow must not be estimated from the Length, but from the Height, of the Descent, in the manner already shewn; because how long soever the inclin'd Plane is, in which the Body falls, it has acquired no more Velocity than what it would do if it had only fallen perpendicularly from the Height of the Plane.

27. To find the Velocity of the battering Ram when it makes its Stroke, we must observe at what Rate its Motion is accelerated; for according to the Number and Strength of the Men that work it, it may be accelerated more or less than Gravity would accelerate it if it was to fall perpendicularly. Therefore we are to observe the Length of the Stroke from the Point most remote from the Wall (or the Thing batter'd) and the Wall, and take notice in what Time the Stroke is made (for when the Men, by a little Practice, have got the Knack of playing the Ram, all their Strokes are made in the same Time) if the Stroke, for Example, be of 4 Foot and made in a quarter of a Second, the *Momentum* of the Ram is equal to what it would be if the same had mov'd uniformly 8 Foot in a quarter of a Second, or 32 Feet in one whole Second. This Force would be quadruple of what Gravity might give the falling Ram (or a Rammer equal in Weight to the Ram) in the same Time; but only equal to what it would give it in one Second or the quadruple of the Time; and only half of what Gravity would give in two Seconds to the same Body falling from an Height of a little more than 64 Feet. If the Time of making the Stroke had been twice as long, or half a Second, then the *Momentum*

Lect. V. *tum* or Force of the Percussion would have been but half as great  
 ~~~~~ &c. the Percussion with the same battering Ram being always in-  
 versely as the Time in which it is accelerated by the same Force
 of Men.

28. To find the Velocity (and consequently the *Momentum*) of the Mall or Hammer at the instant of Percussion, we must consider it first in the most simple manner, as when being rais'd up it falls down again in an Arc of a Circle by its own Gravity: then we are only to consider the Height from which it fell; and we shall know, from what we have said of Bodies falling perpendicularly or on an inclin'd Plane, what Velocity it has at the end of its fall. Thus we may know with what Force the great Hammers, rais'd by the Axis of a Water-wheel, fall upon the Plates or Bars of Iron or Copper that are flatten'd in Iron or Copper Battering Mills or Forges. When the Hammer is mov'd by the Hand, or driven by a Spring as well as by Gravity, it will move quicker, and its Effect will be proportionably greater, therefore by observing the Time of its Fall in such a case, its Velocity may be known. *N. B.* Tho' a Body, moving twice as fast in the same Circle, was able, according to what appear'd in the 15th Experiment of this Lecture, to raise 4 times the Weight; we must not imagine that a Hammer mov'd twice as fast will strike with 4 times the Momentum, whilst a Hammer of the same Weight moving twice as fast, only because of an Handle or Radius twice as long, shall have but double the Momentum according to the 13th Experiment; for it is only the centrifugal Force in the same Circle, which is as the Square of the Velocity, and not the Stroke made along the Tangent. The Effect of the centrifugal Force will only be this, that the Hand which holds the Hammer will feel 4 times the Force endeavouring to pull the Hammer out of his Hand; and would feel but twice that Force if the Velocity was only double by having an Handle twice as long.

It is also to be observed, that in the use of the Hammer it is better to strike with the same Mass of Matter with a double Velocity, than with double Mass and single Velocity; because yielding Substances, as hot Iron, &c. and Nails driven into Wood, give Way to the same Hammer nearly according to the Square of its Velocity, as we shall account for in the Notes. But the Re-*
verse

* Ann. 4.

verse must be done in the Battering Ram, if you would have its greatest Effect. †

Lect. V.
† Ann. 4.

29. IF the Fly be made use of only to give a Blow with one of its Weights after some Revolutions, the Method of finding its Velocity, and consequently its *Momentum*, just at the Stroke, is the same as will serve for the Sling; for tho' the Fly has vastly more Friction than the Sling, we are only to observe what Velocity (without any regard to the Impediments which hinder'd the Velocity from being so great as it would have been without them) it has just at the Stroke; which we may know by comparing several Revolutions, or Parts of Revolutions, together; which will shew us the Degree of Acceleration.

30. Now to shew how to calculate the Force of these Organs when joyn'd with others commonly called the *mechanical Powers* (explain'd in the 2d Lecture) I shall shew what Force may be given by the Machine made of the Fly combin'd to the Screw for stamping the Image upon Coins, as we have above hinted.

LET us suppose the two Arms of the Fly to be 15 Inches long each (measuring from the Center of the Weights to the Axis of Motion) and the Weights to be 50 lb each, and the Diameter of the Axis pressing upon the Dye to be one Inch. If every Stroke be made in half a Second, and the Weights describe an half Circumference, which will in this Case be of 4 Foot, the Velocity will at the Instant of the Stroke be at the Rate of 8 Foot in a Second, and therefore the *Momentum* of it will be 800; but the Arms of the Fly being as Leavers, one *Brachium* of which is 15 Inches long, whilst the other (which is the semi-Axis) is but of half an Inch, we must encrease this Force 30 times, which will give us 24000: an immense Force, equal to 100 lb Weight falling 120 Foot, or near 2 Seconds in Time, or a Body of 750 lb falling 16 $\frac{1}{2}$ Foot or one Second in Time. Some of these Engines for coining Crown Pieces have the Arms of the Fly 5 times as long, and the Weights twice as heavy; and the Effect is 10 times greater.

N. B. *We have allow'd nothing here for the inclin'd Plane of the Screw, because that Declivity only serv'd to help to accelerate the circular Motion of the Weights, which we consider'd in taking the Time of the Description of the Semicircle by them.*

Lect. V. 31. IT would be endless to shew all the Consequences of the two Laws of Motion already explain'd, in the Practice of mechanical Operations; and to apply them to the Explication of all kinds of Motions whether of Bodies on Earth, or of the Planets and Comets in the Heavens: We shall only give a few Instances more; but first we must shew how far the Resistance of Air (which we have hitherto left out of our Computations) hinders the Effects from being such as might be expected from their Causes without it.

THO' the Resistance of *Mediums* is to be consider'd in the Hydrostatical Part of this Work, yet we must say so much of it now as is requir'd for understanding how to allow for the Disturbance that the Resistance of *Mediums* gives to moving Bodies, whether their Motion be owing to Gravity, or any other Cause or Causes.

31. WHEN a Body moves in a Fluid of any kind whatever, or resisting *Medium*, it cannot go on without separating the Parts of the *Medium* to make it self Way; and so much as it bestows of its own Motion on those Parts, so much it loses of its own Motion; so that it will be retarded if its Motion was uniform before; or if it went on with an accelerated Motion, that Resistance (or the Motion given to the Parts of the *Medium*) will hinder the Acceleration from being so great as it would have been, or (according to the Quantity of it) destroy the Acceleration; that is, destroy the additional Motion as fast as it is given to the Body by the accelerating Cause: so that the Body will then move uniformly; as if the accelerating Cause had ceased to act, and the Body should move in a *Vacuum* without any Resistance at all.

32. THERE are two sorts of Resistance in Fluids; the First arising from the Tenacity of the Fluid, that is, from the Cohesion of its Parts; and that Resistance is always as the Velocity of the Body moving in the Fluid; for the swifter the Body moves in such a Fluid the more Parts it has to remove from their Cohesion in the same Time according as it goes thro' a greater Space. That sort of Resistance may be diminish'd by rendring the Medium more fluid, as Oil, Honey and Pitch, &c. become more fluid by heating. N. B. *Such Fluids as have no Tenacity have none of this Resistance.*

THE other sort of Resistance arises from the Quantity of Matter to be remov'd, and that is always proportionable to the Density

or

or specifick Gravity of the fluid *Medium*. Thus Water resists 850 Lect. V. times more than Air, because a Body moving in it thro' a certain Space, has 850 times more Matter to remove; and if it mov'd in Mercury, the Resistance would be 11900 times greater, because Mercury has 11900 times more Matter than Air in the same Space. In respect to the same Body moving with different Velocities in a Fluid, this Resistance is always as the Square of the Velocity. An Instance or two explain'd by Number will make the Thing evident.

LET us suppose the Body A (*Pl. 25. Fig. 16.**) to be moving* Pl. 25. F. 16. in a *Medium* at the Rate of 2 Inches in a Second or from A to B, and that it is to remove the 4 Particles of Matter b, c, g, f, to make its Way, which Particles we will suppose to be an Inch in Diameter. Now it is not enough to consider that those Particles are to be remov'd, but we must also examine with what Velocity they are remov'd in order to find the Quantity of Motion which they receive. Let us then suppose each of them to be remov'd an Inch in a Second or from the Points f, b, c, g, to f, b, c, g, to make Way for the Body A to go between them. Now since it is the same to move all the 4 Particles laid upon one another from f to f, as to move all the Four, one Inch in different Lines, it is evident that the Space ff or one Inch is their common Velocity: One then multiplied by Four the Number of Particles gives 4 for the *Momentum* of the Matter remov'd by the Body A, which consequently must lose as much of its Motion as it has communicated; and therefore in this Case the Resistance will be 4. Again, let the same Body be suppos'd to move twice as fast, that is, from A to B (*Fig. 16.**) in a Second. There must be 8 Particles (that* Pl. 25. F. 16. is, b, c, d, e, f, g, m, n, or as much more Matter) remov'd in the same Time; but as the Body moves twice as fast, it will strike each of them as hard again, which will make them recede to the Points $\beta, \alpha, \delta, \epsilon, \zeta, \gamma, \nu, \nu$, instead of the Points b, c, d, e, f, g, m, n, in the same Time; so that their common Velocity will be 2 Inches instead of 1. But 8 Particles multiplied by 2 will give 16, which is a *Momentum* 4 times as great as what the Matter of the Fluid received before. Therefore the Body moving twice as fast in the same Fluid communicates four times as much Motion to its Parts, and consequently meets with 4 times as much Resistance. Likewise if the Body mov'd 3 times as fast, it would remove 3 times more Matter in the same Time and also dash it 3 times farther; therefore

X x

it

Leſt. V. it would meet with 9 times more Reſiſtance. And this will hold good in any Degree of Velocity of the moving Body ; for the Quantity of Matter remov'd in a certain Time, and the Velocity with which that Matter is remov'd will always produce a *Momentum* in the fluid *Medium*, and conſequently a Reſiſtance, proportionable to the Square of the Velocity of the Body moving in that Medium. N. B. *This Reſiſtance according to the Square of the Velocity is the only Reſiſtance that the Air is found to have, by Experiments of Bodies moving in it : and therefore it has no Tenacity ; a Conſequence of which is, that its Parts do not touch one another.*

HENCE it is, that a Fluid will reſiſt ſometimes as much as a Solid, nay more, when the Velocity of a Body coming againſt it is very great, as we ſhall ſhew by ſome Inſtances that we ſhall give in the Notes.*

* Ann. 5.

S C H O L I U M.

33. WHAT we have ſaid concerning the Fall of Bodies in the Air, and along inclin'd Planes, will not agree with Experiments, becauſe in the Theory we abſtracted from the Reſiſtance of the Air ; but when we make proper Allowances for it, the Experiments will confirm the Theory. According to the beſt Obſervations a Body falling in *Vacuo* ſhould go thro' 16 English Feet and an Inch and a quarter the firſt Second of its Fall ; but in the Air it muſt loſe of that Space in Proportion to the Motion it gives to the Air, which muſt be ſubtracted from its own Motion ; ſo that the more Matter the Body has in Proportion to its Surface wherewith it ſtrikes the Air, the leſs it will loſe of its own Motion. This will be beſt explain'd by giving an account of an Experiment I made by obſerving the Time of Leaden Balls which I let fall from the inſide of the Top of the Cupola in St. Paul's Church.

E X P E R I M E N T XVI.

I took ſeveral Leaden Balls of 2 Inches Diameter, weighing 2 lb Troy, which I let fall from a Board fix'd 2 Foot over the Top of the inner Cupola, and obſerv'd the Time of their Fall very nicely by an Inſtrument which I ſhall hereafter deſcribe, and found that they fell to the Bottom in $4\frac{1}{2}$ Seconds and a very little more. The Height was 272 Feet. Now according to the Theory, the Balls in that Time ſhould have fallen 52 Feet farther, that is, 324 Feet ; therefore

therefore the Resistance of the Air continually taking off something of what Gravity superadded to the Motion of the descending Body occasion'd the Acceleration not to be so quick, and therefore the Body was longer in falling that 272 Feet than it ought to have been; for since a Body falls a Foot the first quarter of a Second of its Fall, if we take the Square Root * of 272 we shall have $16\frac{1}{2}$ quarters, that is, 4 Seconds and $\frac{1}{8}$ for the Time that a Body would fall 272 * No. 15. Feet in *Vacuo*. Suppose now that the Resistance of the Air took off 5 Inches of the Space which the Balls should have fallen the first Second; during the 2d Second the Resistance of the Air must be greater in Proportion to the Square of the Body's Velocity; that is, as the Body then should go thro' 3 Spaces equal to what it went the first Second, the Resistance must be 9 times greater, consequently the Resistance of the Air would take off 9 times 5 Inches, that is, 45 Inches of the Spaces gone thro' by the Body in the 2d Second: so likewise in the 3d Second the Body must lose 125 Inches; in the 4th Second 245; and in the last half Second above 200; which makes about 52 Foot in all, according to what was observ'd in the Experiment.

N. B. *This Calculation is not exact, being given only as an Illustration rather than a Demonstration, and founded upon a Supposition of a Body falling only 16 Foot in the first Second of its Fall, which Number was taken to avoid Fractions.*

SEVERAL Consequences may be drawn from the Resistance of Air in regard to Bodies moving in it, and which are verified by Experiments. The First is, That *the Motion of a heavy Body is not always accelerated, but at a certain Height it becomes equal and uniform in the Air*; because the Resistance of the Air encreasing in the same Proportion as the Spaces encrease (and consequently in a duplicate Ratio of the Times, or of the Velocities) this Resistance may become so great as to destroy as much of the Velocity as should be produc'd, and by that means hinder the Velocity of the moving Body from being encreased any more; * just as if the Body at that Time should cease to be heavy. The Second is, That *Bodies of different specifick Gravities moving in the same Medium have not their Motions accelerated after the same manner*, by reason of the difference of their Bulk, compared to their Weight, which meets with more or less Resistance; because those of a greater Bulk, when the Weight is the same, drive more Air before them than those of a less. * No. 12, and

Lect. V. *THE Third is, That the Motion of heavy Bodies is differently accelerated in different Mediums, and in the most dense Medium it becomes equal soonest; because the more dense the Medium is, the more Difficulty it has to make its Circulations, and it resists Motion the more easily.*

THE Fourth is, That the least Bodies of the same homogeneous Matter fall with less Velocity, and come soonest to an Equality; because that Body which has a greater Surface is more resisted than that which has a less, and the less Bodies have a greater Surface than the great ones in respect of their Weight or Solidity; for we are taught by Geometry, that if a Cube has its Surface, for Example, of one Foot, another Cube eight times as heavy will have its Surface but of four Foot. According to this Principle, the Dust falls very slowly when it is rais'd, Birds sustain themselves in the Air by spreading their Wings; and a Charge of Shot will not go near so far as a Bullet of the same Weight shot from the same Gun with the same Quantity of Powder, tho' both begin to move with equal Velocities.

THE Fifth is, That there is a determinate Height which produces in a heavy Body the greatest Velocity that it can acquire in falling; so that if it should fall from an higher Place it would have no more Velocity; which is evident from the first Consequence, where we have said that the Motion of a heavy Body is not continually accelerated; but that at a determinate Height it becomes equal.

The Sixth is, That there is a determinate Height, the greatest of all those to which the Velocity which a Body has acquir'd in falling, can make the same Body rise up again; because by the foregoing Consequence, there is a determinate Height, which produces the greatest Velocity that a falling Body can acquire, and that Velocity can make it rise up again but about to the same Height.

THE Seventh is, That a Body thrown upwards by a Force greater than the greatest that it can acquire in falling, ought to be longer in falling than rising; because the Velocity of a Body thrown up to any Height whatever, is continually diminish'd; whereas the Velocity of the same Body in its Fall encreases but till it comes to such a Height, it being certain that if it should encrease continually, the Body would be just as long in falling as in rising.

THE

THE Eighth is, That if a Body be thrown downwards by a Force Lect. V.
that exceeds the greatest Force it can acquire in falling, it will
have a retarded Motion; because by the first Consequence the Body
which falls with the greatest Velocity that its Fall would give it,
meets with a Resistance in the Air, equal to its Gravity; and when
it goes with a greater Force, the Resistance of the Air becomes
greater than its Gravity, and must destroy part of the Motion,
which thus will be slackened and retarded.

THIS last Consequence shews why a Cannon Ball shot down-
wards retards its Motion; because such a Ball is put in Motion by
the Force of the Powder which gives it a greater Velocity, than
that which its absolute Gravity would have given it in falling: And
the Seventh Consequence shews likewise the Reason of this Expe-
riment which Father *Mersennus* takes notice of in his *Balistica*, or
Art of throwing heavy Bodies. *Prop. 13.*

THIS Author says, that he has found by several Experiments,
that an Arrow which has been three Seconds in rising, has been five
in descending; and tho' he adds that an Iron Bullet of three Pound-
weight having been shot upwards perpendicularly by a Mortar-piece
a Foot long, has spent as much Time in rising as in descending, *viz.*
Six Seconds; yet it does not follow that it must always happen so,
the Difference not being so considerable in a Bullet as in an Arrow,
whose Motion comes soonest to an Equality, by reason of its
Lightness.

A common Bomb not receiving from the usual Charge of Pow-
der a Velocity greater than the greatest it can acquire in falling, is
as long in coming down as it is in going up. But a Ball of light
Wood or of Cork (which in a *Vacuum* would go much higher
and farther than a Bomb of the same Bigness, because it receives
from the Powder so much more Velocity at first as it has less Mat-
ter) will not go so high as a Bomb, and also will be longer in co-
ming down than in going up, on account of the Air's Resistance
which has more Effect upon those light Bodies for the Reasons a-
bove given.

N. B. *It is not mathematically true, that a Body falling in
the Air ever comes to an equal Motion; but as it always
approaches nearer and nearer to it, we may take it to be
such physically, and reason from it accordingly.*

Lect. V. 34. I shew'd, after explaining the first Law of Motion, how far it would serve to make us acquainted with the Motion of the heavenly Bodies, by shewing in what Manner Gravity and the projectile Force keep those Bodies in their Orbits; but it requir'd the Understanding of the second Law to conceive rightly how they move in Ellipses that have the central Body in one of the *Foci*, and why their Velocities are successively accelerated and retarded.

35. BUT before I proceed to consider this, I must explain some Astronomical Terms, and shew what is meant by saying, that *the Planets and Comets in respect to the Sun describe Area's about it proportionable to the Times; as likewise the Satellites in respect to their primary Planets.* And this is a Truth known and own'd by all modern Astronomers, however they differ in accounting for the Causes of the celestial Motions.


SUPPOSE a celestial Body to move round another in a Curve returning into it self as a Circle or an Oval; as for Example, the Moon about the Earth, whose Orbit we will consider at first as circular, tho' it is really elliptical. If at any Time of a certain Day we observe the Moon's Place in its Orbit to be at $L \dagger$; and the Day after, at the same Time, the Moon is found to be at another Place as L , the triangular Space $T L L$ (being contain'd by the Line or Ray $T L$ drawn from the central to the revolving Body, at the first Observation, the same Line in the Situation $T L$ at the second Observation, and the Arc $L L$ describ'd by the Moon during the Interval of the Observations) is call'd an *Astronomical Area*, and the Ray $T L$, consider'd as sweeping along that Space and carrying the Moon with it, is called the *Radius vector* or sweeping Ray. If some Days after, for Example 14 Days after, we observe the Moon at l , and the next Day at the same Hour and Minute observe it to be at l , the *Area* $T l l$ will be equal to the former *Area* $T L L$ which was describ'd by the Moon and *Radius vector* in the same Time: and this is what is meant by saying, that *revolving Bodies in the Heavens describe about the central ones Area's proportionable to the Times.*

36. Here we are to observe that the Triangles or *Area's* $T L L$, $T l l$; \dagger are not only equal but similar, and therefore the Body L does in this Case describe the equal Arcs $L L$ and $l l$ in equal Times, as well as equal *Area's*; so that the Motion of L round T is equal, neither accelerated nor retarded. Such are the Motions

ons of Jupiter's *Satellites* about his Center, their Orbits being circular (as far as Observations have been made hitherto) except so much as they disturb each other by their Gravity towards one another, and as they are disturb'd by the Sun according to the different Distances of Jupiter from the Sun, or by Saturn whose Attraction becomes sensible, at, and near, its Conjunction with Jupiter. But all these Inequalities may well be neglected here; because, tho' they are certain consequences of the mutual Attraction of Bodies, they are not considerable enough to be observ'd with Telescopes.

37. Now let us suppose the Body T not to be in the Center of the Orbit, as the Earth is not in the Center of the Moon's Orbit, but to be distant from it the whole length CT (Fig. 19. †) ^{† Pl. 25. Fig. 19.} If the Moon or the revolving Body be observ'd at L and L', and so found to have gone thro' the Arc LL' in ones Day's Time; then again if it be observ'd 14 Days after at l, the next Day it will not be at λ (to which Point it wou'd have gone in a Day if its Velocity had not encreas'd) but it will be got quite to l, its Velocity encreasing so as to make it describe so much a greater Arc as it is nearer to the central Body T; otherwise the *Area* last describ'd wou'd not be equal to the *Area* first describ'd; for as much as Tl, the Distance from the central Body in the beginning of the Description of the last *Area*, is less than TL the Distance from the central Body in the beginning of the first, so much must the Arc describ'd in the last be greater; that what this last *Area* wants in length may be made out in breadth.

28. IF, instead of a Circle, the revolving Body moves in an Ellipse, in one of whose *Foci* the central Body is plac'd (as is the Case of the Moon about the Earth, but more strictly so of the Planets and Comets in their Motion round the Sun) and the whole periodical Time of the Body's Revolution be divided into equal Parts, in every one of those Parts of Time the Body (by its *Radius vector*) will describe an equal *Area*, but none of those *Area's* will be similar except those that are describ'd on each side of the Axis of the Ellipse in correspondent Parts of the Curve at equal Distances from the central Body. Let A B C D E F G H I † represent one of the Ellipses which is describ'd by a Planet ^{† Pl. 25. F. 20.} or a Comet round the Sun; P S is the Axis of the Ellipse and S, s, its *Foci*; S the Sun, and A a Planet at the Sun's *Aphelion* (that

 Lect. V. (that is, at the greatest Distance from it) and P the same Planet at the *Perihelion* (or least Distance from the Sun) and the Time of the Revolution be divided into 10 equal Parts; the Planet setting out at A and going towards B, by its *Radius vector* A S, will describe successively the 10 equal *Area's* A S B, B S C, C S D, D S E, E S P, P S F, F S G, G S H, H S I, I S A, of which only every two correspondent *Area's* are alike, as B S C is like I S H, D S E like G S F, &c.

C O R O L L A R Y I.

HENCE follows, that in an excentrick Orbit, such as an Ellipse, the revolving Body moves faster at the *Perihelion* (the Sun being in one of the *Foci*) than at the *Aphelion*; accelerating its Motion from the *Aphelion* to the *Perihelion*, and retarding it from the *Perihelion* to the *Aphelion*.

C O R O L L A R Y II.

HENCE follows also, that the more excentrick (that is, the longer) the Ellipse is, the greater is the Difference of Velocity at the *Perihelion* and *Aphelion*, such is the Case of Comets, which moving in very excentrick Ellipses, go thro' the lower Part of their Orbits with very great Velocity, but move extremely slow near their *Aphelia*.

C O R O L L A R Y III.

THIS shews why a Planet, tho' it be much more strongly attracted in its *Perihelion* than its *Aphelion*, will not be drawn into the Sun; because it acquires a greater centrifugal Force as its Velocity increases, and thereby balances the Sun's increas'd Attraction. So likewise, when the Planet goes from the *Perihelion* to the *Aphelion*, tho' the Sun's Attraction be decreas'd, because of the increas'd Distance, the Planet will not fly out of its Orbit; for the Velocity decreasing, the centrifugal Force decreases also. In the

† Pl. 25. F. Ellipse represented by Fig. 20 †, when the Planet is at P it is six times nearer the Sun S than when it is at A, therefore it is 36

* L. 1. page 20. times more attracted *; but then its Velocity being also 6 times greater, the centrifugal Force increasing as the Square of the Velocity †, becomes 36 times greater. So that the Attraction, or

† Exp. 15. Page 313.

accelerating

accelerating Force * (however it increafes or decreafes, on account *Left. V.* of the different Distance) is always balanc'd by the centrifugal Force of the Planet.

* *L. 1. page 32.*

C O R O L L A R Y IV.

HENCE we fee, why thofe Planets, which are neareft to the Sun, perform their Revolutions in fhorter Time than thofe which are farther off; that their greater Velocity may give them a fufficient centrifugal Force to balance their centripetal Force (or Gravitation) towards the Sun; Regard alfo being had to their Quantity of Matter †.

† *Exp. 12. p. 312.*

THE Satellites of Jupiter and Saturn alfo have their Periodical Times fhorter, as they are nearer to their Primaries, as we have already hinted †.

† *No. 7. page 308.*

THO' the Orbits of the Planets are nearly circular; yet as the *Foci* of an Ellipfe are confiderably diftant from one another, when the Curvature does not much differ from that of a Circle; the Excentricity will be fenfible enough to be obferv'd. Hence it is, that our Winter Half-Year (in the northern Hemisphere of our Earth) in which we go thro' the *Perihelion*, is 8 Days longer than the Summer Half-Year, &c.

39. *KEP LER* was the firft who discover'd, that the Planets, by a Ray drawn from them to the Sun, describ'd *Areas* proportionable to the Times; and guefs'd that the Caufe might be a Gravitation towards the Sun; but he did not demonftrate it. But Sir *Isaac Newton* has demonftrated it in his *Principia*, fhewing, that when a centripetal Force drives one Body towards another, and that firft Body has received the Impreffion of a projectile Force in any other Direction, it will describe round this laft *Areas* proportional to the Times; and *vice verfa*, that if a revolving Body, by a Ray drawn to a central Body, describes *Areas* proportionable to the Times, it is acted upon by a centripetal Force. Then he fhews (from Observations of the Moon's Motion) that the centripetal Force is the very fame as Gravity, which makes our Bodies fall by an accelerated Motion near the Surface of the Earth. We fhall give his Demonftrations and a further Account in the Notes †.

† *Ann. 6.*

Lect. V. 40. IN the mean time I shall endeavour, in the easiest way, to shew how Gravity makes the Planets describe their excentrick Orbits with a Motion uniformly accelerated and retarded; for tho' I shall give no strict Geometrical Account of every thing relating to this Motion, and the Nature of the Curve, yet every Assertion will be a Consequence of the two Laws of Motion, already explained, and their Corollaries.

Plate 25. Fig. 20.

Pl. 25. F. 20. THE Ellipse represented by this Figure, is more excentrick than any of those that are described by the Planets, but not so excentrick as those that are described by the Comets. I took it at a Mean; because, as both the Comets and Planets are retain'd in their Orbits, and continue their Motions from the same Causes, one Explication might serve both.

LET S represent the Sun, A the Planet or Comet, which Gravity (or the mutual Attraction of the Sun and revolving Body) drives towards the Sun in the Direction AS; and let AM represent the Quantity of that Force; that is, the Space, which that Force alone acting, would cause the Planet to go thro' in a given Time. Let the Planet A be supposed to have receiv'd a projectile Force in the Direction $A\alpha$ at right Angles to AS. If Aa expresses the Space which the projectile Force alone would make the Planet describe in the said given Time, and the Quantity of that Force be such, as acting jointly with Gravity, will make the Planet (setting out in the Diagonal Am of the compleated Parallelogram $Aa m M$) describe the Circle $A m \nu$, &c. whose Center is S the Center of the Sun; then a greater *Vis impressa*, or projectile Force, such as $A\alpha$ (the Force of Gravity AM remaining the same) will make the Planet set out in the Diagonal An of the compleated Parallelogram $A\alpha n M$, and describe an Ellipse $An \nu$, &c. whose nearest *Focus* is S; then will the Point A become the *Perihelion* instead of the *Aphelion*, and the Motion will be retarded from A till it comes to the *Aphelion* on the other Side of S; and then from that *Aphelion* accelerated till it comes back to A, from whence it began its Motion. But if the projectile Force be less, than what we have supposed capable of making the Planet move in a Circle, and it be expressed by Aa instead of $A\alpha$, then the Planet

net (beginning its Motion in the Diagonal A B of the compleated Lect. V. Parallelogram A a B M) will describe the Ellipse A B C D E P F G H I A, the Sun S being in the farther *Focus*, and the nearest *Focus* will be at s; so that now A will be the *Aphelion*, and P the *Perihelion*. Pl. 25. F. 20.


Now to shew how the Planet is accelerated in going from the *Aphelion* to the *Perihelion*; let us observe that at first setting out, the Direction of the projectile Force A α is at right Angles with the Direction of Gravity (or the centripetal Force) A S. When the Planet by the joint Action of the two Forces is come to B; the projectile Force has its new Direction along the Tangent B b , which makes an acute Angle with the new Direction of Gravity which now is B S; therefore the Planet will describe a longer Diagonal in the same Time,* *viz.* B C, that is, encrease its Velocity so that the Area B C S may by a Breadth proportionably greater be equal to the longer Area A B S. When the Planet is come to C, the Direction of the projectile Force along the Tangent C c still making an acute Angle with C S the Direction of Gravity, the two Forces conspiring still accelerate the Planet and carry it from C to D in the same Time that it went at first from A to B. The same Forces conspiring still in their Directions D d and D S, when the Planet is at D, will in a Space of Time equal to the former carry it to E: and lastly, the same Forces with their conspiring Directions will still accelerate the Planet in its Motion from E to its *Perihelion* P, where its Velocity is the greatest of all. At P the *Perihelion*, the Direction of the projectile Force is along the Tangent P p , and makes a right Angle with P S the Direction of Gravity; and the Planet from the Action of those two Forces will go to F. When the Planet is at F, the Direction of the projectile Force along the Tangent F f makes an obtuse Angle with F S the Direction of Gravity, and therefore the Motion of the Planet must be retarded,* because the Diagonal F G will be shorten'd, as the Angle f F G opens, and the Forces begin to act against one another. The Angle above-mention'd will still encrease at G, so that the Force of Gravity in the Direction G S will check the projectile Force which now acts in the Line G g , and still retard the Planet more. The Angle continuing to be obtuse at H and I, the Planet is still retarded till it comes to A the *Aphelion*, where its Motion is the slowest of all. *It may perhaps be objected here; that since the Angles a A S, b B S, c C S, &c. only decrease half way from A to P;** and the Angles S P p , * No. 13.

Lect. V. S F f, S b g, &c. only encrease half way from P to A; * the Planet descending towards the Perihelion should not encrease its Velocity after it is come half way, because then the Angles, as S E e, &c. encrease again: neither should it retard its Motion after it has moved half way from P to A, because the Angles made by the Directions of the two Forces no longer encrease but gradually decrease. But then we must consider that there is another Cause of Acceleration and Retardation which does not depend upon the Quantity of the Angle above-mention'd, but continues to encrease after the Angles cease to diminish in the Descent of the Planet towards the Sun; and that Cause likewise continues to decrease in the Ascent of the Planet from the Perihelion to the Aphelion, even after the Angles of the Directions of the Forces cease to encrease: and that Cause is the diminish'd or encreas'd Distance of the Central Body (or the Sun S) the Power of Attraction changing continually in a reciprocal Proportion of the Square of that Distance. If, for Example, we would compare the Velocity of the Planet at D and at E, we must compleat the Parallelogram D δ E d, of which the two Sides D δ and D d represent the Force of Gravity and the projectile Force when the Planet is at D; then we must also compleat the Parallelogram E ϵ P e, in which E ϵ is greater than D δ in the same Proportion as Gravity is greater at E than D; likewise E e must be greater than D d, in Proportion as the projectile Force by its continual Acceleration is there also become greater; and tho' the Angle ϵ E e is greater than δ D d, we shall have a greater Diagonal EP on account of the greater Length of the Sides in the Parallelogram ϵ E e P.

41. To shew how agreeable the Gravitation of Planets and Comets (for what we have said of Planets is as applicable to Comets) towards the Sun, is to the Gravitation of heavy Bodies towards the Earth; let us take a View of the 21st Figure † representing a long Ellipse, or the Orbit of a Comet about the Sun S, when π represents the Perihelion. The Semi-Ellipse $\pi \alpha$ is the Line in which the Motion of the Comet is uniformly retarded, and the Semi-Ellipse $\alpha \pi$ the Line in which the Motion of the Comet is uniformly accelerated. Now, if instead of including the Sun S in the Orbit, we should from one Part of the Surface of the Sun P, project a Body upwards, so that it should rise as far as A, its Motion would be uniformly retarded till it came to A, where it would have its least Velocity; then it would turn again and fall in the Line

Pl. 25. F. 20
*Ann. 7.

† Pl. 25. F. 21.

Line *Ap* accelerating its Motion till it came to *p*, taking up just Lect. V.
as much Time in its Ascent as Descent. And this is what we have 
shewn concerning the Action of Gravity upon Projectiles, abstract-
ing from the Resistance of the Air.

42. BUT now let us take in the Resistance of the Air, and compare it with any resisting Medium, to see what would happen to the Planets, if they mov'd in such a Medium.

WE have shewn, [†] that when Bodies move in the Air, they lose of No. 32.
their Motion by the Resistance of the Air, in Proportion to the Page 340 and
Square of their Velocity; and that that Resistance hinders falling 341.
Bodies from accelerating their Motion, as they would do, if they
fell in a *Vacuum*; because that Resistance continually taking away
some Part of the Velocity, which Gravity superadds to the falling
Body, continually brings the Motion of the Body nearer and near-
er to a Motion of Equality. Now, if the Planets mov'd in a re-
sisting *Medium*, such as the cœlestial Matter, which the *Cartesians*
suppose; the Resistance of that Matter would hinder a Planet from
acquiring that Velocity in its Descent to the *Perihelion*, which is
necessary to make its centripetal Force balance the Force of Gra-
vity; for this last Force would always increase in Proportion to the
Square of the decreas'd Distance, let the *Medium* encompassing the
Sun be of what Nature it would; but the centrifugal Force would
want its proper Increase, if the Planet wanted its requir'd Velocity.
The Consequence therefore would be, that the Planet would change
its Track and come nearer to the Sun, and revolve in a longer El-
lipse. The next Revolution, the Planet coming towards the *Peri-*
helion, and wanting its due centrifugal Force, would be brought
nearer to the Sun by the Attraction (not diminish'd, but increas'd,
because now the *Perihelion* would be nearer); then again would
the Ellipse be changed into a longer, and the next *Perihelion*
would be still nearer; till the Planet for want of its due Velocity in
a Direction along the Tangent, approaching nearer and nearer
every Revolution, would at last fall into the Sun.

NOW, since no such Thing happens, it is evident, there is no such
resisting *Medium*, or cœlestial Matter of a *Vortex*, as the *Cartesi-*
ans suppose to be the Cause of the Motion of the Planets round
the Sun. So far from that, that such a Fluid would destroy the
Motion of the Planets, as we have shewn.

BUT

Lect. V.

BUT what destroys the *Cartesian* Hypothesis at once (as well as the Opinion of those ancient Philosophers, who supposed solid Orbs of crystal to belong to every Planet, and carry it round) is the Observation of Comets, which are neither Meteors, as some of the Ancients suppos'd them, nor Planets straying from one *Vortex* to another, as the *Cartesians* assert; but Planets moving in very excentrick Orbits, which we shall consider more fully in another Place.* Only here we are to observe, that they move freely to and from all Parts of the Heavens; and therefore there can be no crystal spherical Shells, which would stop them; nor Whirlpools of Matter, which would change their Direction by degrees, and at last make them move nearly in the same Plane as the Planets, whose Orbits have all their Planes contain'd in the Breadth of a Zone of a few Degrees. But what is most contrary to the *Cartesian* Hypothesis, is the Motion of a retrograde Comet, such as that of the Year 1682, which moving from East to West, was carried directly against the suppos'd Stream of celestial Matter; and, instead of having its Motion first diminish'd, and then quite stopp'd, and afterwards being carried in a contrary Direction (which must follow, when a Body moving from East to West falls into a Vortex, whose Matter moves from West to East) it accelerated its Motion in its Descent towards the Sun.

* Ann. 8.

43. THO' we may very well call the *Medium*, in which Planets move, a *Vacuum*; yet, since Light is propagated thro' all the celestial Spaces, and some fine *Effluvia* may be separated from the Comets and Planets, there will (strictly speaking) be some Resistance to the Motions of the Planet, tho' not so much by many thousand times as our Air would make; and that Resistance after a great Number of Years, must so alter the Motion of the Planets, as to require the Author of Nature's mending Hand.† If any Alteration has been found in their Orbits, tho' ever so small, since Astronomers began to make accurate Observations (as a great many assert that there has) that will be sufficient to shew, that the World is not *eternal*, if there were no other Arguments against its Eternity.

† This is Sir Isaac Newton's Opinion. See the Queries at the End of his *Opticks*.

THE Sun has been observ'd to have a considerable Atmosphere; as its Surface, on account of the prodigious Heat, must always be throwing out *Effluvia*, those *Effluvia* (except such as are small enough to become Particles of Light, and be darted off with immense

menſe Velocity) floating about round the Sun's Body muſt make a *Leſſ. V. Medium*, at leaſt as denſe as our Air. Now, if a Comet comes near enough to go into the Sun's Atmosphere, it will on the account of the Reſiſtance it meets with, come nearer and nearer to the Sun every Revolution, and at laſt fall into it. Such may in Time be the Fate of the Comet obſerv'd in 1680, which came ſo near the Sun as to be, at its *Perihelion*, no farther diſtant from the Sun's Surface than the 6th Part of the Sun's Diameter. For what we know, many a Comet may have fallen into the Sun without our Knowledge and Obſervation; and perhaps thoſe Bodies may ſerve as freſh Fuel to replenish the Waſte of the Sun in ſupplying the System with Light; for tho' it has been objected, that a Comet would be but a ſmall Supply, yet if it be as big as the Earth, it will be in Diameter the 100th Part of that of the Sun; that is, in ſolidity the Ten hundred thouſandth Part; and that may be as much as the Sun in many Years may loſe in Light; nay, tho' the Comet ſhould be no bigger than the Moon (as moſt Comets are ſuppoſed to be of that Bigneſs) yet it might ſtill be a ſufficient Supply for the Waſte of Light.

44. THO' theſe are but Conjectures, yet it may not be unacceptable to the Reader, to ſhew by a Scheme, how a Comet, when once it comes in to the Sun's Atmosphere, will at laſt fall into it.

Plate 26. Fig. 1.

LET $ABP\pi$ be the Orbit of a Comet, S and F its *Foci*, S the Sun, and $D\beta C$ the Sun's Atmosphere. When from the *Aphelion* A the Comet is come towards the *Perihelion* as far as B , the Reſiſtance of the Sun's Atmosphere hindring ſome of the Acceleration, which the Comet ought to have, the Sun's Attraction will give its Orbit more Curvature at the *Perihelion*, bring it nearer to it ſelf, and make it come to b inſtead of π in its going off, ſo that it will then have leſs Curvature, the Sun's Attraction, at its going off, acting more directly againſt the Direction of the projectile Force. This will make the Ellipſe longer, carry the *Aphelion* to A , and make the *Focus* at f be farther off from the Sun than when it was at F . The next Revolution, when the Comet comes down to β , it will ſtill come nearer to the Sun in its *Perihelion*, and quit it at B in a new Direction, ſo as to go off in an Ellipſe ſtill longer, whoſe farther *Focus* is at ϕ and *Aphelion* at α ; and ſo on, till

Lect. V. till at last it comes down to the Sun in the Line α S. But if a Comet or Planet moves in the Orbit $a p$, whose *Foci* are at the Sun S and at f , and whose *Perihelion* p is quite out of the Sun's Atmosphere, the Motion of the revolving Body will not be sensibly disturb'd in many thousand Years.

OTHER Things relating to Astronomy cannot be well understood, till we have explain'd

The Third LAW of Motion.

45. *To every Action there is always oppos'd an equal Reaction; or the mutual Actions of two Bodies upon each other, are always equal, and directed to contrary Parts.*

WHATEVER draws or presses another, is as much drawn or pressed by that other. If a Man presses a Stone with his Finger, the Finger is also press'd by the Stone. If an Horse draws a Stone tied to a Rope, the Horse (if I may so say) will be equally drawn back towards the Stone: For the stretched Rope, by the same Endeavour to relax and unbend it self, will draw the Horse as much towards the Stone, as it does the Stone towards the Horse, and will obstruct the Progress of the one as much as it advances that of the other. Suppose, for Example, that the Horse is able to overcome an Obstacle equal to 1000 lb Weight, pressing against it with his Breast; when the Horse draws a Stone of 100 lb Weight, he will then be able to surmount an Obstacle but of 900 lb the Stone taking away from the Force of the Horse as much as serves to bring it self forward. We must therefore take care rightly to understand the Term *as much*, and distinguish it from *as far*. If a Body strikes upon another, and by its Force changes the Motion of the other; that Body also (because of the Equality of the mutual Pressure) will undergo an equal Change, in its own Motion towards the contrary Part. The Changes made by these Actions are equal, not in the Velocities (unless in such Cases as the two Bodies have the same Quantity of Matter) but in the Motions or *Momenta* of the Bodies; that is to say, if the Bodies are not hinder'd by any other Impediments. For, because the Motions are equally chang'd, the Changes of Velocities made towards contrary Parts, are reciprocally proportional to the Bodies. *This Law takes place also in Attractions.*

Plate



Plate 26. Fig. 2.

46. IF in a large Vessel of Water AB a Loadstone L *l* be set afloat on a Piece of Cork, and a Piece of Iron or Steel I *i* of the same Weight be likewise set afloat on another Piece of Cork, they will come towards one another and meet at C the middle of their Distance; which shews, that the Attraction is mutual between the Loadstone and the Iron.

IT is well known, that any long Piece of Steel, which has had each End drawn over the Poles of a Loadstone, will with one End attract the Pole that gave it the Virtue, but repell the other Pole; and so likewise with the other End. If then I *i* be the touch'd Piece of Steel, whose End I has receiv'd its Virtue from the Pole L of the Stone, and *i* from the Pole *l*; the Steel and Stone will come together in the manner above-mentioned, when I is plac'd towards L; but if either *i* of the Steel, be placed towards L of the Stone, or *l* of the Stone towards I of the Steel, and they be brought as near to C as the Corks on which they float will allow, then as soon as they are left to themselves, the Stone and Steel will repell one another; which shews, that Action and Reaction, or equal and contrary as well in the Repulsions as Attractions. †

† Ann. 9.

47. A G and B* are two Boats of equal Bigness and Weight,* Pl. 26. F. 5. floating on the Water and at rest, at the Distance G F; a Man in one of them at G pulling a Rope fasten'd at F, by pulling will bring both the Boats together, and they will meet at C their common Center of Gravity, which happens also here (because they are equal) to be the middle of their Distance †. When the Boats † L. 2. are together at C, if the Man pushes the Boat F B from him, that Boat and his own Boat will recede from each other to equal Distances from C. But if the Boat F B had been as big again (for Example had been F B H) and the common Center of Gravity of the two Boats had been at *c*, the Distance of the Boats being G F; then by the Pull of the Man, the Boats would have met at *c*, the biggest Boat going thro' but half the Space which the least would go thro'; likewise if they were push'd asunder from the Point *c*, the Velocity of their Recess would be reciprocally proportionable to their Mass; that is, A G would recede as far again as F B H, the common
Z z Center

Lect. V. Center of Gravity in both Cases being at rest.† It is plain here, that the *Momenta* of both Boats will always be equal, tho' their Velocities are only equal when the Boats are so; and these equal *Momenta*, whereby the Boats are carried towards contrary Parts, shew that Action and Re-action, in all Cases where Bodies act upon one another are equal and contrary. If the Boat A G had been close to a large Ship, and the Man had push'd his Boat off, he would have given as much Motion to the Ship as to the Boat: I say, the Ship would have mov'd *as much*, but not *as far*; because the Velocity of the Ship being as much less as the Ship is bigger than the Boat, would have been insensible to Sight; and therefore the Vulgar in such a Case imagine, that the Ship does not move at all; and much less, that when a Man pushes against the Shore to shove off his Boat, he moves the whole Earth as much as his Boat is mov'd; tho' it is certainly true. In firing a Cannon, the Explosion of the Powder pushes the Cannon backward *as much* as it pushes the Ball forward, only the Quantity of Matter being vastly more in the Cannon than in the Ball, the Recoil is but of a few Feet, whilst the Ball goes perhaps 10000 Feet; in considering the Recoil we must add the Friction of the Carriage of the Cannon against the Earth, which will still diminish the Velocity of the Cannon's Motion. If the Cannon be fasten'd to a Ship, we feel only a hock as we stand in the Ship when the Cannon is fir'd, because all the Matter of the Ship being added to the recoiling Cannon, the Velocity diminishes in Proportion to the Matter added to the Cannon, which makes it insensible to the Sight, and only to be felt by a Shock.

48. ACTION and Re-action are very plainly seen in rowing, swimming and flying; as for Example, when the Man K in the Boat I K (Fig. 4.*) pulls his Oar, he drives the Water towards H, and the Water drives the Boat as much towards D. In swimming, which is nothing but rowing with the Hands and Feet, we are as much push'd forward by the Water as we push the Water back. The same Thing explains the flying of Birds, who are push'd forward by the Re-action of the Air against their expanded Wings, when they strike the Air with them. As for Example, if a Bird strikes the Air downwards with his Wings, with a Force equal to what would raise 10 Pound, the Re-action of the Air will push him up with the same Force; but if the Bird weighs one Pound, the Effect of the Re-action of the Air will make the Bird rise with Force of only 9 Pound; that is, the Bird will rise just as one Pound would

*Pl. 26. F. 4.

would do tied to a String running over a Pulley, by the Force of Leſt. V. the Deſcent of 10 Pound at the other End of the String. If the Bird ſhould ſtrike the Air only with a Force equal to his own Weight, he would for ſome time be ſuſpended in the Air without Motion, as we often ſee Kites, Hawks, and other Birds of Prey.

49. THE above-mentioned Laws of Motion once underſtood, the *Phænomena* of the Tides will be eaſily accounted for; but to make the Matter ſtill eaſier, let us take the following Conſideration or *Lemma* along with us.

If, when three Bodies are moving after one another the ſame Way, with the ſame Velocity, there be an additional Force impreſs'd upon each of them, but greater in the Firſt, leſs in the Second, and yet leſs in the Third; their Diſtances from each other will continually increaſe, tho' they all continue to move the ſame way, and all of them faſter than they did before.

Let us ſuppoſe the three Boats AG, FB, and IK† (Fig. 3, and 4.) to be carried along a Stream from L towards D, floating down with equal Velocity, and that there is but one Man to row the Boat AG, two Men to row the Boat FB, and four Men to row the Boat IK. Now whiſt none of the Men row, the Boats, as they are carried down by the Stream, continue to be at equal Diſtances from one another; ſo that if a Man, ſitting in the middlemoſt Boat at B, can with a long Rod reach the Head of the hindmoſt Boat at G, and the Stern of the foremoſt Boat at I, he ſhall continue to be able to do it while the Men in the Boats do not row; but, if we ſuppoſe, that all the Men at once begin to row, the Motion of all the Boats towards D will be accelerated, but differently, for the four Men in the foremoſt Boat will accelerate it faſter towards D than the two Men in the middlemoſt Boat, and theſe two laſt will make their Boat go on faſter than the ſingle Man in the laſt Boat AG; ſo that the Perſon, who holds a Rod in the middle Boat FB, will no longer be able to reach either the Boat that goes before him, or the Boat that follows him; but will be apt to fancy (when he does not attend to his own Motion) that the Boat before him haſtens away from him, and that the Boat behind him goes backward. This Conſideration will help us to explain the Cauſe of the Tides.

Lect. V. 50. If the Earth was perfectly smooth, without Mountains or Vallies, the Sea would make a watery Shell about it, which Shell would be concentrick to the Earth, if no Body was near it, to alter the Figure of this Fluid by its Attraction. Let $\alpha p \gamma n$ * represent the Figure of the Earth in such a Supposition, C its Center, and A P L N the Surface of the Sea, which is concentrick to the Earth, because equally gravitating towards the Center of the Earth in every Part. Now, let us consider, what Effect the Moon at M (C M being a Distance of 60 Semi-diameters of the Earth) must have. Since Action and Re-action are, equal, as much as the Moon gravitates towards the Sea at L, so much does the Sea gravitate towards the Moon; but as the Sea does also gravitate towards C the Center of the Earth, with much more Force (as it has 40 times more Matter, and is 60 times nearer) the Moon at the Distance M L attracting it in a contrary Direction can only take off from the Gravitation towards the Earth so much as amounts to its accelerating Force at that Distance M L. This will make the Water at L swell up to l ; and at the same Time it will be High-water at A, the Water also swelling up to a on the opposite Side of the Earth, whilst the Water falls at P and N to supply the Rise at l and a . If we consider the Water at L, the Earth at C (reducing all its Weight into its Center of Gravity, as it is not a Fluid to change its Shape) and at the *Antipodes* of L the Water at A, we shall come to the Case of the three Bodies, or three Boats in the preceding *Lemma*; for all these three gravitate towards the Moon at M; but differently according to their Distance, in the following Proportions. The Water at L is distant from the Moon M, 59 Semi-diameters of the Earth; but the Center of the Earth C is distant from the Moon 60 Semi-diameters; therefore as much as 3600, the Square of the Moon's Distance from the Center of the Earth, is a greater Number than 3481 the Square of 59, or the Moon's Distance from the Sea at L; so much is the Attraction of the Moon (that is, the accelerating Force towards the Moon) greater on the Sea at L, than the Earth at C, which makes it come forwarder towards the Moon to l ; or in other Words, this makes High-water at l under the Moon. There is likewise at the same Time High-water at the *Antipodes*, or opposite Part of the Earth at a ; because the Water there being less attracted than the Center of the Earth (in the reciprocal Proportion of the Square of the Distances, that is as much as 3600, the Square of the Distance of the Center of the Earth, is a less Number than 3721 the Square of A M = 61 the Distance

Distance of the Sea at the *Antipodes* from the Moon) must rise at *Lect. V.*
a by being left behind, or not advancing towards the Moon so fast
as the Center of the Earth. For as the three Bodies *L*, *C*, and
A, all tend towards the Moon, but *L* with more Force than *C*,
and *C* with more Force than *A*, the Distance *C L*, as well as the
Distance *C A*, must increase from those Inequalities of Force act-
ing the same Way.

51. I HAVE often heard it objected, that it did not seem proba-
ble, that the Moon should raise the Water at one Part of the Earth,
as *L*, by attracting the Water more than the Earth; and at the
same time raise it at the opposite side of the Earth, as at *A*, by at-
tracting it less than the Earth; but the whole Objection will va-
nish, by explaining the meaning of the Word *to raise* the Water,
which here is equivocal. In respect of the Earth, what is farther
removed from the Center of the Earth, is said to be *rais'd*, and in
that Sense the Water at *a* is raised as well as at *l*; but in respect
of the Moon at *M*, if the Water at *L* is said to be *rais'd*, because
it comes to *l* nearer to the Moon, the Water at *A* going to *a*,
farther from the Moon, should rather be said to be *depress'd*, or
left behind, as it is less attracted than the Earth. If we consider
the Earth drawn towards *M*, so that the Part *B* \propto *B* of its Sur-
face, is brought to *B* \propto *b*, whilst the Water remains at *A*, or comes
on towards *M* more slowly than the Surface at *B* \propto *b*, there will
happen the same Thing to an Inhabitant at *a*, as if the Water
(without any Regard to the Moon) had risen at *A* from *A* to *a*.
To make this still plainer; let us suppose an Inhabitant of the Earth
at *A* to stand in the Sea near the Shore, so as to have the Water up
to his Middle, his Feet being towards *a*, and his Head towards *Z*
the *Zenith*; if the Moon be in the *Nadir* at *N* (that is, under the
Earth in respect to the Man at *A*) the Earth on which the Man
stands will be drawn downwards whilst the Surface of the Water
does not descend so fast; the Consequence of which will be that
the Man will be up to the Neck in Water whereas he was only
up to the Middle before; and if he knows nothing of the Moon's
Action in drawing down the Bottom on which he stands, he will
say that it is the Water that rises—and this is really what is cal-
led the coming in of the Tide when we have the Moon at our *An-*
tipodes.

Lect. V. 52. FOR the sake of such Readers as may not be so well satisfied with what has been said of the Tides hitherto; I'll give an hydrostatical Solution of it, anticipating only one plain hydrostatical Proposition, namely, that *all Liquors of the same specifick Weight,** which being contained in different Vessels have a Communication, stand at the same Height; but if in any of the Vessels there be a Liquor specifically lighter than the rest, it will rise so much higher as it is specifically lighter to restore the Æquilibrium.* Now we must observe, that the Points L N A P being equally distant from the Center of the Earth C are equally high, and the Waters at those Places, which we suppose to communicate with one another do gravitate towards the Center of the Earth in the Lines L C, N C, A C and P C in the equal Columns L γ , N n , A α , and P p ; whilst no other Body but the Earth attracts the Waters. But when the Moon is at M, the Columns of Water at L γ become less heavy towards C than they were, because the Moon attracts them in a contrary Direction (*viz.* from L to M) which will have the same Effect as if the Water there was specifically lighter, as it would be if it lost its Saltness. But the Columns at P and N do not become less heavy towards C; because the Moon drawing at right Angles to P C and N C does not diminish the Tendency of the Waters at P and N towards C; and therefore they retain their full Gravity, which makes them overweigh the Columns at L which have lost some of their Gravity towards C: the Waters therefore will fall at P and N and rise at L, till the Waters from L coming up from l have as much added to them in Quantity as they have lost in specifick Weight; and thereby the Æquilibrium will be restor'd. Likewise the Columns of Water at A, which have less Tendency towards C, than the Waters at P and N, do (as it were) become specifically lighter; and therefore, in order to maintain the Æquilibrium, they must receive an additional Quantity of Water from P and N, which raises them up to a .

53. I am aware of an Objection here, which seems at first to be of great Force; namely, That since the Waters at L, by the Moon's Attraction, have a Tendency towards the Moon in the Direction L M, the Waters at A must also receive a Tendency towards the Moon in the Direction A M by the Moon's Action, tho' that Action is less at A than at L; and consequently that instead of their Gravity

** One Body is said to be specifically heavier than another, when it contains more Matter under the same Bulk, or as much Matter under a less Bulk.

or Tendency towards C being diminished, it ought to be encreased, be the Addition ever so small; and therefore they ought to press more towards C than the Waters at P and N, and so be lower at A, instead of rising up to *a*. Lect. V.

IN answer to this, we are to consider that as the Earth at C is more attracted towards the Moon than the Water at A, it is the same Thing as removing the Center of Gravity of the Earth from C to *c*, and by that Means the Water at A will be so much less attracted towards C on account of the greater Distance A *c* * instead of A C, that even with the additional Force given by the Moon¹⁷ in the same Direction it will have less Tendency towards the Center of the Earth than the Water at P and N, and consequently it must be rais'd up to *a* by the Flow of Water from P and N. L. 1. N°.

54. THERE is yet another Objection which seems to destroy the whole Foundation of this reasoning upon the Tides, and to make the Case quite different from the Example of the 3 Boats which we have given to illustrate it. And it is this, *viz.* If the Earth C and the Waters at L and A (as in the Case of the 3 Boats) were all carried towards the Moon, the Earth and Moon must in Time come together; but we cannot observe the least Approach towards it, the Distances of the Earth and Moon varying only according to the Nature of the Curve in which the Moon moves, and the joint Actions of the Sun and Earth according to the different Position of those 3 Bodies, and the different Distance of the Earth from the Sun as it goes round it in its elliptical Orbit.

IN Answer to this, we must observe that it is an undeniable Consequence of the mutual Attraction of all Bodies, and the 3^d Law, that the Earth must gravitate towards the Moon as well as the Moon towards the Earth; but we are not to expect to see the Earth go nearer to the Moon, because another Cause destroys that Effect. For the Moon and Earth (as we have before observed *) * L. 2. N°. describing similar Ellipses round their common Center of Gravity,³² by the Velocity of their Motion acquire such a centrifugal Force as balances (or takes off the Effect of) their centripetal Force towards the common Center of Gravity. Hence only a Tendency is left, and not a visible Effect in the Earth it self. But in respect of the Waters, that Tendency must produce a visible Effect, tho' they do not leave the Earth; because there is a Communication from

Lect. V. from the Water at P and N to the Water at L and A. And as we have already demonstrated that the Waters at L and A have a less Tendency towards C the Center of the Earth than the Waters at P and N; the Effect of that Tendency will be, that the Waters must rise at A and L and fall at P and N to maintain the *Æquilibrium*. But if there was no Communication between P, N, and A, L, there would be no visible Effect; that is, the Water would not rise at L and A, as it really does not in Lakes, and such Seas as have no Communication with the Ocean, unless they have a vast Extent.

55. THE next Thing we have to shew relating to the Tides, is the Rise and Fall of the Water successively in any one Place; for Example, if it be high Water at *Portsmouth* at Noon, it will be low Water a little after Six in the Evening; then about Midnight, or Six Hours after, high Water again; at Six in the Morning low Water; and, about Noon the next Day, high Water again, &c. From what has been said it appears, that the Waters about the Earth do by the Moon's Attraction form themselves into an oblong Spheroid, whose longer Axis produced goes thro' the Center of the Moon; as the Spheroid $L N A P^*$, whose Axis is $a l$, and which produced goes thro' the Center of the Moon at M. Now let us suppose L to be any Sea-Port on the Ocean, as suppose at *Lucaio* one of the *Bahama Islands* in the Latitude of 27 North, and that C is the Axis as well as Center of the Earth; and that *Lucaio* is at Noon at L where the Water is under the Moon M. As the Earth turns from West to East, in Six Hours, *Lucaio* will be carried to P where the Water is low (for the Spheroid of Water will be immoveable in respect of the Moon, which we suppose at rest at M, whilst all the Countries of the Earth go round in the Circle L P A N or other Circles parallel to it): in six Hours more *Lucaio* will come to A where it will have high Water; six Hours after that, it will be at N, where the Water is low; and lastly, it will return to L the next Day at Noon, and so have high Water.

* Pl. 26. F. 5.

THE real Fact is, that if it be high Water in any Place this Day at Noon, it will not be high Water next Day till about 53 Minutes, that is, near an Hour after Noon to-morrow. The Cause of which is, that the Moon does not remain at rest in the Heavens as we have just now supposed, but moves in its Orbit 13 Degrees and $10\frac{1}{2}$ Minutes, as for Example thro' the Arc M *m* in 24 Hours (Fig. 6)

(* Fig. 6) so that when any Place of the Earth, as *Lucajo*, has gone from *L* to *L*, then to *l*, and to *γ*, and lastly to the same Place at the same Hour at *L*; the Moon being no longer at *M* but at *m*, the high Water has changed with the Moon's Place, and the long Axis of the Spheroid is no longer *L l* but *N n*; and therefore *Lucajo* must move quite to *N* where the high Water is towards the Moon, the Spheroid of Water being now represented by the prick'd Oval, whose Axis produc'd is the Line *n N m*; and the Time of that Motion will be about 53 Minutes, at the Rate of 15 Degrees in an Hour, which every Country of the Earth moves from West to East.

Lect. V.
* Pl. 26. F. 6.

56. BESIDES this alternate Rise and Fall of the Water twice in 24 Hours, it is observ'd that at the Time of full and new Moon, the Tides (then called Spring-Tides) are greater than at the Time of the Quadratures, that is, when to us the Moon appears halv'd; and as the Water rises higher at the Tide of Flood, so it sinks lower at the Tide of Ebb, when the Moon is Full and when it is New. Those Tides that happen at the Quadratures, are called *Neap-Tides*. And both are accounted for by the joint Action of the Sun and Moon.

IF the Action of the Sun on the Waters, and the Reaction of the Waters be consider'd, as we have done in respect of the Moon, we shall find (for the same Reasons) that the Waters will rise under the Sun and at the *Antipodes*; but as the Attraction of the Sun, on account of its immense Distance, is above five times less than that of the Moon, ** all the Effects, *ceteris paribus*, will be above five times less. Now, since both the Moon and Sun act upon the Waters, when their Actions co-incide (as they do at *Full* and *New Moon*) the Water rises one fifth higher, and falls one fifth lower; and when their Actions are contrary to each other (as they are at the Quadratures) the Tides rise and fall one fifth less. Sir *Isaac Newton* has shewn * that the Moon is capable of raising the Water of the Ocean 10 Foot, and the Sun only 2 Foot; therefore when both their Actions joyn, the Water will rise 12 Foot;

* Princip. L.
3. Prop. 38.

** The visible Surface of Bodies decreases as the Squares of their Distances encrease, and the Power of the Attraction of Bodies (that is, their accelerating Force) decreases in the same Ratio; * therefore, if the Sun and Moon were of the same Density, their Attractions on the Earth and Seas would be equal when their apparent Diameters are equal; but the Sun having about 5 times less Matter than a Moon of the same Bulk would have (that is, being about 5 times less dense than the Moon) has 5 times less absolute Force, and therefore must have 5 times less Power to move the Sea.

* L. 1. Ann.
11.

Lect. V. and where they act against each other, the Water will rise only 8 Foot. But this will be better understood by a Figure.

Plate 26. Fig. 7.

Pl. 26. F. 7. A C L is the Earth, whose Center is at C ; M or *m* the Moon, whose Action reduces the Water to the oblong Spheroid *l n a p* whose Axis produced goes thro' the Moon. S is the Sun, which, if there had been no Moon, would have raised the Water up to *A* and on the opposite Side to *L*, so as to reduce it into the Spheroid *A P L N* less oblong than that of the Moon. Now, when the Sun and Moon are in the same Line, (as when the Moon is full at *M*, or new at *m*) the Axes of the two Spheroids co-incide so that the Water, which would have been raised by the Moon but to *a* and *l*, by the Sun's Additional Force will be raised up to *α* and *λ*, 12 Foot instead of 10 Foot, and consequently depressed so much lower at *p* and *n*. But if the Moon being at *M* or *m*, the Sun was at *s* or *σ*, (that is, at the Quadratures, when the Moon appears halv'd) the Sun's Action would be capable of raising the Water up to *P* and *N*, if it acted alone ; but as the Moon at the same time raises the Water at *a* and *l*, all the Effect of the Sun's Force will only be to hinder the Water from rising so high under the Moon and on the opposite Side of the Earth as the Moon wou'd occasion it to do without that Check : so that the Water, instead of rising up to *a* and *l*, only rises up to *A* and *L*, 8 Foot instead of 10, and the Water is only depressed to *P* and *N* instead of falling as low as *p* and *n*. Thus are the *Neap-Tides* produced by the contrary, as the *Spring-Tides* are by the joint Actions of the Sun and Moon.

As the Sea-shores are very irregular, the Sea of various Depths, and Rivers run swifter or slower into the Sea with and against Tide, and as the Water runs differently into Bays, Gulphs, and Streights, we must not expect the same Regularity of Tides as we have mention'd, any where but in the free Ocean ; but if Circumstances and Facts are rightly examined, we may solve all the *Phænomena* relating to the Tides from the Principles above explained. That learned and indefatigable Astronomer Dr. *Halley* has given an Instance of it in the first Volume of the *Miscellanea curiosa*, and *Philosophical Transactions*, No. 162. where he has enlarged upon Sir *Isaac Newton's* Theory, and applied it to several Cases, which without it would be altogether unaccountable ; shewing by a Solution

tion of very difficult *Phænomena*, that the Theory will reach all Lect. V. Cases known and faithfully related. I refer those who are curious to know more of this Matter to the 38th Proposition of the 3d Book of Sir *Isaac Newton's Principia*, and Dr. *Halley's* Dissertation above mentioned. There they will see why the Water rises to prodigious Heights as 40, 50, nay above 60 Foot in several Places, as at *Chepstow* above *Bristol*, at *St. Malo*, at *Auranches*, in *Normandy*, *Cambaja* and *Pegu* in the *East Indies*. Why there are no sensible Tides in some Places as the *Mediterranean*, the *Caspian Sea*, the *Black Sea*, and the Lakes. Why the Tides are so extraordinary at *Tonquin* in *China*, where there is but one Flood and Ebb in 24 Hours, and twice in each Month no Tide at all. Why the Moon must be a little past the South to make high Water in some Places; and why the highest Spring Tides are not precisely on the *New* and *Full Moons*, nor the *Neaps* on the *Quarters*, but generally the third Tides after them; on account of the Resistance that the Water meets with, which hinders it from following the Moon so fast as it would do if it covered the Surface of a smooth Earth. I shall explain but one Thing more before I leave this Subject, *viz.* the *Phænomena* of the *Æquinoctial* Tides, which are higher near the Vernal and Autumnal *Æquinox* than at any other Time; and least of all at the *Solstices*.

Plate 26. Fig. 8.

57. Let 1 C 2 D represent the Orbit of the Earth, and S the Sun Pl. 26. F. 8. in the Middle; 1 and 2 the Earth at the Time of the Summer and Winter *Solstice*; 3 and 4 the Earth at the *Æquinox*. First, let us consider what must happen to any particular Place, as *Lucajo* above mention'd when it is Summer *Solstice* for that Country, as at 1. If the Moon be in the Line A S, that is, whether it be *New* or *Full*, in the *Syzygy*, as Astronomers call it, A L will be the long Axis of the watery Spheroid, and the Waters will be highest at L and A. Now *Lucajo* being at Noon at L will have high Water, and that in a Place where the Spheroid is highest; but at Midnight *Lucajo* will come to L where indeed it will have high Water, but not near so high as if it had been at A the *Antipodes* of L; the Parallel of *Lucajo*, that is the Plane of it (as well as those of all the other Parallels or Circles in which the several Countries in the World revolve about the Earth's Axis) making an Angle with the Axis of the watery Spheroid, the Whirl of the Surface of the Earth is

LECT. V. is oblique to the Motion of the Water caus'd by the Sun and Moon's Action, and consequently the Rise of the Water is abated by the Shores carrying it another Way. At the Winter *Solstice* when the Earth is at 2, *Lucajo* being at *l* at Noon has indeed high Water; but the Water is not near so high as at *b* the *Antæci* of *Lucajo*: But at Midnight *Lucajo* is carried to λ where it has a great Tide, the Water being there in the highest Place of the Spheroid.

C O R O L L A R Y.

HENCE follows, that at the Summer *Solstice* the Northern Countries have the Flood in the Day Time higher than at Night; but the Reverse at the Winter *Solstice*: and the contrary happens to the Southern Regions.

WHEN the Earth is at 3 or 4 where the *Æquinoxes* are, the Water is highest at C and D, the Sun being then in the Plane of the *Æquator* Æ Q ; so that every Country, that has had high Water in the Day Time as in the Hemisphere Æ 4 Q C , will have as high Water in the Night 12 Hours and a half after; as in the Hemisphere Æ 3 Q D . The centrifugal Force of every Place helping also to raise the Water, every Parallel of Latitude having its Plane parallel to the Axis of the Spheroid. But this will appear still plainer by the 9th Figure,* where P *p* is the Axis of the Earth, Æ Q the *Æquator*, Æ N p n Q P the watery Spheroid raised by the Sun S and New Moon M, L, *Lucajo* having high Water at Noon; it is evident that when *Lucajo* comes to *l* at Midnight it will have the Water as high as it had it at Noon. The same may be said of any other Place, as the *Antæci* of *Lucajo* at N, which, when they come at Midnight to *n*, will have as high Water as they had at Noon at N. So likewise will those that inhabit at Æ find the Water in $12\frac{1}{2}$ Hours as high at Q.

* Pl. 26. F. 9.

N. B. As the Sun is nearest the Earth in the Winter, we have the highest Tides in that *Solstice* which makes the Winter of our Northern Regions; and hence also follows that the greatest Spring Tides are not precisely at the Time of the *Æquinoxes*, but a little before the Vernal, and a little after the Autumnal *Æquinox*.

Concerning




Concerning PENDULUMS.

58. A *Pendulum* is a heavy Body of any kind, which being fastened to a String or Wire hangs down from a Point where it is suspended; as the Pendulum P* hangs from the Center C by the String C P. Now if such a Body be removed from its lowest Point P to any other Point as p ; as soon as it is loosed, it will not only come down again to P from whence it came, but will go on till it rises up to π to an Height equal to that from which it fell, except so much as it is hindered by the Resistance of Air and Friction at C; otherwise it would move forwards and backwards for ever. That Descent and rise of the Pendulum, as for Example, from p to π , is called an *Oscillation* or *Vibration*. *Pl. 26. F. 10.

THE Uses of Pendulums are many; but they are most commonly applied for measuring Time. The whole Doctrine of Pendulums, that is, all that relates to their *Phænomena*, naturally follows from what we have said concerning the Fall of Bodies.

59. I have already shew'd, that if there be an inclin'd Plane, as A B (* Fig. 11.) whose Height is A C and Base C B; a Body falling along the inclin'd Plane from A will be got no farther than D, whilst another Body setting out from the same Point A is fallen down to C the whole Height of the Plane; and that the Point D is found by drawing a Perpendicular from C (the Bottom of the vertical Line which measures the Height of the Plane) to the Plane. If therefore there be several other inclin'd Planes from the Point A to the horizontal Line C B, as A E, A F, A G, &c. the Perpendiculars C H, C I, C K, will shew the Points on those Planes to which a Body falling from A will come in the same Time; now since all the Angles in a Semi-Circle are right Angles, upon the Line A C with the Distance A P (the Half of A C) may be described round the Point P a Semi-Circle which will pass thro' the Angular Points D, H, I, K, &c. Let the whole Circle be completed. The intercepted Parts of the Lines (or Planes) A B, A E, A F, A G, namely A D, A H, A I, A K are now Chords of the Circle. Draw in the other Semi-Circle, the Lines O C, N C, M C, L C, respectively parallel to (therefore equal and equally inclin'd with) the Chords above mention'd. Now, from what has been said it is evident that a Body will fall thro' all those last Chords *Pl. 25. F. 11.
† No. 18. P.
329, 330, 331.

Lect. V. Chords in the same Time as well as thro' the others. Therefore
 if a Circle be set upright, and there be ever so many Chords drawn to the lower part of the Diameter, a Body will fall in the Diameter (the whole Length of it) and in any of the Chords in the same Time, those Chords that are most inclin'd being shortest, and so *vice versa*. Now, since small Arcs of Circles differ but little from their Chords, Bodies falling in small Arcs of Circles, instead of falling in their Chords, will describe those Arcs in Times that are sensibly the same**. For Example; let a pendulous Body C by a String C P hang at P the Center of the Circle; then let it be raised from C to M or any Point between C and M (or the same on the other Side of C) as soon as you let it go it will fall to C in an Arc of the Circle instead of the Chord, and sensibly in the same Time, whether it fall from L, K, or I, because the Chord L C or I C differs from its Arc L C or I C very little more than the Chord C K differs from its Arc C K. But if the Pendulum was to fall in the Arc N M C or M L C, it would be something longer in getting to the Bottom C, than if it had only fallen from K, because the Arc N M C or M L C does exceed in Length the Chord N C or M C much more than the Arc C K does exceed its Chord C K. But if the Pendulum falling from M, be only hinder'd in its Motion by the Air, it will rise nearly up to H which is as high as M, and then there will scarce be any Difference between the Time of the Pendulum's Rise and Fall.

C O R O L L A R Y I.

HENCE follows, that if a Pendulum be made to swing in great Arcs of a Circle, and being left to it self does continually diminish its Vibrations by the Resistance of the Air, the Times of the Vibrations will be a little longer at first than at last; but if the Body makes but small Arcs at first, all its Vibrations without any

** I don't mean that a Body will fall in the Arcs in the same Time that it falls in the Chords; but that in Arcs not very unequal, a Body will fall sensibly in the same Time. For a Body will fall faster in an Arc than its Chord, because it sets out with a greater Declivity; the Time of the Fall in the Arc being to the Time of the Fall in the Chord nearly in the Proportion of 785 to 1000. The Descent along the Chord being perform'd in the same Time as the Descent along the Diameter; it must follow that the whole Vibration, or the Fall and Rise in two Chords, taking up twice the Time, must be perform'd in the same Time that a Body wou'd fall 4 Diameters, or 8 times the length of the Pendulum: But Mathematicians have demonstrated, that the Time in which a Pendulum vibrates in an Arc of a Circle, is to the Time in which it wou'd fall the 4 Diameters above mention'd, as the Periphery of a Circle to those 4 Diameters; or, the Time of the Fall in an Arc: is to the Time of the Fall in a Chord: as a Quarter of a Circle: to its Diameter; that is, nearly, as 785: to 1000.

sensible

sensible Error may be reckon'd *isochronal* (that is perform'd in the same Time) till it come to rest. Lect. V.

C O R O L L A R Y II.

HENCE follows also, that whatever be the Metal or Kind of the Body which is made a Pendulum, provided the String be of the same Length, all the Vibrations, if small, will be perform'd in the same Time (since all Bodies tend to fall with the same Velocity †) L. No. 8. by counting the Vibrations one may measure Time very exactly. For tho' such Bodies, as are specifically lighter than others, will sooner come to rest by the Resistance of the Air, and so they will have fewer Vibrations, yet every one of those Vibrations will be perform'd in the same Time.

N. B. This, and what has been said before will be made more evident by Experiments.

EXPERIMENT XVI. *Plate 26. Fig. 12.*

Pl. 26. F. 12.

60. E C D is a triangular Board of Wood set upright, with an Arc of a Circle drawn from the Center *b* round *o* to *g*, and another drawn round the Center *a* to *g* on the other Side, and the Divisions on each Side are equal. From *a* and *b* let the two Balls or Pendulums hang so as just to touch one another when they are at lowest, which they will do because the Points *a*, *b*, are just distant from one another the Distance of the Balls Centers. If the Ball A be raised to 7, 8, or 9, and the Ball B to 1, 2, or 3; and you let them go both at the same Time, they will meet exactly at *o*, tho' they have described unequal Arcs; and so they will, let the Arcs be as different as they can be made upon this Instrument, at least as far as Sight can determine, when the biggest Arc is not of above 20 Degrees, as it cannot be in this Instrument.

EXPERIMENT XVII. *Plate 26. Fig. 13.*

61. Take 3 Balls P, P, and Π, one of Lead, one of Ivory, and one of Cork, and, having fasten'd each of them to Strings of equal Lengths, let them hang from the horizontal Wire W on the Top of the Stand S; then raising them all up to equal Heights, they will, when you let them go at the same Time, perform their
Vibra-

Lect. V. Vibrations in the same Time; for tho' the Cork will come sooner to rest than the Ivory, and the Ivory than the Lead; and while the Lead is still vibrating in the Arc P p, the Ivory vibrates only in the lesser Arc P p, and the Cork in an Arc still less, viz, $\Pi \pi$, one may observe that they always come to the lowest part of the Vibration at the same Time.

Pl. 27. F. 1.

EXPERIMENT XVIII. Pl. 27. Fig. 1.

62. ABOUT the Pin C for a Center let the Pendulum P swing, having first lifted it from F to P, and it will describe the Arc P F e rising up to e sensibly as high as P in the horizontal Line P e. Then at the Point A, the Middle of the said horizontal Line, stick in the Pin A, and the String of the Pendulum will be check'd and stopp'd by that Pin; but the remaining Part of the String below from A to F will be at Liberty, so as to let the Pendulum rise up to B a Point in the horizontal Line above mention'd in an Arc of a Circle whose Radius is A F. If the Pin be fix'd at a, the Pendulum will rise in the Curve F D; and if two Pins be fixed at b and c, the Pendulum will rise in the Curve F E, made up of three different Arcs; of which the first has its Center at C, the next at b, and the last at c. This shews (as has been already explained) That *whatever Curve or Curves a Pendulum rises in after its Descent, it will go up to the same Height from whence it fell, except what must be allowed for the Resistance of the Air*; the Velocity of the Pendulum at the lowest Point F being the same as it would have acquir'd in falling directly from A: and, that that Velocity is capable of carrying the Body (in such a Direction as Gravity cannot act against it) twice the Space A F by an uniform Motion in the same Time that it falls thro' the Space A F by an accelerated Motion, as has been already proved; † but it will be made visible by the following Experiment.

† No. 15. P.
324.

EXPERIMENT XIX. Plate 27. Fig. 2.

63. To a String about 40 Foot long from the Center C hang the Pendulum P, which when at Liberty will hang down to p and swing in the Arc p q. From the lowest Point p on the Arc p q measure two Foot or the Arc p π , which will be a Line sensibly horizontal, the vers'd Sine of such an Arc being but about half an Inch. At the Distance of half a Diameter of the Pendulum beyond π fix a ver-

vertical Obstacle or wooden Plane $O \pi$; and another such on the other Lect. V.
Side of p at the Distance of about one Foot and an half, but in an horizontal Position, so that the Surface of it may be even with the Bottom of the Pendulum, namely, in the Line $p \pi$. Let the Pin N be fixed perpendicular to the String (that is, in an horizontal Position) at the Height of one Foot above p . A Mark or strong Point, as n must also be made in the Wall or Plane behind the String of the Pendulum a little above p . Then having raised the Pendulum up to P in the Line $H b$, that is, just one Foot high, let it go again and observe the very Instant of Time that a Mark in the String of P comes to the Point n , that you may just then let fall the Body H from the horizontal Line $H b$ to the Obstacle M ; and you will find that the Ball H strikes M at the very Instant that P strikes the Obstacle $O \pi$ at π , P moving horizontally the Space $p \pi$ which is two Foot, while H falls the Height $H M$ which is but one Foot. This may be done so exactly that the Sound of the two Strokes will be but as one.

64. It has been observed, that a Pendulum, whose Length is 39 Inches and 2 Tenths (English Measure*) from the Center of the * Ann. 9.
Ball to the Point of Suspension, will perform one Vibration in one Second of Time; that is, 3600 in an Hour. It has been tried with a Pendulum of 50lb Weight made of a lenticular Figure the better to cut the Air, and a fine Steel Wire was used instead of a String; and with such an Instrument the Vibrations have continued a whole Day. Hence it is that Pendulums will serve to measure Time equally and exactly, Regard being had to the Length of the Pendulum, be the Weight of any kind, or Bigness.

65. THE shorter Pendulums are, the shorter will be the Time of their Vibrations; tho' the Times of Vibration will not be as the Lengths of the Pendulums, but as the square Roots of their Lengths: as for Example, a Pendulum of 13 Foot and near an Inch long (that is, 156,8 Inches) which is 4 times as long as a Pendulum that swings Seconds, will perform its Vibrations in 2 Seconds; and if we would have a Pendulum to go twice as fast as the Second-Pendulum, that is, to swing twice in a Second, we must make it 4 times shorter; that is, 9 Inches and 8 Tenths. The reason of it is evident from the following

DEMONSTRATION. *Pl. 27. Fig. 3.*

Pl. 27. F. 3.

A E B G, and D F B are two Circles whose Diameters A B and D B are to one another as 4 to 1. Now it has been demonstrated,* * L. 5. No. 15.
B b b that

Lect. V. that if a Body fall thro' a certain Space, as A B, in any determinate Time; it will fall thro' only $\frac{1}{4}$ of that Space, as D B, in half that Time. But it has also been proved that Bodies will fall in the

L.5.No.59. Chord of a Circle in the same Time that they fall in the Diameter; therefore a Body at E, will fall thro' the Chord E B in twice the Time that a Body at F will fall thro' the Chord F B. Now as the Chord E B differs from the Arc E B (or is shorter) in the same Proportion as the Chord F B differs from its Arc F B (both being supposed Arcs of the same Number of Degrees) the Difference will be proportionable in the Descent of the two Pendulums C E and c F; consequently E will be twice as long coming down to B, as F coming down to B; but the whole Vibration which carries up E to G, and F to f will, in both, be double the Time of their respective Falls. Therefore a Pendulum, to vibrate twice as fast as another, must be $\frac{1}{4}$ times shorter. Q. E. D.

Pl. 27. F. 4.

EXPERIMENT XIX. Pl. 27. Fig. 4.

ON the horizontal Wire W of the Stand represented in the 13th Figure of Plate 26, whose Top is here expressed in Fig. 4, hang a Pendulum C P whose Length is 39, 2 Inches, and upon the same Wire a little forwarder hang the Pendulum c p of a quarter of that Length. Let both Pendulums fall at the same Time from any Height to which they have been raised, as P and p, and you will find that p will be got to f (having performed its whole Vibration) when P is only come to B; when P is come to F, then p is come back from f to p; when P is come back to B, p is got to f; and at last when P is come back to P, p will also be come to p from whence it first set out; so that the two Pendulums will begin to fall together, every other Stroke the short one vibrating in half Seconds, whilst the long one vibrates in whole Seconds.

Pl. 27. F. 5.

EXPERIMENT XX. Pl. 27. Fig. 5.

TAKE an Iron Bar, either square or round, provided it be of the same Thickness in all Parts, whose Length must be 58, 8 Inches, measuring from a little Hole to receive the Wire of the Stand to its End; and having thrust the Wire through its Hole at A, and suspended a Second-Pendulum at C, let both go down from P and B at the same Time, and their Vibrations will be *isochronal*, that is, performed in the same Time. This shews that any streight, even Rod, square or round, of any Metal whatsoever, will perform its Vibra-

Vibrations in the same Time that a Pendulum of $\frac{2}{3}$ of the Length Lect. V.
of it vibrates ; as if the whole Matter of the Bar had been collected
into the Point *p*, which is therefore called the Center of Oscillation.

N. B. *If any Person holding a Bar, as A B, at its End A, should strike a Blow with it, the greatest Stroke possible with the same Strength would be made when the Point p strikes against the Obstacle ; for this Reason, that Point, which is here the Center of Oscillation, may also be called the Center of Percussion. (a)*

66. A Pendulum is very successfully applied to Clock-Work for measuring equal Time, as may be seen by looking into any Clock. But to shew in what manner it regulates the Motion, one Example will be sufficient, which I give here in describing a little Clock or *Chronometer*, which Mr. *George Graham* contrived and made for me some Years ago, so exact as to measure a small Part of Time very nicely, even to the 16th Part of a Second.


To one End of an horizontal Axis is fixed a contrate Wheel of 120 Teeth, and, at the other End there is a little Barrel which on its outside receives the String of a little Weight to turn the Wheel round ; the Wheel with its Teeth moves an horizontal Pinion of 15 Leaves (or Teeth) fixed to the Bottom of a vertical Axis on whose Top is the Balance Wheel, which is a contrate Wheel of 15 Teeth cut askew ; that is, one Side of every Tooth is perpendicular to the Plane of the Wheel, and the other cut obliquely with a Curve. Just over this last Wheel there is a little Axis or Rod of Steel placed horizontally, which has two little Pallets at right Angles to each other, so placed as to be alternately struck by the upright Part of the Skew-Teeth on the opposite Sides of the Balance Wheel, in such manner that not one of the Teeth of that Wheel can pass by without a Stroke be made against each Pallet to give a quarter of a Turn to the Axis above-mentioned. At the other End of this Axis is fixed a Wire, at whose End is a Brass Bob whose Center of Gravity is distant from the horizontal Axis 2,45 Inches, that is, near two Inches and a half, or the 16th Part of the Length of a Pendulum vibrating Seconds. Therefore, as the Weight carries the Wheel-Work round, the Skew-Teeth of the Balance Wheel must strike the Pallets fixed to the horizontal Axis about which

(a) *In some sort of Bodies the Center of Oscillation and the Center of Percussion is not the same ; but the Discussion of that Matter being foreign to my present Purpose, I shall say no more of it here.*

Leſt. V. the little Pendulum moves ; and as that Pendulum, on account of its Length, vibrates 4 times in a Second, † and there muſt be 2 Strokes againſt the Pallets for every Tooth that paſſes by in the Balance Wheel, ſo there muſt be 2 Swings of the Pendulum, or half a Second in Time, for every Tooth of the great Wheel that is brought forward by the Pivion (whoſe Number of Teeth is equal to that of the Balance Wheel) and as the great Wheel has 120 Teeth, there will be performed 240 Vibrations of the Pendulum whiſt the great Wheel goes round ; therefore an Hand or Index fixed to the great Wheel will go round in One Minute pointing to 60 large Diviſions for Seconds on the Dial Plate, which Diviſions are again ſubdivided into 4 to ſhew the Quarters of Seconds. But then beſides this, there is a Quadrant of a Circle of a *Radius* equal to the Length of the Pendulum, which is divided into 4 Parts by 5 little Braſs Pins, fixed to an horizontal Axis ; which Quadrant ſerves not only as a Detent to ſet the Machine a going in a 16th Part of a Second, but alſo to ſtop it in the ſame Time, the Pins ſtopping the Pendulum in any fourth part of its Vibration ; for in fixing the Pins, Regard was had to the Time of every fourth part of the Vibration, the 2 Spaces between the Pins next to the loweſt Part of the Quarter of the Circle which the Pendulum vibrates in $\frac{1}{4}$ of a Second, being ſo much greater as the Pendulum moves ſwifter in that Part of its Vibration.

THIS Chronometer is of great Uſe for meaſuring ſmall Parts of Time in Aſtronomical Obſervations, the Time of the Fall of Bodies, the Velocity or running Waters, and fit for many other Purpoſes, where a ſmall Space of Time is to be meaſured nicely, as from 3 or 4 Seconds to a Minute or two ; but it is not adapted to meaſure long Spaces of Time very exactly ; becauſe, tho' it ſeems to vibrate exactly a quarter of a Circle every Vibration, yet it does not really do ſo, and the Difference of the Lengths of the Vibrations in ſuch large Arcs makes a Difference in Time ; and however ſmall that Difference is, yet a great many of them create a ſenſible Error. In this Machine, there is ſometimes an Error of $\frac{1}{4}$ of a Second in 14 Seconds. This is a Fault which all Clocks are liable to, that have ſhort Pendulums which ſwing large Arcs of a Circle.

67. BUT there is a Curve, which is neither a Circle nor an Ellipſe, in which all the Vibrations of the ſame Pendulum, whether long or ſhort, are performed in the ſame Time, and when Clock-Work is

is regulated by a Pendulum moving in that Curve, a long Space of Lect. V. Time may be exactly measured. This Curve is a *Cycloid*, of  which here follows the Description.

68. IF a Circle in a vertical Position rolls along the horizontal Line A B* (as the Wheel of a Carriage does upon the Road) till it has made ^{* Pl. 27. F. 6.} one Revolution, that Point of the Circle, as A, which touched the Plane, will rise up from the Line and be at *a* when the Circle has rolled half way, having described the Curve A *a*, from whence (as the Circle goes on) it goes downwards in the same kind of Curve, till it comes to touch the Line A B at B; the Point C, which was uppermost at the beginning of the Motion, having been down at *c* when A was at *a*, and now being returned to *k* when the Circle is in the same Position as before its Revolution began. The whole Curve A *a* B so described is called a *Cycloid*, and the Circle A C the *generating Circle*, and the Line A B the *Base of the Cycloid*. † † Ann. 10.


IT is evident from the Formation of the *Cycloid*, that the Base of the *Cycloid* is equal to the Circumference of the generating Circle. Several other Properties of the *Cycloid* have been demonstrated by Mathematicians, † whose Demonstrations it would be tedious to repeat here; therefore we shall take them for granted and mention them as we have Occasion for them in our Conclusions ^{† Dr. Wallis, Craig, Dutton, Bar-nouilli, &c.} about Pendulums.

69. Let us now invert the *Cycloid*, so as to have A B the Base at Top and the *Cycloid* underneath.* It has been demonstrated, ^{* Pl. 27. F. 7.} that if from any Point in the *Cycloid*, as E or *p*, there be drawn a Line parallel to the Base, and from the Point where that Line cuts the generating Circle when it has performed half its Revolution (that is, when it is at G) there be drawn a Chord such as *e* D or *q* D, the intercepted Arc of the *Cycloid*, as E D or *p* D will be double the Chord *e* D or *q* D; and so one half of the *Cycloid*, as A D will be double the Diameter G D of the generating Circle. Now since it has been shewn that a Body will fall in the same Time in the Diameter G D, and in the Chords *q* D, *e* D, a Body must fall in the Arcs A D *p* D and *e* D in equal Times; because each of them is the double of the above-said Lines G D, *q* D and *e* D: so that when a Pendulum swings in a *Cycloid*; all the Vibrations, however unequal, are isochronal.

Now

Lect. V. Now, in order to make a Pendulum swing in a *Cycloid*, there have been contrived cycloidal Cheeks on each Side of the Center of Suspension, that the String applying it self to them should shorten the Distance of the Pendulum from the Point of Suspension, so as to make it describe a cycloidal Arc instead of a circular one. The cycloidal Cheeks are thus made. Let the half *Cycloid* A D be transferred to C B so that the Point D may be on B, and the half *Cycloid* B D be transferred to C A so that the Point D may be on A. Now if there be a couple of Cheeks of Wood or Metal cut convex so as exactly to fit those Semi-cycloids, and a Pendulum as P of the Length C D, double the Diameter G D, has its Point of Suspension just between the said Cheeks at their Point of Contact C, such a Pendulum will no longer vibrate in the Arc H D I (whose Center is C and Radius the Length of the Pendulum) but in the *Cycloid* A p D F B by the shortening of the String as it applies it self to the cycloidal Cheeks as at C S p.

THERE is another Way to make a Pendulum vibrate in Arcs so nearly cycloidal as to have no Error in their Measure of Time; and that is, by having the Pendulum very long, and making it only vibrate very small Arcs. For Example, let C D be a half-Second Pendulum, that is, 39,2 Inches long, and let it only vibrate from P to δ about 4 or 5 Degrees. It is plain that the Circle and Cycloid co-incide at D and some Degrees on either Side; therefore the Pendulum that swings in the circular Arc H D I, describing only the Part of it P δ , may be considered as swinging in a cycloidal Arc; and if it should sometimes go from D only to δ , it would describe that little Arc in the same Time as the greater Arc D δ . N. B. It is better to use the long Pendulum than the cycloidal Cheeks; because if the String or Wire, by which the Pendulum hangs, strikes strongly on the cycloidal Cheeks, the Reaction of those Cheeks, which are in some measure elastick, will add to the Force of Gravity by which alone the Pendulum must swing to have its Vibrations isochronal in a *Cycloid*; so that when any such Shock happens, the Vibration will be too quick. But the long Pendulum without a very great Shock (such as would send it up to p) will not so sensibly vibrate out of the *Cycloid* as to cause any Error in the Measure of Time. So that is now used very successfully in Clocks for keeping Time at Land; the Shocks at Sea being too great to have a long Pendulum Clock go so well in a Ship.

70. BEFORE we quit this Subject, it may not be improper Lect. V. to mention a remarkable Property of the *Cycloid*, which is this, *viz*,  that it is *the Line of the swiftest Descent*; that is, if a Body is to move from A to D,* there is no Line of any kind which can be* Pl. 27. F. 7. drawn from the upper to the lower of those Points, along which a Body will descend so fast as in the Semi-cycloid A D; neither an Arc of a Circle, nor even the streight Line A D, tho' it be so much shorter; for the Body at first setting out from A descends in so steep a Direction (that is, is so much acted upon by Gravity) that it acquires a great Velocity so as to carry it the more swiftly in the lower Part of the Curve which is less steep: and a circular Arc which would be more steep at Bottom is less so at Top. If an inclin'd Line as A F, be drawn from A beyond the lowest Point of the *Cycloid*, and a Body goes down that Line, whilst another runs from the same Point along the *Cycloid*, the greater Celerity of the Body in the *Cycloid* will still appear more evidently. This Property, and that of having the Descent of a Body from any Part of the (inverted) *Cycloid* to the Bottom, exactly in the same Time, will be illustrated by the following Experiments.


EXPERIMENT XXI. Pl. 27. Fig. 8.

Pl. 27. F. 8.

71. THE Machine B H M D D, represented by the Figure, is made of Wood about 10 Inches high, two Foot long and 2 Inches thick. From I to F it is cut into a Channel as wide at Bottom as at Top, in the Form of a Semi-cycloid inverted, the lowest Point being at F, from whence the Channel is continued horizontally for the Length of a Foot from F to G, the Walls of the Channel being H H and I I, and this Channel is very smooth and divided into two Channels by a thin upright Brass Partition L L from the Top between H and I to the farther End at G. Upon the Partition are mark'd Divisions, which from F to G are equal, but unequal from F to I, in such manner as to denote equal Heights, H the Beginning of the Channel is 9 Inches above the Level of F, O O are two wooden Stops for the Channels, and to be fixed in any Place by means of a Skrew at the End of each of them. The whole Instrument may be set upright and horizontal by means of three Skrews such as C, C, and the Plummet N M.

Two Brass Balls, of half Inch Diameter each, are to move in the two Channels.

FIX

 Lect. V. Fix the two Stops exactly at F, and the Brass Balls, tho' you let them go from different Parts of their respective cycloidal Channels, will strike them at the same Time; which will also be very sensible to any one who holds a Finger in each Channel at F, whilst another Person lets the Balls run down from unequal Heights.

Now let the Stops be shifted, and one of them be fixed 4 Inches beyond F towards G, and the other 6 Inches beyond F; and let the two Balls fall at once, the one from the Height of 4 Inches, and the other from the Height of 9. That which falls from 4 Inches in Height, will go 4 Inches beyond F; and that which falls from 9 Inches in Height, will go 6 Inches beyond F, each striking the Obstacle in its own Channel exactly in the same Time as the other. *Four* and *Nine* are the Spaces fallen thro', whose Roots 2 and 3 express the respective Velocities of the Balls, which are shewn by one of them running 4 Inches in the horizontal Part of its Channel, the other runs 6 Inches in the other horizontal Channel.

Pl. 27. F. 9.

EXPERIMENT XXII. Pl. 27. Fig. 9.

A E C *b* is another Machine with a cycloidal Channel, wherein the whole *Cycloid* is hollowed in from B to *b*, its lowest Part being at C. There is a freight Channel A *a* moveable round the Point A as a Center to apply on the back-side of the *Cycloid* as the Machine stands upright, so that its Channel may be in the Plane of any Chord drawn from the Point A to any Part of the *Cycloid*. Let this Trough or Channel be first set in the Line A C; then let one of the Balls fall from B in the cycloidal Channel, whilst the other begins at the same Time to fall from the same Level in the freight Channel, and you will observe, that the Ball running down in the *Cycloid* will be a good Way beyond C, when the Ball running in the freight Channel is just got to it; as may still be made more sensible by fixing one of the Stops above-mentioned (in the Explanation of the last Figure) over-against C in the freight Channel, and the other between C and *b* in the cycloidal Channel. If the freight Channel be fixed beyond the Point C as at A *a*, the Ball in the *Cycloid* will come from B to *b* and be down again at C when the other is come from A in a freight Line to G.

72. WHEN a Pendulum of a proper Length to swing Seconds is applied to a Clock; there is yet another Error which it is subject to,

to, arising from the Nature of the Materials. For all Metals are Lect. V.
 subject to be dilated by the Heat and contracted by Cold; so that
 if the Rod of the Pendulum (that is, the Wire whereby it hangs
 instead of a String) be adjusted to its right Length in the Winter
 Time, it will become too long in Summer by the Heat, and there-
 fore vibrate too slow; as on the contrary, if it had been adjusted in
 Summer, it would be shortened in Winter by the Cold, and by its
 quicker Vibrations make the Clock go too fast. The Difference of
 Length in a long Pendulum Rod will sometimes amount to about
 the 40th Part of an Inch, which in many Vibrations will make a con-
 siderable Alteration in Time. But there is a Contrivance of a Skrew
 to raise or sink the Weight of a Pendulum; so that one may keep
 it always to its due Length, regulating it with a Thermometer.

*The Way to measure the least Alteration of Dimensions in Me-
 tals so as to make it sensible, I shall shew in the Notes.** * Ann. 11.

73. BESIDES all this, Pendulums are liable to alter the Times of
 their Vibrations on account of the Figure of the Earth, which
 is an oblate or flatted Spheroid like a Bowl, the Earth (as we have
 already mentioned) being about 31 Miles higher at the Æquator
 than the Poles. For when we travel Southward, we shall find our
 long Pendulum Clocks (besides what was allowed for the Heat)
 to go too slow, as Dr. *Halley* and several other Astronomers have
 observed, who were obliged to shorten the Pendulums of their
 Clocks before they could make them keep true Time. Mr. *Richer*,
 in order to examine this *Phænomenon* fully, made Observations on
 the Vibrations of a simple Pendulum (that is, a Pendulum not
 applied to a Clock) for 10 Months together, and found, that a Pen-
 dulum to vibrate Seconds under the Æquator must be about one
 10th of an Inch shorter than in our northern Latitude; without
 which the Clock, which it regulates, will go too slow between two
 and three Minutes in a Day.

FROM these Observations on Pendulums Sir *Isaac Newton*
 shews what the true Figure of the Earth is; tho' contrary to the
 Opinion of some Gentlemen of the Royal Academy of Sciences at
Paris, who assert that the Earth is an oblong Spheroid (like an
 Egg) higher at the Poles than the Æquator about $\frac{1}{6}$ of the whole
 Diameter. *I shall, in the Notes, shew how these latter came to
 be mistaken.** * Ann. 12.

Lect. V. 74. WHEN we consider what Velocity the diurnal Rotation of the Earth must give to Bodies on its Surface, especially under the Æquator or near it (where Bodies are carried round at the Rate of about 1040 Miles in an Hour) we may easily see the Necessity of its being higher under the Æquator; because unless there was a Descent towards the Poles, the centrifugal Force would drive all the Waters towards the Æquatorial Regions, which would thereby be drowned, whilst not only the Polar Regions, but even the temperate Zones, would be left dry. And if the Earth was ever in a fluid State, or its Substance was soft and yielding, the Author of Nature giving it the diurnal Rotation that it has, it must from a spherical put on a spheroidical Figure, swelling under the Æquator till it came to have the Figure that it now has. A swifter Revolution round the Axis would still make the Difference greater between the Æquatorial and Polar Diameters. And this is observed in *Jupiter*, which, by some nice Observations of that late ingenious Astronomer the R. Mr. *Pound*, appears to have its Æquatorial Diameter $\frac{1}{2}$ longer than its Polar Diameter, agreeable to its swifter diurnal Revolution, which is performed in about 9 Hours.

EXPERIMENT XXIII.

UPON an Axis of Iron, that could be made to turn swiftly (by means of the Wheel of the whirling Tables, whose String went round a Pulley fix'd to the said Axis) I slipp'd on two Iron Hoops, whose Planes intersected at right Angles, representing two Colures, which, being of a Spring-Temper, stood at first in such manner as to be $\frac{1}{6}$ part longer in that Diameter that co-incided with the Axis, than in their Æquatorial Diameter; this Proportion being the same that Mr. *Cassini* supposes to be between the Axis and Æquatorial Diameter of the Earth. Two circular Plates, to which the said Hoops were rivetted, had square Holes, thro' which the Axis pass'd, so that the two Poles of the oblong Spheroid, which the Hoops describe in their Revolution, might approach together in such Manner, as to let them put on the Form of a true Sphere, when, by the whirling, the Æquatorial Diameter of the Machine swell'd and over-power'd the Elasticity of the Hoops. A greater Degree of Swiftmess turn'd the Sphere into an oblate Spheroid of Sir *Isaac Newton's* Figure: A Velocity still greater makes the Disproportion of the Diameters, such as that of *Jupiter*; and still the Æquatorial Diameter increases with the centrifugal Force.

ANOTHER

ANOTHER Hoop with a Catch, representing the *Æquator*, Lect. V. shews (in the Experiment) the Increase of the *Æquatorial Circumference*; and an Index apply'd to the Frame, shews the Increase of the Diameter.


As soon as the Revolution of the the Machine ceases, the Colures, Meridians or Hoops, return to their elliptical Figure, whose vertical longest Diameter is the Axis of Revolution.

THOSE that would know more concerning Pendulums, may consult *Huygen's* Posthumous Works, and Dr. *John Keil's Introductio ad veram Physicam* at the latter End; where there are several Propositions concerning the circular Pendulum, which I have not room to say any more of in this Place, having already made this Lecture too long; but yet I cannot put an End to it, before I have acquitted my self of my Promise, to give an Account of the *Bow* or *Spring*, so far as it is a mechanical Organ.

75. THIS Engine is so universally known, and the Practice of it so generally understood, and withal so natural and easy, that all Mankind, in all Ages of the World, and in every Quarter of the yet habitable Earth, even among the most rude, ignorant and barbarous, as well as among the most civil, polite, and learned, have been found expert and skilful in the Use thereof; and even in all that vast Tract of Land, extending so far towards the North, and towards the South Pole, wherein scarce any of those mechanical Engines which depend upon the Balance, or Beam, was known, namely the Leaver, the Capstane, the Pulley, the Screw or the Wedge, nay where the Use of the Roller and Wheel was altogether unthought of; yet even through all this vast Tract of Land, and in all the adjacent Islands, the People were found very dextrous and skilful in the Use of the *Bow*, and knew how to convey the Arrow by means thereof so as to hit and pierce the design'd Mark; so that it seems to be an Engine most obvious and easy to be contrived by Mankind: And, yet after all, I do not find that any one of the Ancients and Moderns has reduced the Nature, Properties and Power of Springs to a certain Theory, and Calculation; tho' divers have written much concerning it, and have made many Attempts, and Experiments about it; till Dr. *Hook* of late reduced it to a Certainty, by shewing,

Lect. V. *That the Power of a Spring, of what kind soever, does encrease in the same Proportion with its Tension*, whether it be by Compression or Condensation, Distension or Rarefaction; so that if the Strength of one Pound, moves it one Degree out of its natural State, then two Pounds will move it two Degrees, and three Pounds move it three Degrees, and so on. Now this mechanical Organ, contrary to all others I have already specified, receives all the Power or Motion which is impressed on it, and retains it till it be left at Liberty, or no longer any farther extended, or forced with greater Power; but then, instead of proceeding and going forwards in that way in which it had been moved, it recoils, and goes quite back again in a contrary Direction; and by *Re-action* exerts all the Power and Motion it has received upon the Bodies it meets with in its way as it goes back, and carries them with great Swiftneſs; as having received all the Power that had by Tension been impressed on the *Bow* by the Strength that bent it. By this means all the Strength of the Arm exerted to bend the *Bow* one way, is by the *Bow* suddenly communicated the contrary way to the Arrow, which makes it move with that Swiftneſs and Speed after it is discharged, continuing the Motion by the first Law. Now, here possibly may be enquired, how it comes to pass, that the *Bow*, which in its bending, or drawing, is only moved by a slow and heavy Motion (nay, it may be, is kept some Time still, and without any Motion at all) should yet retain so great a Quantity of so swift a Motion or a Power of producing it, in another Body, tho' as to Sense it seems to have no Motion at all of its own; whereas all the other Organs, which produce or effect such a powerful or rapid Motion, have that Motion sensibly acting in them, when they produce their Effect, proceeding directly forward in the straight Line in which they moved before the Stroke; so that there is nothing but the Continuation of the whole acquir'd Motion.

To which I answer, that tho' the physical Reason of the Power of Springs, may not possibly appear so evident to Sense as that of a Weight lifted up and then let fall; yet it is much of the same Nature, as it may be made plainly evident; for as in the lifting up of a Weight to a certain Height, and thence letting it fall, the Power of Gravity makes that Weight descend back again to the Place whence it was lifted, with greater Force as it is lifted higher; and so in its Descent enables it to exert as much Force or Pow-

er as was spent in the lifting it up to so great a Height: So a Spring, *Lect. V.*
which has a Power like Gravity, or an Activity equivalent to it; 
and within it self, but with this Difference, that it acts any way
as it is directed (whereas Gravity only acts downwards) so as in its
Return to exert all that Force and Power, which was spent in
bending it to that Tension; and as the Motion of Gravity or De-
scend, whatever the Velocity of Motion was that rais'd the Weight,
carries it back to the Place whence it was rais'd, by one uniform Pro-
gress of Acceleration, which is peculiar to the Power of Gravity,
as we have shewn; so the Recoil of the Spring, with whatever
Velocity it has been bent, unbends it self by such Degrees of Pow-
er and Velocity, as are proper and peculiar to the Nature of
Springs, which, as I mentioned before, are so much the more pow-
erful, by how much the more they are bent; so that to make the
Comparison exactly to answer, we must not conceive a single
Weight to be lifted up, as in the Example of Gravity, because
the Power of Resistance of Gravity, in such a Weight, is the same
at the last as at the first, as to Sense; for the same Power that lift-
ed such a Weight one Foot high, at beginning, will lift it one Foot
higher after it has lifted it to that Height, or a third Foot after a
second; for Instance, if the Force of a Pound Weight will lift it
one Foot high, that Force of one Pound continu'd, will lift it ano-
ther Foot high, and (continu'd) a third Foot high: But in a Spring
it is otherwise; for if the Force of one Pound be requir'd to bend
the Spring one Inch, then there must be the Force of two Pounds
to bend it two Inches, and three Pounds to bend it three Inches;
so that the Power of a Spring may be more properly represented
by the lifting up of a Chain, or a String of Beads, or Weights
that lie on the Ground; for in lifting up the first Link, or Bead,
there is requir'd one Pound, for instance, but lifting it higher
there rise two Links or Beads, and so the Force of two Pounds
is requir'd; and if higher, three Links or Beads rise, and so on:
Or more exactly yet, 'tis represented by lifting a Cylinder of
Timber endwise out of the Water, which is of near the same spe-
cific Gravity with the Water, or forcing Water to rise in a
Pipe by a forcing Plug; for in these Cases, as the Cylinder is rais'd;
higher above the Surface of the Water, out of which it is rais'd,
so does the Gravity of it increase, and the Power that lifts it
must increase in the same Proportion, which is exactly the same as
that of the Spring; and so either of these Examples becomes a
most proper Instance to illustrate the Power of the Spring, and the
Manner

Lect. V. Manner of its acting, and tho' the Cause of the Necessity of the Increase of the Power to move the one, be more sensible and intelligible than the like Necessity of Increase to bend the other; yet if we further proceed to shew the true Reason and physical Explication of the Power of Gravity, we shall find it to be as difficult, and as far remov'd from our Sense, as the physical Cause and Reason of the Power and Increase thereof in the Spring; it is enough therefore for Geometry, to know, that the *Phænomena* are so, and thence to examine and demonstrate the Consequences that will follow. Now the Power of Gravity in accelerating the Velocity or Motion of falling Bodies, as it was first found out, and discover'd by *Galilæo*, so was the perfecting of the Theory thereof, as to the Motion of Descent of heavy Bodies in inclin'd Planes, and the Curvature of a Parabola, very far promoted by him, and many curious Theories concerning it deduc'd, as we have shewn in this Lecture; and the same Theory has been farther improv'd, and carried on by *Torricellius*, and others also since him. But all these proceed upon a Supposition, that the Power of Gravity in all the Points, or the Perpendiculars of Descent are mathematically equal and uniform; and indeed, as to all Motions of terrestrial heavy Bodies, they are found so as to Sense; but as to mathematical Exactness and physical Theory, they are really otherwise; that is, the Perpendiculars of Descent are not parallel one to another, but converging and growing nearer together, as they approach nearer to the Center of the Earth; and the Power of Gravity in every Point of those Lines, is not equal and uniform, but stronger as it is nearer the Earth, and weaker as it is farther off, * according to a certain Proportion, which makes the Curve Line of a projected Body not Part of a parabolical Line, as *Galilæo*, *Torricellius*, and the rest have suppos'd; but perhaps, Part of an Ellipse, one of whose *Foci* is in the Center of the Earth. Notwithstanding which, the Theories and Problems deduc'd geometrically from them upon such *Phænomena* as they had then discovered, and upon the Hypothesis they had thereupon taken, are highly valuable; and the Deductions truly made and demonstrated, which is the utmost Performance that can be expected from Geometry. But now these Things have been carried much farther by Sir *Isaac Newton*; the Power of Gravity is determin'd and brought to a Certainty, by what Degrees it decreases the farther it acts from the central Body; by which Means the true Reason of the Motion of the Celestial Bodies, and the Lines or Orbits in which they pass,

*L. I. No. 17.

pass, and the Velocities with which they move in their Orbits, is Lect. V.
perfectly adjusted, as we have fully explain'd.

BUT in this, the Example taken from Gravity is (as I said before) defective, for solving the Effect of a recoiling Spring, upon a Body mov'd by it; for in the Spring, the farther it is bent the more powerful it is; but in Gravity it is the contrary.

THE Effect therefore of the Power of the Spring, in this Particular, must be sought out from the Theory of the Nature of Springing, which is, as I but now mentioned, stronger and stronger, as it is the more bent, that is, in the same Proportion with the bending; and consequently (since Action and Re-action are equal) the Velocity of the Body mov'd by it, will always be in the same Proportion with the Degree of bending, whether it be longer or shorter, that is, the farther the Spring is bent, the quicker is the Motion of its Recoil, and the less it is bent, the slower. If it be bent one Degree, the Velocity is as One; if two Degrees, the Velocity is as Two; if three, as Three; and so for any other Degree: And consequently, the Velocity being always proportion'd to the Space it is mov'd by it, the Time of the Body's moving, whether a longer, or a shorter Space, will always be the same, or *isochrone*, which is the same with the Motion of a Pendulum, whose Weight is mov'd in a Cycloid, as we have shewn.

THIS would lead me to explain several Things concerning *Elasticity*, and the Laws of elastick Bodies in their Congress; as also the Effect of that Property in the Air for the Conveyance of Sounds; but these Things I shall treat of in the second Volume, this being already swell'd to too great a Bulk.

ANNOTATIONS upon the Fifth LECTURE.

1. [3. Page 286.—*the Body will go thro' the whole Diagonal by the Action of the said two Forces.*]

† Pl. 28.
F. 1.

IF Three, Four, Five, or any Number of Forces act upon a Body at the same Time, the same Rule will serve to shew the Line in which the Body will move, which will be the Result of all the Forces compounded. As for Example, if four Forces, whose Quantities and Directions are represented by the Lengths and Positions of the Lines AB, AC, AD, AE, † act at once upon a Body at A; first let us draw Ba and a C respectively equal and parallel to the Lines AB, and AC, and having so compleated the Parallelogram AB a C, let us draw the Diagonal Aa, in which the Body would move if only the two Forces AB and AC acted upon it. Now, having shewn, that the Body, by only one Force, whose Quantity and Direction are represented by the Length and Position of Aa, would move just as it does by the Action of the two Forces AB and AC; we may consider them as only one Force Aa, which comparing with the Force AD, we shall have the Diagonal Aa of the compleated Parallelogram Aa a D, being the Line in which the Body acted upon by the three Forces AB, AC and AD will move, just in the same manner as if only two Forces, as Aa and AD had acted upon it. Three Forces therefore being thus reduc'd to one, represented by Aa, we may combine Aa with AE the fourth Force, and so have the Diagonal Aα of the compleated Parallelogram Aa α E, the Line in which a Body will move by the joint Action of four Forces.

N. B. *Which soever of the four Forces you begin with, the Line Aα will always be the Diagonal of the last Parallelogram.*

In the same manner may be found the Result of the Actions of any Number of Forces.

[—*the Space in which the Body moves is carried, &c.*]

When we do not attend to this, we often make very wrong Judgments concerning Motion and Percussion, mistaking an oblique Stroke often for a perpendicular one, and a perpendicular for an oblique one, and drawing false Consequences from *Phænomena* on account of those Mistakes.

Pl. 28. F. 2. This will more evidently appear by a Scheme. (See Plate 28. Figure 2.)

GHBA and ABDC are two equal Squares or Parallelograms standing upon

upon another, and separated from each other only by the Plane or Line AE ; Annotat. and there is a Body moveable upon the Line GA from G to A . First, *Lect. V.* let us suppose the Body to fall * from G to A in this Situation of the Parallelograms; and the Parallelogram $ABDC$ will receive the Blow of the falling Body at A , the Stroke being made in a Direction perpendicular to AE . Then let the Parallelogram $GHBA$ be set before the other, so that the Line GA may be directly before AC , and the Line HB before BD (or, to consider it more mathematically, let those Lines respectively co-incide) then the Body falling from G to A will also describe the Line AC perpendicular to AE . Thirdly, the Parallelograms being still before each other, let the Body fall again along the Line GA now before AC ; but at the very Beginning of the Fall, let the Parallelogram $GHAB$ (which has the Body) be moved horizontally in the Direction AE , so as to come into the Situation $BEFD$, when the falling Body has mov'd its Length GA , whereby it will be got to D instead of C , the Parallelogram $ABDC$ being all this while immovable. By this means the Body, carried in its Motion by its own Parallelogram, will have described on the Parallelogram $ABDC$ the Diagonal AD , whilst it has describ'd on its own Parallelogram only the Line GA , now represented by BD . So that a Person, suppos'd to stand in the Parallelogram $GHBA$ when it is before $ABDC$, and not to be sensible of the Motion of the Parallelogram in which he is carried, will suppose the Motion of the Body to have been *absolutely* in the Line BD perpendicular to AE , when it was only *relatively* so, but *absolutely* in the Line AD oblique to AE , as we have shewn in the Lecture. † Fourthly, let us suppose the Parallelogram † No. 3. $GHAB$ to stand upon the other $ABDC$, and to be mov'd horizontally on the Line AE , whilst the Body is falling from C to A , so as to be in the Position $HIEB$, when the Body is come down to A , which is now transferr'd to B . A Person, who does not attend to this lateral Motion, will imagine the Stroke at B to be made in the Direction HB perpendicular to AE ; and the more so, because the Force of the Stroke will be no greater than if the Body had mov'd in the Perpendicular HB , on account of its Obliquity. For tho' by the Motion of the Parallelogram, which made the Body go in the Diagonal GB , its Velocity (consequently its *Momentum*) is increas'd in the Ratio of GA to GB ; yet the Force of the Stroke, on Account of the Obliquity, must again be diminish'd in the Ratio of GB to GA , or of the *Radius* to the *right Sine* of the Angle of Inclination, as we have shewn in the Case of oblique Traction.*

Fifthly, Let us suppose the Parallelogram $GHBA$ upon the other Parallelogram $ABDC$, as we suppos'd at first; but the Body to move down the Line HB . In that Case the Body will strike the Parallelogram $ABCD$ at B in the Direction HB perpendicular to AE ; and if the upper Parallelogram be set before the lower one, the Line BD and BH

* Tho' we make use of the Word (to fall) we must suppose the Body to come down by an equal, not accelerated Motion, to render the Explanation the easier; tho' it is equally true, when we consider the Motion as accelerated.

Annotat. will be describ'd at the same Time by a Motion of the Body perpendicular to A E.

Pl. 28. F. 2. Sixthly, Let the Parallelogram G H A B be fix'd before the Parallelogram A B D C, which we will suppose to move behind it whilst the Body goes down from H to B, which Points are now in the Position of B and D; so that the Body may be come down its whole Fall to D, when the Parallelogram A B D C is come to the Position B E F D; the Point D being come to F, and the Point C to D, where it meets the Body, instead of the Point D, to which the Body came before. In this Case, a Man carried along in the Parallelogram A B D C, if he was not sensible of his own Motion, would imagine the Body to have really describ'd the Diagonal E D oblique to A E, which was only *relatively* describ'd; for the Body *absolutely* describ'd no other Line, but B D perpendicular to A E. For when the Parallelogram A B D C is come into the Position *a b d c*, the Body will have descended from B to *i*, which is a Point common to the Diagonal *b c*, and the perpendicular Line of Descent B D, and consequently seem to have describ'd *b i*, when it has only describ'd B *i*: then, when the moving Parallelogram is come into the Position $\alpha \beta \delta \kappa$, the Body will be at the Point *2*, common to the Diagonal $\beta \kappa$, and the Line of Descent B D, seeming to have describ'd $\beta 2$, when it has only describ'd B *2*: and when the Parallelogram A B D C is come to the Position B E F D, a Person carried in it, and not sensible of his own Motion, will imagine the Body to have describ'd the whole Diagonal E D oblique to A E, when it has really describ'd the Line B D perpendicular to A E.

Lastly, let us suppose the Parallelogram G H B A immoveable upon A B D C, which last is to move horizontally the Length B E, whilst the Body falls from H to B. In this Case, the Stroke will be receiv'd by the Parallelogram A B D C at the Point A instead of B, the Point A being now come to the Place of B just as the Body comes to it. The Stroke will also seem oblique as if it had come in the Diagonal I B, and the more so, because its Force will be less than if the Stroke had been made at the Point B of the moving Parallelogram, in the Ratio of I B to H B, that is, of the *Radius* to the *right Sine* of the Angle in which the Parallelogram A B D C by its lateral Motion receives the Stroke.

N. B. The Difference betwixt this Case and the 4th is this: That tho' the Obliquity of I B is the same as that of G B, yet the Force of the Stroke is greater in that Case than in this. For in that, the lateral Motion of the Parallelogram, that carries the Body, is a second Force added, the Combination of which Forces do really carry the Body in the Diagonal and increase its *Momentum*, whose Effect is only diminish'd on account of the Obliquity of the Angle G B A: but in this last Case, the Body describing no more than the Perpendicular H B, has no Increase of *Momentum*, the lateral Motion of the Parallelogram A B D C adding no Force to the Body, but on the contrary lessening the Effect of its Force by receiving

ceiving the Stroke of it obliquely, and that in the Proportion of I B to H B. Annotat. Lect. V.

If the Parallelogram A B D C had mov'd slower; as for Example, if it had only been in the Position *a b d c*, when the Body had run down its whole Length H B, it would have receiv'd the Stroke on its Point α now at B, and the Obliquity would have appear'd less, the Direction of the Motion seeming now to be in the Line K B, which also would have taken away less from the Force of the Stroke. But if A B D C had mov'd swifter than we first suppos'd; then the Obliquity of Motion would have seem'd greater in Proportion to the Obliquity of the Stroke, whose Force would have been more diminish'd. Pl. 28. F. 2.

Many are the Uses of the Truth which we have been explaining, not only in Mechanics, but all mix'd Mathematicks. I'll give one Instance of it in Opticks, by shewing how the annual Motion of the Earth about the Sun is a certain Consequence of some Observations on the fix'd Stars, made by that curious and accurate Astronomer the Rev. Mr. *James Bradley*, *Savilian* Professor of Astronomy in the University of Oxford. Those that understand Astronomy and spherical Trigonometry, may read his own Account of his Observations, and the Consequences he draws from them, in the *Philosophical Transactions*, No. 406. But as the Principles hitherto explain'd in this 1st Volume of my Course, do not qualify the Reader for understanding Mr. *Bradley's* Account; I shall just explain so much of it as will shew the Application of what we have said of oblique Motion, and at the same Time put that important Discovery of the Motion of the Earth out of Doubt.

It is proper, in order to be the better understood, that I should make a Digression here concerning the Opinions of Philosophers about the Motion or Stability of the Earth, and the Methods which have been us'd for the settling that Point.

Tho' the first Notion, that the earliest Observers of the Heavenly Motions, after discovering the Roundness of the Earth, must (in all probability) have, is, that the Sky with the Sun, Moon, and Stars, turn round the Earth, as a fix'd Center; yet some Philosophers very early taught, that the Sun was in the Center of the System, and that the Earth and all the Planets revolv'd about the Sun; as *Pythagoras*, *Philolaus* and others. But *Ptolemy* and his Followers, maintain'd the different Opinion, which agrees with the common Notion of the Vulgar, who imagine the Earth a vast Body in comparison of the Sun and Planets, and immoveable, as their Senses seem to inform them. But as Astronomers came to observe the Motions of the Planets, whose apparent Irregularity could not be accounted for upon the Supposition of the Stability of the Earth, and the Sun was discover'd to be vastly bigger than the Earth; likewise that the Diurnal Motion of the Earth about its Axis, would solve all the *Phænomena* of Day and Night, and all the other Motions ascrib'd to the *primum Mobile*, as the annual Motion solves all the *Phænomena* of the Seasons, and clears up all the Difficulties relating to the Stations and Retrogradations of the

Annotat. Planets; most People receive the Opinion of the Earth's Motion reviv'd by
 Lect. V. *Copernicus*, as the most reasonable Supposition; but still, without a Demon-
 stration, this could be but a Supposition; tho' the Use of Telescopes, inven-
 ted in our latter Ages, very much increased the Probability of the *Copernican*
Hypothesis, by shewing by the Phases of Mercury and Venus, that they must
 indubitably move about the Sun, and not the Earth. Philosophers therefore
 endeavour'd to make such Observations as might give a Demonstration one
 Way or other; and this was chiefly aim'd at by endeavouring to find an
 annual *Parallax* in the fix'd Stars; the Manner of which will be easily
 Pl. 28. F. 3. understood by the 3d Figure of Plate 28.

E O *e* represents the Orbit of the Earth, whose longest Diameter is
 E *e*, S a fix'd Star, which an Observer at E sees by the visual Ray S E,
 which is at right Angles with the Diameter E *e*, and with the Plane of
 the Ecliptick E O *e*: Let us suppose this Observation to be made in *June*;
 then half a Year after, when the Earth and Observer are gone from E
 to *e*, all the Distance E *e* from the Point E where the visual Ray S E is
 perpendicular, if the same Star be again observ'd, it will no more be seen
 by a perpendicular Ray as *s e*, but by the oblique Ray S *e*, the Angle
 from the Perpendicular being S *e* s, which is equal to E S *e*, the Angle
 under which the Diameter of the Earth's Orbit would be seen if the Ob-
 server was at the Star S; and this is called the *parallactick Angle*. It is
 evident, that if the Star was as far again from the Earth, as at Σ in-
 stead of S, the Star would be seen by the Ray Σ *e*, and the *parallactick*
 Angle would be as little again; and therefore, that the farther the Star
 is in Proportion to the Diameter E *e*, the less will the *parallactick Angle*
 be: So that if the Star be at an immense Distance from the Earth, the
 Diameter E *e* seen from the Star (that is, in respect to the Distance of
 the Star) is but a Point; therefore the *parallactick Angle* vanishes. Now
 the Fact is really so; for when the Earth is come to *e*, the Star being at
 an immense Distance is seen by the Ray *s e*; which, tho' it makes an
 Angle with the Ray S E, yet as to Sense it seems parallel to it. This
 has made the *Ptolemaicks* triumph in their Opinion, and say—that the
 Orbit E O *e* is only imaginary, and that the Earth never stirs from the
 Place E; because the Star appears always in the same Place, the right
 Angle which S E makes with the Plane of the Ecliptick (describ'd by
 the Sun, and not the Earth) never changing. The *Copernicans* have always
 answer'd, that the want of Instruments sufficiently exact has been the
 Reason, that the Angle of the *Parallax* could not be observ'd, and accord-
 ingly have endeavour'd to fix the most accurate Instruments in the best
 manner for that Purpose. Mr. *Flamsteed* fix'd an Arc of a Circle to a
 Wall, in order to observe a *Parallax* in the Pole Star, which he thought
 to be of 40 or 45 Seconds; but that Observation could not be depended
 upon, because it is not probable, that the Wall, to which he fix'd his In-
 strument, should remain exactly in the same Position in all its Parts, from
 one Solstice to the other. Dr. *Hook* pretended to have found a sensible
Parallax in the fix'd Stars, observing it in the bright Star in the Head of

Draco,

Draco, by a better Method, not depending upon a fix'd Instrument; but with a Telescope of 36 Foot, which could easily be rectified for that Purpose; but his Observations, which had been look'd upon as very accurate, were erroneous, as has appear'd by Mr. *Bradley's* which we are going to mention. Sir *Isaac Newton* was the first that demonstrat'd the Motion of the Earth; but he did it from the Laws of Gravity, and not from astronomical Observations. He was very cautious of giving Credit to such Observations as others deduc'd the Parallax of the fix'd Stars from; and when Mr. *John Rowley* had ground an Object-Glass to fix in one of the Towers of *St. Paul's Church*, at the Top of the Newel of the Stair-Cafe, which would serve as a very long Telescope to observe the Stars near the Zenith, in order to determine their Parallax, he forbade the Glass to be put up, lest Conclusions should be drawn in so material a Point from precarious Observations, looking upon them to be such; because the Building might in settling alter the Situation of this Telescopick Tower. Another Way is mention'd by Dr. *Gregory* in the 9th Section of the 3d Book of his *Astronomy*, which is this: Suppose the two Stars S and Σ^* to be in a Line, so that an Observer in the same Line at E , mistakes them for one Star S ; it is certain, that if the Observer be removed to e , or any sensible Distance from E , the Star will appear double, because now the two must subtend the Angle $Se\Sigma$; and such an Observation does not depend upon the fixing of an Instrument to an immoveable Point, or much Apparatus; a Telescope arm'd with a Micrometer, being sufficient for that Purpose. There was Hope of making an Advantage of such an Observation, because Mr. *Cassini* had observ'd several Stars appearing double or triple at one Time, and single at another; as the first of *Aries*, that in the Head of the first of the *Gemini*, that in *Orion's* Sword, and some of the *Pleiades*; but upon further Examination they did not appear single and double at fix Month's Distance, and in a Manner agreeable to what must be expected from the Parallax; but the Phenomenon of the Stars appearing single or double was owing to some other Cause: And the Distance of the fix'd Stars is so very great, that even the Removal of the Earth from its Place 160 Millions of Miles, is not able to make us discover Stars that are behind one another.

*Pl. 28. F. 3.

As a great many learned Men did not understand Sir *Isaac Newton's Principia*, and therefore did not acquiesce in his Proof of the Earth's Motion deduc'd from the Laws of Gravity; and those that did understand his Book, were yet willing to have a collateral Proof from astronomical Observations; several Persons have since attempted to observe a Parallax in the fix'd Stars; and in the Year 1725, the late honourable *Samuel Molyneux* Esq; with an Instrument, made and contriv'd by that curious Member of the *Royal Society*, Mr. *George Graham* (so much more exact than Dr. *Hook's*, that without it we should still remain in great Uncertainty as to the Parallax of the fix'd Stars) began to observe the bright Star in the Head of *Draco* (mark'd γ by *Bayer*) as it pass'd near the Zenith. Mr. *Bradley* also observ'd it along with him; and from many Observations


Annotat. vations made with great Care, it appear'd, that the Star was more north-
 Lect. V. erly 39 Seconds of a Degree in *September* than in *March*, just the contrary
 Way to what it ought to be from the annual Parallax of the fix'd Stars :
 That is, the Observers, who in *September* saw the Star at S in the Line
 ES, did in the *March* following, instead of seeing it at S in the Line eS,
 see it at σ in the Line e σ . This strange Appearance puzzled the Obser-
 vers, and Mr. *Molyneux* died before the Cause of it was discover'd. Mr.
Bradley with another Instrument of the same kind, made also by Mr. *Gra-*
ham, observ'd the same Appearances, not only in that, but several other
 Stars, and being by many Trials fully satisfied, that the *Phænomenon* was
 not owing to any Error in the Instrument; he consider'd what might be
 the Cause, and found that it was really owing to the Motion of the
 Earth, and the progressive Motion of Light. So that now the Motion of
 the Earth is settled by astronomical Observation. This Discovery, of so
 great Consequence to Astronomy, which has been attempted in vain for
 many Ages, and at last made in the Year 1728, is yet mention'd no-
 where, that I know of, but in the *Philosophical Transactions*, in a Letter
 to Dr. *Halley*, from the Discoverer, whose Modesty is equal to his great
 Skill in Astronomy. Had this been found out by some Foreigners, we
 should have had whole Volumes written about it in a pompous Manner,
 before this Time. It is surprizing, that in the Time of five Years, since
 the Publication of it, none of the learned Gentlemen abroad, have taken
 any Notice of it, either to object against, or acquiesce in it.

This shews us, that the Parallax of the fix'd Stars is much smaller,
 than has been hitherto suppos'd by those, who have pretended to deduce
 it from their Observations; for instead of many Seconds, it does not a-
 mount to one. Hence follows, that the above-mentioned Star in *Draco's*
 Head, is above 400000 times farther from us than the Sun; that Light
 moves as far from the Sun to the Earth in 8 Minutes and 13 Seconds,
 and that the Motion of the Earth in its annual Orbit, is slower than
 that of Light 10210 times; tho' the Earth moves at the rate of about
 56000 Miles in an Hour.

Tho' Mr. *Bradley's* Letter is very clear to Astronomers; yet for the
 sake of those, who have but a superficial Knowledge of the Celestial Sci-
 ence, I will (according to my Promise) endeavour to shew in the plainest
 Manner, how the progressive Motion of Light, compar'd with the Mo-
 tion of the Earth, makes a Star (which, after having been observ'd in the
 Zenith, is left behind by the Motion of the Earth) to appear, as if it
 had mov'd forward, contrary to what we might expect.

Plate 28. Fig. 4.

Pl. 28. F. 4. Let us suppose a Star to be at S, from which Light is propagated eve-
 ry Way in right Lines; but we will here consider only two of those
 Rays or Lines, viz. SE and Se. Ee is a Line, in which the Observer,
 together with the Earth, is carried in the annual Orbit; which Motion
 (to

(to make our Explanation the easier). we will suppose rectilinear; and al- Annotat.
 so consider the Motion of Light to be only 6 times swifter than the Mo- Lect. V.
 tion of the Earth; so that a Particle of Light shall go through the Line  AE or Se, in the same time that the Earth goes through the Line Ee. Pl. 28. F. 4.
 It will then follow, that whilst a Particle of Light goes through any
 Part of the Line Se, as S 1, 1 2, 2 3, &c. the Earth will go through a
 correspondent Part of the Line Ee, as Ef, fg, gb, &c. Now if the
 Earth was at Rest at E or e, an Observer from it would see the Star in
 the Line ES, or Se, by means of the Particle of Light which should
 come from S to E, or from S to e; but he would never see the Star in the
 Line es, as if it was at s. But if we suppose the Earth setting out
 from E to move in the Line Ee, at the same time that two Particles of Light
 set out from the Star S, the one to move in the Line SE, and the other in
 the Line Se; ** it will follow that the Earth will have left the Point E,
 before the Particle of Light coming in the Line SE can be arriv'd at the
 Observer's Eye, which will be at e, when the above-mentioned Particle
 of Light is at E; but then the other Particle, which came in the Line
 Se, will come to the Observer's Eye, just as the Earth comes to e, and
 strike it in the Direction se, as if the said Particle had come from s; be-
 cause, tho' it really comes in the Line Se, perpendicular to Ee, the
 Motion of the Observer in the Line Ee, makes him receive a perpendi-
 cular Stroke, as if it really was oblique; as we have shewn in the last Pl. 28. F. 2.
 Case of Fig. 2. (Page 382.) For if we suppose the Parallelogram ASeE
 with the Earth at the Angle E to move laterally in the Direction Ee,
 whilst the Particle of Light coming from the Star, moves in the Line Pl. 28. F. 4.
 Se, with such a Velocity as to go through S 1, whilst the Earth goes
 through Ef; 1 2, whilst the Earth goes through fg; 2 3, whilst the
 Earth goes through gb; 3 4, whilst the Earth goes through hi; 4 5, whilst
 the Earth goes through ik; and lastly, 5 6, whilst the Earth goes through
 ke, or what is the same, to have 6 times the Velocity of the Earth;
 then the Diagonal SE being carried laterally, when the Particle of Light
 is come to 1, and the Earth (carrying the Parallelogram ASeE) to f,
 the Point a in the Diagonal will co-incide with the Point 1, in the Ray
 Se, or Line of Motion of the Particle of Light, which by that Means
 will seem to have mov'd in the Line Sa; when the Earth is at g, b in
 the Diagonal will co-incide with 2 in the Perpendicular, and the Particle
 seem to have describ'd ab; when the Earth is at h, c will co-incide with
 3; when the Earth is at i, d will co-incide with (or be brought to) 4;
 when the Earth is come to k, d will co-incide with 5; and lastly, when
 the Earth is at e, E co-incides with 6, and the Particle of Light seems to
 have describ'd the Ray or Diagonal SE, which must be now transfer'd
 to its Parallel se, because the Earth is now remov'd to e; so that the Star
 which becomes visible to the Observer by the Line or Ray Se, does (on
 account of his Motion in the Line Ee) appear to him to come in the

** Because of the immense Distance of the Star, we must consider the Line SE, as equal
 to Se.

Annotat. Line se ; whereas if the Observer's Place had been immoveable, he must
 Lect. V. have seen the Star at S , and not at s . If then the Observer, as the
 Earth carries him along, looks through a Telescope, whose Axis makes
 Pl. 28. F. 4. the Angle SEe with the Line of the Observer's Motion Ee , when he
 comes to e , he will see the Star at s , which he could not do, if his Te-
 lescope was directed in the Line EA or eS perpendicular to Ee .

N. B. Since the Line Se is really 10210 times longer than the Line Ee ,
 because Light moves so many times faster than the Earth, how small
 must the Angle $AES (=Se s)$ be? And how nice the Instrument to
 observe such an Angle? Yet we find by repeated Trials, that the In-
 strument does perform it; which shews Mr. Graham's great Accuracy,
 which will be remember'd as long as Mr. Bradley's great Sagacity in
 drawing the Consequence.

2. [7. Page 308. Those Parts of the Vortex, which are nearest to the
 Sun, will move fastest, &c.—The Squares of the Periodical Times, &c. are
 as the Cubes of their Distances, &c.] Sir Isaac Newton has in the 9th
 Section of the second Book of his *Principia*, consider'd all possible Cases
 of Vortices, of which I have only taken that, which is mentioned in the
 11th Corollary of Prop. 52, as the most obvious. I refer the Curious to
 that Section; which would be too long to insert here at full Length; but
 the latter Part of it (*viz.* the 53d Proposition, which is applied to the Car-
 tesian Hypothesis to shew its Inconsistency, making it appear, that it is
 impossible for the Planets to move in Vortices) I thought proper to trans-
 cribe here; as well as Part of his general Scholium, at the End of his
 Book, which is to the same Purpose.

PROPOSITION 53d of Sir Isaac Newton's Principia, BOOK 3d.

*Bodies carried about in a Vortex, and returning in the same Orb, are of
 the same Density with the Vortex, and are moved according to the same
 Law with the Parts of the Vortex, as to Velocity and Direction of Mo-
 tion.*

For if any small Part of the Vortex, whose Particles, or physical Points,
 preserve a given Situation among each other, be suppos'd to be congeal'd;
 this Particle will move according to the same Law as before, since no
 Change is made either in its Density, *Vis insita*, or Figure. And again,
 if a congealed or solid Part of the Vortex be of the same Density with
 the rest of the Vortex, and be resolv'd into a Fluid, this will move accord-
 ing to the same Law as before; except in so far as its Particles now be-
 come fluid may be moved among themselves. Neglect therefore the Mo-
 tion of the Particles among themselves, as not at all concerning the pro-
 gressive Motion of the whole, and the Motion of the whole will be the
 same as before. But this Motion will be the same with the Motion of
 other

other Parts of the *Vortex* at equal Distances from the Center; because the Solid, now resolv'd into a Fluid, is become perfectly like to the other Parts of the *Vortex*. Therefore a Solid, if it be of the same Density with the Matter of the *Vortex*, will move with the same Motion as the Parts thereof, being relatively at Rest in the Matter that surrounds it. If it be more dense, it will endeavour more than before to recede from the Center; and therefore overcoming that Force of the *Vortex*, by which being, as it were, kept in *Æquilibrio*, it was retain'd in its Orbit, it will recede from the Center, and in its Revolution describe a Spiral, returning no longer in the same Orbit. And by the same Argument, if it be more rare, it will approach to the Center. Therefore it can never continually go round into the same Orbit, unless it be of the same Density with the Fluid. But we have shewn in that Case, that it would revolve, according to the same Law, with those Parts of the Fluid, that are at the same or equal Distances from the Center of the *Vortex*. Annotat. Lect. V.

Cor. 1. Therefore a Solid revolving in a *Vortex*, and continually going round in the same Orbit, is relatively quiescent in the Fluid, that carries it.

Cor. 2. And if the *Vortex* be of an uniform Density, the same Body may revolve at any Distance from the Center of the *Vortex*.

S C H O L I U M.

Hence it is manifest, that the Planets are not carried round in corporeal *Vortices*. For, according to the *Copernican Hypothesis*, the Planets going round the Sun, revolve in Ellipses, having the Sun in their common *Focus*; and by *Radii* drawn to the Sun, describe Areas proportional to the Times. But now the Parts of a *Vortex* can never revolve with such a Motion. Let AD, BE, CF (*Pl. 28. Fig. 5.) represent three Orbits described about the Sun S, of which, let the utmost Circle CF be concentric to the Sun; and let the *Aphelia* of the two innermost be A, B; and their *Perihelia* be DE; therefore a Body revolving in the Orb CF, describing by a *Radius* drawn to the Sun, Areas proportional to the Times, will move with an uniform Motion. And, according to the Laws of Astronomy, the Body revolving in the Orb BE, will move slower in its *Aphelion* B, and swifter in its *Perihelion* E; whereas according to the Laws of Mechanics, the Matter of the *Vortex* ought to move more swiftly in the narrow Space between A and C, than in the wide Space between D and F, that is, more swiftly in the *Aphelion*, than in the *Perihelion*. Now these two Conclusions contradict each other. So at the Beginning of the Sign of *Virgo*, where the *Aphelion* of Mars is at present, the Distance between the Orbits of Mars and Venus, is to the Distance between the same Orbits at the Beginning of the Sign of *Pisces*, as about 3 to 2; and therefore the Matter of the *Vortex* between those Orbits ought to be swifter at the Beginning

E e e

ginning


Annotat. *ginning of Pisces*, than at the Beginning of *Virgo*, in the *Ratio* of 3 to
 Lect. V. 2. For the narrower the Space is, through which the same Quantity of
 Matter passes in the same Time of one Revolution, the greater will be
 the Velocity with which it passes through it. Therefore, if the Earth, be-
 ing relatively at Rest in this celestial Matter, should be carried round by
 it, and revolve together with it about the Sun, the Velocity of the Earth,
 at the Beginning of *Pisces*, would be to its Velocity at the Beginning of
Virgo in a *sesquialteral Ratio*. Therefore the Sun's apparent diurnal Mo-
 tion at the Beginning of *Virgo*, ought to be above 70 Minutes; and at
 the Beginning of *Pisces* less than 48 Minutes. Whereas on the contrary,
 that apparent Motion of the Sun is really greater at the Beginning of
Pisces, than at the Beginning of *Virgo*, as Experience testifies; and therefore
 the Earth is swifter at the Beginning of *Virgo* than at the Beginning of
Pisces. So that the Hypothesis of *Vortices*, is utterly irreconcilable with
 Astronomical *Phænomena*, and rather serves to perplex than explain the
 heavenly Motions.

GENERAL SCHOLIUM.

The Hypothesis of *Vortices* is press'd with many Difficulties, that eve-
 ry Planet by a *Radius* drawn to the Sun, may describe Areas propor-
 tional to the Times of Description, the periodick Times of the several Parts
 of the *Vortices* should observe the duplicate Proportion of their Distances
 from the Sun. But that the periodick Times of the Planets may obtain
 the sesquiplicate Proportion of their Distances from the Sun, the periodick
 Times of the Parts of the *Vortex* ought to be in the sesquiplicate Pro-
 portion of their Distances. That the smaller *Vortices* may maintain their
 lesser Revolutions about *Saturn*, *Jupiter*, and other Planets, and swim
 quietly and undisturb'd in the greater *Vortex* of the Sun, the periodick
 Times of the Parts of the Sun's *Vortex* should be equal. But the Rota-
 tion of the Sun and Planets about their Axes, which ought to corre-
 spond with the Motions of their *Vortices*, recede far from all these Pro-
 portions. The Motions of the Comets are exceeding regular, and go-
 vern'd by the same Laws with the Motions of the Planets, and can by
 no means be accounted for by the Hypothesis of *Vortices*. For Comets
 are carry'd with very excentric Motions through all Parts of the Hea-
 vens indifferently, with a Freedom that is incompatible with the Notion
 of a *Vortex*.

3. [9. Page 314. *There are many more Experiments relating to central Forces, &c.—for the sake of the Curious we shall mention a few more in the Notes.*] To shew experimentally how a Planet accelerates its Motion as it approaches to the Sun, and retards, as it recedes from the Sun; fix the whirling Table (in the Condition represented by the 11th Figure of

*Pl. 24. F. 11. Plate 24. *) to its Frame, that it may be easily turn'd round in its hori-
 zontal Situation. Then take off the Square, or rectangular Piece S, and
 fasten

fasten it to the Table at the Place *Mm*, by means of a Screw in the Annotat. under Edge of it, which goes through the Table, and is fix'd by a Nut Lect. V. under the Table. Having so done, instead of the two Balls *T, M*, take  one of them as *M*, and having plac'd it within an Inch of the short Part ^{Pl. 28. F. 11.} of the *Square* at *M*, pass its String through one of the Side-Holes of the Center-Piece *C*, and so up through the middle Hole; then having tied a little Wire *w* cross the String, so that when the centrifugal Force acts upon *M* (while the Table turns round) the Wire may hinder the Ball striking against the End of *Mm*. Let the Table be whirl'd about in the Direction of the Dart, whilst the Hand is holding the other End of the String loosely at *x*, and the Ball will press on the *Square Vv* at *M*. Pull the String a little to make the Ball approach nearer to the Center, and you will see and hear it strike on the other Side at *m*, which shews its Motion to be accelerated; but if then (the Table still going round with the same Velocity) you slack your Hand, the Ball in receding from the Center will strike against *M*, which shews it to be retarded, because it moves slower than the Parts of the Table under it, as it mov'd faster than the Table in the other Case.

The Conclusion from the 15th Experiment (*viz.* that in the same Orbit the centrifugal Force is as the Square of the Velocity) would follow from any other Variation of the periodical Times; as for Example, if the String of the Pulley *H* * be put on the Groove of 3 Inches, and that of the Pulley *K* on the Groove of 2 Inches, each Planet being of equal Weight, and at the same Distance from the Center; the swiftest Planet *P* will (in its Tower) raise a Weight of $2\frac{1}{4}$ Pounds, whilst the slowest *p* raises only 1 Pound; which Weights are to one another as 9 and 4; that is, as the Squares of the periodical Times 2 and 3 reciprocally, or the Squares of the Velocities directly. ^{* Pl. 25. F. 1.}

In a word, let the *Central Forces* differ in any manner whatsoever, one may compare them together by the Help of what has been already explained; for *they are always in a Ratio compounded, of the Ratio of the Quantity of Matter in the revolving Bodies, of the Ratio of the Distances from the Center, and lastly of the inverse Ratio of the Squares of the periodical Times.* And the Comparison is thus made. Multiply the Quantity of Matter in each Body by its Distance from the Center, and divide the Product by the Square of the periodical Time, and the Quotients of the Divisions will be in the said compound *Ratio*, that is, as the central Forces.

EXPERIMENT. Plate 25. Fig. 1.

Let the Wheel-string be on the 2 Inch Groove of the Pulley *K*, and ^{Pl. 25. F. 1.} the 3 Inch Groove of the Pulley *H*, which will make the periodical Times of the Planets *P* and *p* be to one another as 2 to 3², by the Addition of 4 Ounces to the Weight-bearing Piece in the Tower *S*, the whole Weight will be of 6 Ounces, and let the Planet *P* be of 2 Ounces, and at 8 Inches

Annotat. from the Center. The Planet p must be of 4 Ounces at 12 Inches from
 Lect. V. the Center, and the Weight that draws it in the Tower s , must be of
 8 Ounces. Turn the Wheel, and both Weights will rise at the same
 Pl. 25. F. 1. moment.

Multiply 2 (the Weight of the Planet P) by 8 in Distance, and the Product is 16, which being divided by 4 (the Square of its periodical Time) gives 4 for the Quotient (for $2 \times 8 = 16$, and $\frac{16}{4} = 4$.) Then multiply 4 (the Weight of the Planet p) by its Distance 12, and the Product 48 being divided by 9 (the Square of the periodical Time of p) you will have $5\frac{1}{3}$ (for $4 \times 12 = 48$, and $\frac{48}{9} = 5\frac{1}{3}$.) Now these Numbers 4 and $5\frac{1}{3}$ are to one another as 12 and 16, or as 6 and 8, which are the Weights in the Towers that express the Quantities of the central Forces.

Whenever the Quantities of Matter are equal, you need only divide the Distances by the Squares of the periodical Times to determine the Proportion of the central Forces to one another.

And in that Case, if the Squares of the periodical Times are to one another, as the Cubes of the Distances, the Quotients of the Divisions will be in an inverse Ratio of the Squares of the Distances; and you will find the central Forces to be in the same Ratio; which is the Case of the Planets and Comets in the Heavens.

This may be very well illustrated by the following

EXPERIMENT. Pl. 25. Fig. 1.

Let the Wheel-string go over the 3 Inch Groove of the Pully K , and the 6 Inch Groove of the Pully H , to make the periodick Time of the Planet P be 3, while that of p is 6. Let P and p be of 4 Ounces each, P being 5 Inches from the Center, and p at the Distance of 8 Inches from the Center. Then if P is fasten'd to 10 Ounces in its Tower S , and p to 4 Ounces in its Tower s , both Weights will rise at once when the Wheel is turn'd about.

The central Forces are here as 4 to 16, and the inverse Ratio of the Squares of the Distances is the Ratio of 25 to 64 (which last Number should strictly be 62 $\frac{1}{2}$, and may be so if the Distance from the Center be made the square Root of 62 $\frac{1}{2}$, but we choose 8 Inches to avoid Fractions) near enough for an Experiment. The Squares of the periodical Times are 9 and 36, which also agree near enough with the Cubes of the Distances, viz. 125 and 512; for 12 would become only 100, the exact Ratio wanted, or much nearer, if the Distance from the Center had been the square Root of 62 $\frac{1}{2}$, as we said before.

4. [16. Page 328. Others, &c. alledge, that the Momentum is not as the Product of the Mass by the Velocity of the moving Body, but as the Mass multiplied into the Square of the Velocity.

[24. Page

[24. Page 336. *The Weight, &c. multiply it by the Velocity, it will give us the Momentum, &c.* Annotat. Lect. V.

Till within these few Years, all Mathematicians and Philosophers agreed in the Opinion, that the *Momentum*, or moving Forces of Bodies, is made up of the Mass, or Quantity of Matter, multiplied into the Velocity, as we have explain'd in the 2d Lecture.* Monf. *Leibnitz* (if I am rightly inform'd) was the first that rejected the old Opinion, affirming, that the Force of Bodies in Motion, was made up of the Mass multiplied by the Square of the Velocity, applying it to the Fall of Bodies, and saying, that the Stroke made by falling Bodies was always proportionable to the Heights from which they fell, which Heights † are as the Squares of the Velocities. But his Error lay, in not considering the Time (as we have shewn) for the Velocities alone are not the Cause of the Spaces describ'd, but the Times and the Velocities together; for otherwise the Effect would be greater than the Cause, which is absurd. Several ingenious Men have endeavour'd to defend Monf. *Leibnitz's* Opinion by subtle, tho' fallacious, Arguments, having first deceiv'd themselves by misapplied Observations and Experiments.

Some have distinguish'd the Actions exerted on Bodies (as for Example, the Force of Gravity) into a *living* and a *dead* Force (*vis viva*, & *vis mortua*) calling that a *living Force*, whereby it produces a visible Effect on a Body, and a *dead Force*, that which is destroy'd by a contrary Cause; as when a Body is kept from falling by an Obstacle, or being in the Scale of a Balance is kept from descending by a Counterpoise in the opposite Scale. Were we to allow of this Distinction, yet the common Experiment of the most simple mechanical Organ, I mean the Balance, shews that the *vis viva* and the *vis mortua*, are both in the simple Proportion of the Velocity multiplied into the Mass. For Example, 4 Pounds being plac'd at the Distance of 6 Inches from the Center of Motion, and 2 Pounds at the Distance of 12, will have a *vis viva*, if the Balance be put into a swinging Motion. Now these Forces appear to be equal, because, with contrary Directions, they soon destroy one another; but they are to one another in the simple Ratio of the Velocity multiplied into the Mass, viz. $4 \times 6 = 24$, and $2 \times 12 = 24$. Whereas if the Forces in this Case had been as the Mass into the Square of the Velocity, the Weight 2 plac'd at near 8 Inches and an half, would have destroy'd the Motion of the Weight 4 at 6 Inches Distance, and have reduc'd it to an *Æquilibrium*. That is, $6 \times 6 \times 4 = 144$, and $8,426 \times 8,426 \times 2 = 144$. The Case is the same in the *vis mortua*; for then 2 at the Distance 12, keeps 4 in *Æquilibrio* at the Distance 6, and the least Alteration either of Weights or Distances, will destroy the *Æquilibrium*. The same thing is true in the *Leaver*, *Pulley*, *Axis in Peritrochio*, *Inclin'd Plane*, *Wedge*, and *Screw*.

The *Momentum* of Bodies appearing so plainly by this common Experiment to be the Product of the Mass by the Velocity, others have said, that the particular Make of the mechanical Organs was the Cause of this Phenomenon,

Annotat. *Phænomenon*, and that this Action of Bodies upon one another, by means
 Lect. V. of Instruments must be call'd *Pressure*, distinguishing between *Force* and
Pressure, allowing *Pressures* of Powers to be to one another reciprocally as
 the Masses multiplied into the Velocities; but denying that Forces are
 in that *Ratio*. Now to shew, that Forces, which we also call the *Mo-*
menta of Bodies, are in the *Ratio* above-mentioned, let us impartially
 consider the following Experiment communicated first to me by Mr. George
 Graham.

EXPERIMENT. Plate 24. Fig. 1.

Pl. 24. F. 1. The Machine represented by the Figure has been already describ'd
 (Page 291.) only we must suppose the Arc F E to be divided into 24
 Degrees on each Side, counting from the Edges A D and C B of the flat
 square Pendulum, instead of 18 mark'd in the Figure; and that the Pen-
 dulum weighs 2 Pounds, and the Weight W also 2 Pounds.

Now, if the Pendulum without the Weight W be drawn up to the
 24th Degree on the Side of E, when you let it go, it will rise to 24 on the
 other Side; but if a Person, who holds the String L in his Hand (whilst
 the Weight W hangs half an Inch above A B C D, when at the lowest
 Place) lets go the String just at the moment that the flat Pendulum
 comes to the lowest Place, the Pendulum receiving thereby an Addition
 of Matter equal to its own, goes only to 12 on the Side of F. If we
 let the Pendulum fall from 20 Degrees, and W be let down upon it at
 the lowest Part of the Vibration, then it will (thus loaded) go but to 10
 Degrees; and when you let it go from 12, if it receives the Weight W
 at Bottom, it will go but to 6.

Now to compare together the two Opinions, we must examine, which of
 them gives the *Momentum* the same before and after the Reception of W.

OLD OPINION.

| In the first Half of the Vibration. | | | In the last Half of the Vibration. | | |
|-------------------------------------|------|---------|------------------------------------|------|---------|
| Velocities | Mass | Momenta | Velocities | Mass | Momenta |
| 24 | 2 | 48 | 12 | 4 | 48 |
| 20 | | 40 | 10 | | 40 |
| 12 | | 24 | 6 | | 24 |

NEW OPINION.

| In the first Half of the Vibration. | | | | In the last Half of the Vibration. | | | |
|-------------------------------------|----------|------|---------|------------------------------------|----------|------|---------|
| Vel. | Sq. Vel. | Mafs | Momenta | Vel. | Sq. Vel. | Mafs | Momenta |
| 24 | 576 | 2 | 1152 | 12 | 144 | 4 | 576 |
| 20 | 400 | | 800 | 10 | 100 | | 400 |
| 12 | 144 | | 288 | 6 | 36 | | 144 |
| | | | | instead of | | | |
| | | | | 1152 | | | |
| | | | | 800 | | | |
| | | | | 288 | | | |

But

But to have the *Momenta* as they ought to be according to the Annotat. new Opinion, the Velocities, after the Reception of W on the Pendulum, Lect. V. ought to have been $17-$, $14,1+$, and $8,48+$; for $17- \times 17- = 289$, and $14,1+ \times 14,1+ = 200$, and $8,48+ \times 8,48+ = 72$; which Numbers respectively multiplied by the Mass of the Pendulum and Weight, will give for Products 1152, 800, and 288, the same *Momenta* that the Pendulum had in the first Half of the Vibration.

N. B. As it requires a little Practice and Dexterity to let W fall just upon the Middle of ABCD, one must make several Trials, and have a Circle drawn upon the Plate ABCD, to shew the exact middle of it; but the Experiment is conclusive, tho' W does not fall precisely into the Circle; for when it falls beyond it (for Example, when you let the Pendulum go from 12) then will the compound Pendulum go as far beyond 6 towards F, and come as far short of 6 towards E. The same happens, whatever Height the Pendulum falls from. The Experiment also succeeds equally, whether the Bottom of the Weight and upper Surface of the Plate be rough or smooth, soft or hard. Nay, there is another Way of making the Experiment, easier than letting fall the Weight, and that is done by fixing the String that holds the Weight so, that the Plate can but just slip under it in its Vibration, and then screwing the curv'd upright Piece w x w to the Plate, represented now by a b c d, in such manner, that the curv'd Piece, or Semi-cylinder, which is something bigger than the Circle w w (Base of the Weight when on) shall, at the Bottom of the Vibration, embrace the Weight W, and carrying it along, without its falling upon the Plate. In this last Case the Experiment will also succeed, whether the Weight has its cylindric Surface of Lead, Iron, Brass, or even soft Leather. I made the Experiment these different Ways, in order to remove several Objections; tho' they were of so little Force, that they may be easily answer'd without; and they are as follows.

Object. 1. The Shock made by the Fall of the Weight W, hinders the compound Pendulum from going so far as it ought to do.

Answer. The Pendulum is in an horizontal Motion, when it receives the Weight W, which by its Fall gives a perpendicular Stroke, which (by what has been said of two Forces acting at the same time upon a Body)† L. 32. can neither advance nor retard it towards the Side where it is going, but No. 85. only make it go lower, if the Strings that suspend the flat Pendulum yield a little. For Example, let e f (* Plate 24. Fig. 1.) be the Line in which * Pl. 24. F. 1. the Pendulum moves at the lowest Part of its Vibration, which we will consider as a straight Line, taking the Tangent instead of a small Arc; let the Weight W fall in the Line w e. Now, if we suppose the Strings to yield, so as to become longer by the whole Depth e g, then will the Pendulum be acted upon by two Forces, as e f and e g, which will make it go in the

Annotat. the Diagonal el ; if the Strings stretch but to b , then will the Body go
 Lect. V. in the Diagonal em ; if they stretch only to i , the Body will go in the
 Pl. 24. F. 1. Diagonal en ; and lastly, if the Strings either don't stretch at all, or by
 their Re-action restore themselves wholly, then is the whole Effect of the
 Shock of the Fall of W destroy'd, and the Body will go in the Line ef
 as if there had been no such Stroke. Now in all these Cases, the compound
 Pendulum goes from the Line eg to the Line fl in the same Time,
 and the Edge of the Plate must shew the same Number of Degrees upon
 the graduated Index below it. To these Objectors also, the Success of the
 Experiment when soft Leather is glued under the Lead, must give
 Satisfaction; but making the Experiment without letting the Weight fall
 at all (when the little Semi-cylinder $w x w$ of the Plate takes the Weight
 along with it) must be a full Conviction.

Object. 2. The Friction of the Weight W , and the Plate $ABCD$ one
 against another, occasions the Loss of Motion, and hinders the compound
 Pendulum from going to those Distances, that are agreeable to the new
 Opinion.

Answer. This Objection is too loose and general; for the Objectors
 ought to shew how much Motion Friction can destroy in other Cases, and
 that there is as much Motion lost here. Whereas, upon Examination,
 the suppos'd Cause will appear unequal to the Effect. For the Motion
 lost here, according to the new Opinion, must be half of the whole Motion
 of the compound Pendulum; whereas Friction, at most, never destroys
 above a Third. Besides, the Friction of one Part of the Pendulum
 against another, does no way affect the Motion of the whole Pendulum, or
 of the System of those two Bodies, that Way in which it is carried. For
 Sir Isaac Newton has demonstrated in his *Principia*, Book the 1st, in the
 Corollaries to the Laws of Motion—*That if Bodies, any how mov'd*
among themselves, are urg'd in the Direction of parallel Lines by equal accele-
rative Forces; they will all continue to move among themselves, after the
same Manner as if they had been urg'd by no such Forces. Now, this is
 the Case of our compound Pendulum, all whose Parts for the small
 Time of the Shock or Friction, may be consider'd, as carried along in
 right Lines; that is in the Line ef and its Parallels. The Answer by way
 of Experiment is, that when I have made the Bottom of W very smooth,
 as also the Surface of the Plate (as I was desired to do by some of the
 Objectors) it has been more difficult to make the Experiment; but the
 Success has been the same: For whenever the Weight W had its Center
 a little behind the Center of the Circle $w w$, perhaps by slipping back-
 wards, the compound Pendulum came short of the 12, 10, or 6 Degrees
 forwards towards F ; but then it exceeded as much when it came back
 towards E ; whereas the Diminution of Friction, ought to make it go be-
 yond those Numbers according to the new Principle.

Tho'

Tho' this is sufficient (at least as it appears to me) to convince any Person that is attentive to the Thing, of the Truth of the old Opinion; yet, in justice to the two ingenious Professors of *Leyden* and *Utrecht*, I ought to mention an Experiment, which they made in Opposition to this Experiment of the flat Pendulum.

EXPERIMENT. *Plate 28. Fig. 6.*

Pl. 28. F. 6.

ABC is a vertical Board, having an upright Piece *AD* fix'd to it with two Screws *yy*, at the Top of which Piece there is fix'd an horizontal Arm *DE*. From the said horizontal Arm are let down two fine Fiddle-strings *Cm*, *cn*, which by the two little Hooks *mn* sustain the Brass Cylinder *W*. This Cylinder is a Box with a Cone *H* screwing into it, to shut up any Weights contain'd in it, and at the same time to cut the Air the better, when the Cylinder moves with the conick End foremost. *tt* is a little Tail or Ruler fix'd to the other End of the Cylinder on its Axis continued, with several vertical Holes in it, to receive the Pin *P*, which goes through it behind *rr* a strong Brass Plate fix'd at right Angles to the Board *ABC* by Means of its Return *R* screw'd tight against the Board, which Plate has an Hole to transmit the Tail *tt*, whereby the Spring *S* (fasten'd also to the End of the Cylinder) is bent. When the Pin *P* is suddenly pull'd out, the Cylinder is shot forward by the unbending of the Spring, in the Line *Hh*; and the sliding Ruler *On* being drawn out, so as to have its first Division begin where the String *cn* hangs, that String in the Motion of the Cylinder will successively pass over the Divisions upon the Ruler, and shew how far the Cylinder is shot forward in every Experiment. The Cylinder, with its Tail, Spring, and Head, weighs just one Pound; but by putting different leaden Cylinders into it, it may be made to weigh 2, 3, or 4 Pounds; and likewise the Spring may have different Degrees of Force given to it, according as the Tail *tt* is drawn farther through the Plate *rr*, towards *R*, and kept in its Place by the Pin *P*. N.B. *The 7th Figure represents Pl. 28. F. 7. Part of the Cylinder with its Tail and Spring, and the rectangular Plate, drawn by a larger Scale, and mark'd with the same Letters.*

When the Spring bent to the same Degree, does, by pulling out the Pin *P*, successively shoot forward the Cylinder differently loaded; you will find, by observing the String *cn*, to what Distances the Cylinder is thrown. Now it always happens, that in those Cases, whatever the Weight of the Cylinder is, the Product of its Mass by the Square of the Velocity, is always the same. That is, if the Cylinder weighing 1, the Spring be adjusted so, as to throw it to the Degree 8 mark'd on the Ruler or Tangent Line *no*; the Cylinder loaden with 2, will by the same Bent of the Spring be thrown to 5,65, &c. if loaden with 3, it will go to 4,62, &c. and if loaden with 4, it will go to 4. These Gentlemen therefore alledge, that they have made Use of Elasticity in their Experiment, as I have made use of Gravity in mine; and that as their Experiment is agree-

Annotat. able to the new Opinion, and mine is agreeable to the old ; therefore no-
 Lect. V. thing is to be concluded from either of them.

Pl. 28. F. 6. { Indeed, one would imagine so at first ; and one would think, that (ac-
 cording to the old Opinion) the Spring which shoots the Weight 1 to 8
 Degrees, ought to shoot the Weight 4 only to 2 Degrees, whereas we
 see it shoots to 4 Degrees, since the same Force must produce the same
Momentum. But here lies the Fallacy. If the Force of the Spring acted
 no longer upon the Weight 4 than it does upon the Weight 1, it
 would shoot it but to 2 ; but the Spring acts as long again upon the
 Weight 4 as it does upon the Weight 1, as is evident, because the
 Weight 4 moves but with half the Velocity of the Weight 1, the Spring
 quitting the Plate *rr* as soon again, when it drives forward the Weight
 1. So that an equal Cause acting twice the Time, must produce the same
 Effect, as a double Cause acting during the same Time as the single one.

When a Truth has been prov'd by plain Reasoning and simple Experi-
 ments ; there is no Occasion to take any Notice of such subtle Reasonings,
 and complex Experiments, that are made use of to perplex it, for the Con-
 firmation of that Truth. But, if the Experiments made to establish one
 Opinion, and overthrow another, tho' they have not answer'd that End, yet
 lead us to some new Discovery, or clear up some other Truth not gene-
 rally known ; they certainly merit our Attention and thorough Examina-
 tion. Dr. *s'Gravesande*, and Dr. *P. Muschenbroek*, are too curious in ma-
 king, and too faithful in relating their Experiments, not to have us cre-
 dit the Facts : Therefore, tho' I deny the Conclusion, which they deduce
 from the Impressions made on Clay, or other soft Substances, by Spheres,
 Cones, or Cylinders, I think, there are very useful Consequences to be drawn
 from their Experiments in respect of Percussion, and the Resistance of soft
 Bodies. I likewise believe Professor *Polenus* to have made his Experiments
 carefully, tho' I have not the Honour to know him. Therefore I will,
 in the Beginning of my next Volume, consider this Dispute fully, and
 hope to shew the entire Fallacy of the Reasonings, which have been pub-
 lish'd in Favour of the new Opinion, and fully explain, why the Experi-
 ments alledged, are not conclusive in that Point. It is not proper to do
 it now ; because, I have not yet explain'd the Laws of the Congress, or
 Shock of Bodies, which this Volume, already swell'd to too great a Bulk,
 allows me no Room to do. I hope I shall then be able to settle the Dis-
 pute, by proving the three following Propositions.

1. *That the Cause, which accelerates Bodies in their Motion downwards,
 and retards them in their Motion upwards, does (abstracting from the
 Resistance of the Air) not accelerate them in their Fall with more
 Difficulty, or slower than it retards them in their Rise, which is a
 Supposition made in favour of the new Opinion.*

2. *That Impressions, or Pits of equal Capacities or Depths, made on soft
 Substances by hard Bodies, by striking them with unequal Velocities, are
 not*

not proportionable to the Momenta, or Forces of those percutient Bodies; tho' they be proportionable to the Masses multiplied into the Squares of the Velocities. Annotat. Lect. V.

3. That Experiments of the Congress of soft and elastick Bodies do not prove the new Opinion, but confirm the Old.

In the mean time, I refer the curious Reader for Satisfaction to the *Philosophical Transactions*, where he will find Dissertations upon this Subject, in the following Numbers. No. 371, 375, 376, 396, 400, and 401.

In No. 371. There is a Paper of Dr. Henry Pemberton, to shew the Experiments of *Polemus* inconclusive for the new Opinion.

In No. 375. I endeavour to confirm the old Opinion, by an old Experiment on the Balance, and some Experiments of the Congress of Steel Balls.

In No. 376. I endeavour to shew the Fallacy of *Polemus's* Experiments by two new ones.

In No. 396. Mr. John Eames, F.R.S. makes Remarks upon the new Opinion, relating to the Forces of moving Bodies, in the Case of Collision of *Non-elastick Bodies*: And in the same *Transaction* shews, in another Dissertation, that the Proof for the new Opinion, drawn from the Doctrine of Composition and Resolution of Forces (when a Body by its Impulse bends several Springs) equally proves both Sides of the Question, and thereby proving too much, does in reality prove nothing at all; and therefore is far from deserving the Name of a Demonstration.

In No. 400. The said Mr. Eames has made Remarks upon some Experiments in *Hydraulicks*, which seem to prove, that the Forces of equal moving Bodies, are as the Squares of their Velocities.

In No. 401. There is a Letter of the late Rev. Dr. Samuel Clarke, concerning the Proportion of *Velocity* and *Force* in *Bodies in Motion*.

Yet I cannot but blame the Doctor's uncivil Treatment of those Gentlemen, who defend the new Opinion, in Expressions very rude, and not at all relating to the Argument it self: Neither can I be of his Opinion, in his taxing Dr. s'Gravesande with "intending to raise a Dust of Opposition against Sir ISAAC NEWTON's Philosophy;" since I know that Professor does, and always did, very much esteem Sir Isaac Newton's Works; and that both he, and Dr. Petrus van Muschenbroek, Professor at *Utrecht*, do teach the *Newtonian* Philosophy, tho' they differ from him in relation to the *Momentum* of Bodies.

[25. Page 337. *The Pile may enter into the Earth farther than in that Proportion, &c.*—Page 338. *Nails, &c. give way to the Hammer nearly according to the Square of the Velocity, &c.*—But the Reverse must be done in the *Battering Ram*.

Tho' the Experiments made by the ingenious Professors *Polemus*, s'Gravesande, *Muschenbroek*, and others, do not prove the *Momenta* of Bodies, to be as the Mass into the Square of the Velocity, for the Reasons given in the

Annotat. *Philosophical Transactions*, which I have refer'd to, and for other Reasons
 Lect. V. (which I reserve for that Part of my Work, wherein I shall consider the
 Congress of Bodies, and which, I hope, will be satisfactory to those Gentlemen themselves) yet they have a great deal of Merit, because they do certainly shew, that the same Body striking soft Substances with different Velocities, does make Impressions upon them, in Proportion to the Squares of those Velocities; and thereby give us a Principle to direct the Practice of some mechanical Operations, which was not very well known before. Accordingly, a one-handed Hammer, striking upon a hot Iron, when it has 4 times the Velocity of a two-handed Hammer, which is four times heavier, will do twice the Work in forging the Iron, tho' the *Momentum* is the same in both; but the Introcession, or Yielding of the Iron, is not the same in both. So in driving Piles, when the Earth is equally yielding, if we raise the Rammer four times as high, the Pile will enter into the Earth four times easier; but when the Pile (tho' shod with Iron) strikes upon a Rock, or is driven in stony Earth, the Case is not the same, but in a mix'd Proportion between that of the Mass into the Velocity, and the Mass into the Square of the Velocity.

* Pl. 28.
 F. 8.

Now in a Battering Ram, the Effect will be greater, when a great Quantity of Matter moves with a small Velocity, than when the same *Momentum* is produc'd by a little Quantity of Matter which moves with a great Velocity. The explaining of this will give some Light into the Dispute concerning the new and the old Opinion. For, if we suppose *ACDE* * a Piece of Wall (whose horizontal Section is represented here) struck by a Cannon Ball, or a Battering Ram at *B*, the Effect in the first Case will be this, that the Cannon Ball (supposing it to have a sufficient Velocity) will go thro' the Wall, making the Hole *abBc* thro' the Wall, or at most, the Hole *dcBe*, in the Form of a truncated Cone, shaking a little more of the Wall, than the Parts of it, which are very near that Hole, because the Velocity is so great, that the Motion has not time to be communicated very far, whilst the Bullet is passing. But if a Battering Ram, with the same *Momentum*, but a Velocity so much less as it has more Matter, strikes the Wall at *B*, it will shake it round about to a great Distance, the Motion having Time to be communicated; and, if the successive Blows be continued, so that the second may be given, before the Shaking given by the first is over; after several Blows, a large Piece of Wall will be thrown down, such as *FGHI*. This is what Sir *Christopher Wren* has often practis'd in the beating down of old Buildings and Ruins; in order to erect new Edifices in the Place where old ones formerly stood: And he said, he found no Way so expeditious for demolishing old Walls, whose Mortar was almost as hard as the Stone it self.

L. 2. No. 9.
 and Ann. 11.

In War indeed, Cannon is preferable to the ancient Battering Rams, for many Reasons; some of which have been given in *Lect. 2d. No. 9.* and the 11th *Annotation*. And in those Cases we are to observe, that the Besiegers shoot obliquely, choosing rather to strike the Walls obliquely from different Batteries, tho' with less *Momentum*, than the second Ball may beat down what the first has shak'd: This being more effectual, than

than to make many small Holes in a Wall by shooting perpendicularly. Annotat. To illustrate this yet further, I will here repeat what I have already said Leſt. V.
in the *Philosophical Transactions* (No. 376.) viz. “ That when a Door
“ is half open, and moving very freely on its Hinges; if a Pistol be
“ fir’d againſt it, the Ball will go through the Door without moving
“ out of its Place; but if we take a large Weight of Lead, and throw it
“ againſt the ſame Door, with the ſame Force as the Pistol Bullet mov’d,
“ the Door will be remov’d from its Poſition, and carried out of the Place
“ on its Hinges by the Stroke; becauſe in the firſt Caſe, the Motion of
“ the Ball is communicated but to a few Parts of the Door, and in the
“ laſt it is diffuſ’d all over it. Nay, the Door will be mov’d by the
“ Stroke, even tho’ there ſhould be a prominent Part in the Lead no
“ bigger than a Pistol Bullet, in order to ſtrike the Door upon no more
“ of its Surface, than the Bullet has done.

5. [32. Page 342. *A Fluid will reſiſt ſometimes as much as a Solid, &c.*] Let us ſuppoſe a Piece of Wood, as a Plank one Inch thick, to be of the ſame ſpecifick Gravity as Water; and, that Water of the ſame thickneſs covers a Surface of Clay; and let us examine what will be the Effect of a Bullet ſtriking the Solid or the Fluid. Let us again ſuppoſe, that the Bullet ſtriking the Water with the Velocity 1, paſſes through it, overcoming by its *Momentum* the Reſiſtance as 1, which ariſes from the Denſity of the Water, and alſo another Reſiſtance as 1, which ariſes from its Tenacity. If the Bullet, with the ſame Velocity, ſtrikes the Wood, whoſe Reſiſtance from its Denſity is as 1, but whoſe Reſiſtance from the Tenacity is as 9; that is, whoſe whole Reſiſtance is 10; it is certain, that the Bullet can not penetrate through the Wood with the Velocity 1, but it muſt have the Velocity 10, to overcome the Reſiſtance above-mentioned.

Now, if the Bullet, with the ſame Velocity 10, comes againſt the Water, it will meet with a Reſiſtance, as 100, on account of the Denſity, † and alſo the Reſiſtance 10, on account of the Water’s Tenacity, * † L. 5. 32. that is, 110. Then if the Bullet with the ſame Velocity ſtrikes the * Page 340. Wood, and we ſuppoſe the Reſiſtance of the Wood, from its Quantity of † 341. Matter, to increaſe as the Square of the Velocity, as it does in the Water (tho’ it does not increaſe in ſo great a Proportion) the whole Reſiſtance of the Wood will be 190, that is, 100 from its Denſity, and 90 from its Tenacity. Here the whole Reſiſtance of the Water, is to the whole Reſiſtance of the Wood, as 110 to 190, or as 11 to 19. If the Velocity of the Bullet be doubled, that is, becomes 20, the whole Reſiſtance of the Water will be $400 + 20 = 420$; and the Reſiſtance of the Wood will be $400 + 180 = 580$: Then will the Reſiſtance of the Water be to that of the Wood, as 42 to 58, or 21 to 29, the Difference being now leſs than before. If the Velocity of the Bullet be 40, the Reſiſtance of the Water will be $1600 + 40 = 1640$, and that of the Wood $1600 + 360 = 1960$; and then the Reſiſtance of the Water will be to that of the Wood, as 164 to 196, or 41 to 49, the Difference being ſtill leſs. Therefore,

Annotat. fore, as the Velocity of the Bullet increases, the Resistance of the Water
 Lect. V. continually comes nearer to that of the Wood. Now, if the Wood be
 less dense than the Water, the Resistance of the Water on increasing the
 Velocity of the Bullet) will soon be greater than that of the Wood.
 Likewise, tho' the Wood be as dense as the Water, if the Resistance, on
 account of the Quantity of Matter, does not increase so fast as it does in
 the Water (as I said above, and Experience shews) by increasing the Bul-
 let's Velocity, the Resistance of the Water will soon overtake that of the
 Wood. The following Experiment has been made to confirm this Assertion.

A Sail was spread horizontally in a Pond about $2\frac{1}{2}$ Feet under the Sur-
 face of the Water, and a Musket with a small Charge being fir'd oblique-
 ly to the Surface of the Water, but perpendicularly to an half Inch Deal
 Board fix'd under the Water over the Sail, the Bullet went through the
 Board under the Water, and after that through the Sail-cloth. The
 Experiment was made again with a greater Charge, and the Bullet struck
 against the Board without going through it, making but a small Impres-
 sion in the Board, yet its Roundness was alter'd more than the Resistance
 of the Board seem'd likely to have caus'd, and therefore the Bruise in the Lead
 was thought owing to the Resistance of the Water. Upon increasing the
 Charge a third time, the Ball fell upon the Sail without reaching the
 Board, and was much beaten out of Shape. At last putting in a Charge
 pretty near equal to the Proof (that is, the Weight of Powder equal to
 the Weight of the Bullet) the Bullet was beaten to pieces upon the Sur-
 face of the Water. N.B. *I cannot exactly tell in what Proportion the
 Charges were increas'd, because I did not see the Experiment my self; but I
 had the Account from Persons of Credit that saw them.* That the Resis-
 tance of the Water was greater than that of Wood appear'd by this, viz.
 that when the Shot was made against a Plank of $1\frac{1}{2}$ Inch thick in the
 Air, the Bullet went through more and more easily the more the Charge
 was increas'd.

This great Resistance of Water is farther confirm'd, by an Accident,
 that happen'd to my self.

Some Years ago, on a rejoycing Day, the Honourable Colonel *Samuel
 Horsey*, with several other Persons, and my self, went out upon the *Thames*
 in a great Barge belonging to the *Vintners Company*, where we play'd
 off some Fire-works. It happen'd, that a Water-rocket (whose Property
 is to go under Water several times and rise again, and at last burst on
 the Top of the Water) came up, when it was ready to burst, under the
 Stern of the Barge, being thereby prevented from coming up to the Sur-
 face of the Water, and bursting where it was, gave the Barge a great
 Shock, so as to give it a sensible Lift, which I plainly felt, being directly
 over the Place, and wonder'd at the Force of so small a Quantity of
 Powder, there being much less than an Ounce of Powder to make
 the Report. A little after, some Gentlemen and Ladies, being in ano-
 ther Barge, of less than half the Bigness of ours, came near us, and de-
 fir'd me to throw some of the Water-rockets between the two Barges,
 that

that they might the better observe their Motion. I threw several; but at last one of them in its last Rise, stopping under the Middle of their Barge, broke there, and made so great an Hole in the Barge's Bottom, that there was only time for the Company to get out of their Barge into ours, and the Watermen to row to Shoar from the Middle of the Thames (at Mortlack, where it is but narrow) before the Barge was half full of Water. Because in the Explosion of the Powder, whose Velocity, in its Expansion, is exceedingly great, the Water resisted, like a Solid, and consequently the Powder made its Way through the Bottom of the Barge, whose Planks were less dense than the Water, and not very thick.

After this, to try the Effect of the Explosion of Gun-powder under Water; I loaded a Water-rocket, so that it should break under the Water, and having set Fire to it, threw it into a Pond, that cover'd an Acre of Ground: And so great was the Shock, that several Persons, who stood round the Pond, felt it like a momentaneous Earthquake.

Annotat.
Lect. V.

6. [39. Page 349. *Sir Isaac Newton has demonstrated it, &c.*]

Take Sir *Isaac Newton's* two Demonstrations in his own Words.

BOOK I. SECT. 2. PROPOSITION I. THEOREM I.

" *The Areas, which revolving Bodies describe by Radii, drawn to an immoveable Center of Force, do lye in the same immoveable Planes, and are proportional to the Times in which they are describ'd.* Plate 28. Fig. 9.] Pl. 28. F. 9.

" For, suppose the Time to be divided into equal Parts; and in the first Part of that Time, let the Body by its innate Force describe the right Line AB. In the second Part of that Time, the same would (by Law 1.) if not hinder'd, proceed directly to c, along the Line Bc equal to AB; so that by the Radii AS, BS, cS drawn to the Center, the equal Areas ASB, BS c, would be described. But when the Body is arrived at B; suppose, that a centripetal Force acts at once with a great Impulse, and turning aside the Body from the right Line Bc, compels it afterwards to continue its Motion along the right Line BC. Draw cC parallel to BS meeting BC in C; and at the End of the second Part of the Time, the Body (by Cor. 1. of the Laws) will be found in C, in the same Plane with the Triangle ASB. Join SC, and, because SB and Cc are parallel, the Triangle SBC will be equal to the Triangle SB c, and therefore also to the Triangle SAB. By the like Argument, if the centripetal Force acts successively in C, D, E, &c. and makes the Body in each single Particle of Time, to describe the right Lines CD, DE, EF, &c. they will all lye in the same Plane; and the Triangle SCD will be equal to the Triangle SBC, and SDE to SCD, and SEF to SDE. And therefore, in equal Times, equal Areas are describ'd, in one immoveable Plane: And, by Composition; any Sums SADS, SAFS, of those Areas, are

" one

Annotat. " one to the other, as the Times in which they are describ'd. Now,
 Lect. V. " let the Number of those Triangles be augmented, and their Breadth
 " be diminish'd *in infinitum*; and (by COR. 4. Lem. 3.) their ultimate Peri-
 Pl. 28. F. 9. " meter ADF will be a curve Line; and therefore the centripetal Force,
 " by which the Body is perpetually drawn back from the Tangent of this
 " Curve, will act continually; and any describ'd Areas SADS, SAFS,
 " which are always proportional to the Times of Description, will, in this
 " Case, also be proportional to those Times. Q. E. D.

" Cor. 1. The Velocity of a Body attracted towards an immoveable
 " Center, in Spaces void of Resistance, is reciprocally as the Perpendicu-
 " lar let fall from that Center on the right Line that touches the Orbit. For
 " the Velocities in those Places A, B, C, DE are as the Bases AB, BC,
 " CD, DE, EF, of equal Triangles; and these Bases are reciprocally
 " as the Perpendiculars let fall upon them.


" Cor. 2. If the Chords AB, BC of two Arcs, successively describ'd
 " in equal Times, by the same Body, in Spaces void of Resistance, are
 " compleated into a Parallelogram ABCV, and the Diagonal BV of this
 " Parallelogram, in the Position, which it ultimately acquires, when those
 " Arcs are diminish'd *in infinitum*, is produced both ways, it will pass
 " through the Center of Force.

" Cor. 3. If the Chords AB, BC, and DE, EF of Arcs describ'd
 " in equal Times, in Spaces void of Resistance, are compleated into the
 " Parallelograms ABCV, DEFZ; the Forces in B and E, are one
 " to the other in the ultimate Ratio of the Diagonals BV, EZ, when
 " those Arcs are diminished *in infinitum*. For the Motions BC and EF
 " of the Body (by Cor. 1. of the *Laws*) are compounded of the Motions
 " Bc, BV, and Ef EZ; but BV and EZ, which are equal to Cc
 " and Ff, in the Demonstration of this Proposition, were generated by
 " the Impulses of the centripetal Force in B and E, and are therefore pro-
 " portional to those Impulses.

" Cor. 4. The Forces by which Bodies in Spaces void of Resistance,
 " are drawn back to rectilinear Motions, and turned into curvilinear Or-
 " bits, are one to another as the versed Sines of Arcs describ'd in equal
 " Times; which versed Sines tend to the Center of Force, and bissect the
 " Chords, when those Arcs are diminished to Infinity: For such versed
 " Sines are the Halves of the Diagonals mention'd in Cor. 3.

" Cor. 5. And therefore those Forces are to the Force of Gravity, as
 " the said versed Sines to the versed Sines perpendicular to the Horizon of
 " those parabolick Arcs, which Projectiles describe in the same Time.

" Cor.

“ *Cor. 6.* And the same Things do all hold good (by *Cor. 5.* of the Annotat.
 “ *Laws*) when the Planes in which the Bodies are mov’d, together with the *Lect. V.*
 “ Centers of Force which are placed in those Planes, are not at Rest, 
 “ but move uniformly forward in right Lines.

“ PROPOSITION II. THEOREM II.

“ *Every Body, that moves in any curve Line describ’d in a Plane, and*
 “ *by a Radius, drawn to a Point, either immoveable, or moving forward*
 “ *with an uniform rectilinear Motion, describes about that Point Areas pro-*
 “ *portional to the Times, is urged by a centripetal Force directed to that*
 “ *Point.*

“ *Case 1.* For every Body, that moves in a curve Line, is (by *Law*
 “ *1.*) turned aside from its rectilinear Course, by the Action of some
 “ Force, that impels it. And that Force, by which the Body is turned
 “ off from its rectilinear Course, and is made to describe, in equal Times,
 “ the equal least Triangles *SAB, SBC, SCD, &c.* about the im- * *Pl. 28. F. 9.*
 “ moveable Point *S* (by *Prop. 40. Book 1. El. and Law 2.*) acts in the
 “ Place *B*, according to the Direction of a Line parallel to *cC*, that is,
 “ in the Direction of the Line *BS*; and in the Place *C*, according to
 “ the Direction of a Line parallel to *dD*, that is, in the Direction of
 “ the Line *CS*, &c. And therefore acts always in the Direction of
 “ Lines tending to the immoveable Point *S*. *Q. E. D.*

“ *Case 2.* And (by *Cor. 5.* of the *Laws*) it is indifferent, whether the
 “ Superficies in which a Body describes a curvilinear Figure be quiescent,
 “ or moves together with the Body, the Figure describ’d, and its Point *S*,
 “ uniformly forwards in right Lines.

“ *Cor. 1.* In non-resisting Spaces or Mediums, if the *Areas* are not
 “ proportional to the Times, the Forces are not directed to the
 “ Point in which the *Radii* meet; but deviate therefrom *in consequentia*,
 “ or towards the Parts to which the Motion is directed, if the Descrip-
 “ tion of the *Areas* is accelerated; but in *antecedentia*, if retarded.

“ *Cor. 2.* And even in resisting Mediums, if the Description of the
 “ *Areas* is accelerated, the Directions of the Forces deviate from the
 “ Point in which the *Radii* meet, towards the Parts to which the Mo-
 “ tion tends.

SCHOLIUM.

“ A Body may be urged by a centripetal Force compounded of several
 “ Forces. In which Case the Meaning of the Proposition is, that the
 “ Force which results out of all, tends to the Point *S*, &c.

G g g

I have

Annotat. I have contriv'd a Machine to represent this mechanically, wherein a
 Lect. V. little Brass Ball representing a Planet, moves in an Ellipse about a Body
 representing the Sun in one of the *Foci* of that Ellipse, describing by a
Radius vector Areas proportional to the Time. N. B. I shall give a Fi-
 gure and Description of it at the End of this Volume.

— That the centripetal Force is the same as Gravity.

Sir *Isaac Newton* demonstrates this in the 4th Proposition of the third Book of his *Principia*.

“ PROP. 4. THEOR. 4. That the Moon gravitates towards the Earth;
 “ and by the Force of Gravity is continually drawn off from a rectilinear Mo-
 “ tion, and retain'd in its Orbit.

Sir *Isaac*, having given an Account of the Moon's mean Distance, ac-
 cording to several Astronomers, in the Beginning of this Proposition, goes
 on in these Words. “ Let us assume the mean Distance of 60 Semi-dia-
 “ meters in the *Syzygies*; and suppose one Revolution of the Moon, in
 “ respect of the fix'd Stars, to be compleated in 27 d. 7 h. 43 m. as
 “ Astronomers have determined; and the Circumference of the Earth to
 “ amount to 123249600 *Paris Feet*, as the *French* have found by Mensu-
 “ ration. And now, if we imagine the Moon, deprived of all Motion,
 “ to be let go, so as to descend towards the Earth, with the Impulse of
 “ all that Force, by which (by *Cor. Prop. 3.*) it is retained in its Orb;
 “ it will, in the Space of one Minute of Time, describe in its Fall
 “ $15 \frac{1}{21}$ *Paris Feet*. This we gather by a *Calculus*, founded either upon
 “ *Prop. 36.* Book 1. or (which comes to the same Thing) upon *Cor. 9.*
 “ *Prop. 4.* of the same Book. For the versed Sine of that Arc, which
 “ the Moon, in the Space of one Minute of Time, would by its mean
 “ Motion describe at the Distance of 60 Semi-diameters of the Earth, is
 “ nearly $15 \frac{1}{12}$ *Paris Feet*, or more accurately 15 Feet, 1 Inch, and 1
 “ Line $\frac{4}{5}$. Wherefore, since that Force, in approaching to the Earth, in-
 “ creases in the reciprocal duplicate Proportion of the Distance, and,
 “ upon that Account, at the Surface of the Earth, is 60×60 times great-
 “ er, than at the Moon; a Body in our Regions, falling with that
 “ Force, ought, in the Space of one Minute of Time, to describe
 “ $60 \times 60 \times 15 \frac{1}{12}$ *Paris Feet*, and, in the Space of one Second of Time,
 “ to describe $15 \frac{1}{12}$ of those Feet; or more accurately, 15 Feet, 1 Inch,
 “ and 1 Line $\frac{4}{9}$. And with this very Force, we actually find, that Bo-
 “ dies here upon Earth do really descend. For a Pendulum oscillating
 “ Seconds in the Latitude of *Paris*, will be 3 *Paris Feet*, and 8 Lines $\frac{1}{2}$
 “ in Length, as Mr. *Huygens* has observ'd. And the Space, which an
 “ heavy Body describes by falling in one Second of Time, is to half the
 “ Length of this Pendulum, in the duplicate *Ratio* of the Circumference
 “ of a Circle to its Diameter (as Mr. *Huygens* has also shewn) and is there-
 “ fore

“ fore 15 *Paris* Feet, 1 Inch, 1 Line $\frac{3}{4}$. And therefore the Force, by Annotat.
 “ which the Moon is retained in its Orbit, becomes, at the very Surface *Lect. V.*
 “ of the Earth, equal to the Force of Gravity, which we observe in
 “ heavy Bodies there. And therefore (by Rule 1. and 2.) the Force by
 “ which the Moon is retained in its Orbit, is that very same Force, which
 “ we commonly call Gravity. For, were Gravity another Force different
 “ from that; then Bodies descending to the Earth, with the joint Im-
 “ pulse of both Forces, would fall with a double Velocity, and in the
 “ Space of one Second of Time, would describe $30\frac{1}{2}$ *Paris* Feet; al-
 “ together against Experience, &c.

S C H O L I U M.

“ The Demonstration of this Proposition may be more diffusely ex-
 “ plained after the following manner. Suppose several Moons to revolve
 “ about the Earth, as in the System of Jupiter or Saturn; the periodic
 “ Times of these Moons (by the Argument of Induction) would observe the
 “ same Law which *Kepler* found to obtain among the Planets; and there-
 “ fore their centripetal Forces would be reciprocally as the Squares of the
 “ Distances from the Center of the Earth, by *Prop. 1.* of this *Book*. Now,
 “ if the lowest of these was very small, and was so near the Earth, as
 “ almost to touch the Tops of the highest Mountains; the centripetal
 “ Force thereof, retaining it in its Orb, would be very nearly equal to
 “ the Weights of any terrestrial Bodies, that should be found upon the
 “ Tops of those Mountains, as may be known by the foregoing Computa-
 “ tion. Therefore, if the same little Moon should be deserted by its cen-
 “ trifugal Force, that carries it through its Orbit, and so be disabled
 “ from going onwards therein, it would descend to the Earth; and that
 “ with the same Velocity as heavy Bodies do actually fall with, upon the
 “ Tops of those very Mountains; because of the Equality of the Forces
 “ that oblige them both to descend. And, if the Force by which that
 “ lowest Moon would descend, were different from Gravity; and if that
 “ Moon were to gravitate towards the Earth, as we find terrestrial Bodies
 “ do upon the Tops of Mountains, it would then descend with twice the
 “ Velocity, as being impelled by both these Forces conspiring together.
 “ Therefore, since both those Forces, that is, the Gravity of heavy Bodies,
 “ and the centripetal Forces of the Moons, respect the Center of the Earth,
 “ and are similar and equal between themselves, they will (by Rule 1.
 “ and 2.)† have one and the same Cause. And therefore the Force, which
 “ retains the Moon in its Orbit, is that very Force, which we commonly
 “ call Gravity; because, otherwise, this little Moon at the Top of a
 “ Mountain, must either be without Gravity, or fall twice as swiftly as
 “ heavy Bodies use to do.

† See the Beginning of Sir Isaac's third Book.

7. [40. Page 351, 352. Since the Angles, &c. — only decrease half way from A to P, &c. And the Angles, &c. — only increase half way from

G g 2

P to

Annotat. P to A, &c.] This will be made evident by proving, that the Angle *Lect. V.* made by a Line drawn from either of the *Foci*, to any Point in the Periphery of an Ellipse, and the Tangent to the Curve in that Point, will be the least possible, or at its *Minimum*, when that Point is at either of the **Pl. 28. F. 10.* Ends of the lesser Axis; that is, *SHT* * is less than *SI t*. And this we shall prove by the Help of the two following *Lemmata*.

L E M M A I.

In every quadrilateral Figure, inscribed in a Circle, the Sum of the two Rectangles form'd by the opposite Sides, are equal to the Rectangle form'd by the two Diagonals of that quadrilateral Figure.

This is a known Property of the Circle, and has been demonstrated by *†Pl. 28. F. 11. Ptolemy*, and others; that is, $\overline{AC \times DB} + \overline{BC \times AD} = \overline{AB \times DC}$ †.

L E M M A II.

If an Arc of a Circle be divided into two equal, and into two unequal Parts. I say, that the Sum of the Chords of the two equal Arcs, will always be greater than the Sum of the Chords of the two unequal Arcs. That is, if the Arc ACKB is divided into two equal Parts at C, and into unequal Parts at K. $AC + CB$ is greater than $AK + KB$.

In order to prove this; through C and the Center *k*, draw the Diameter CD, and the Diagonals and Chords AB, DK, AD, BD. It is evident (by the 2d and 30th of the 3d of *Euch.*) $AD = BD$, and they may be substituted for one another. It is evident also, that DK will always be less than CD (by *Euch.* 14th of the 3d.) Now by the last Lemma $\overline{AK \times DB} + \overline{KB \times AD} = \overline{AB \times DK}$, now since DK has been prov'd less than DC, it must follow, that $\overline{AB \times DC}$ is greater than $\overline{AB \times DK}$; therefore $\overline{AC \times DB} + \overline{BC \times AD}$ is greater than $\overline{AK \times DB} + \overline{KB \times AD}$; and since the Factors DB and AD are equal, it must follow, that $AC + CB$ is greater than $AK + KB$.

Pl. 28. F. 10. Let a Circle in *Fig. 10.* be described passing through the Points S, H, *s*, 'tis evident (by *Euch.* 19th of the 3d) that the Angle *SH s* and *SM s*, are equal; and 'tis evident by the 2d Lemma, that $\overline{SH} + \overline{H s}$, is greater than $\overline{SM} + \overline{M s}$. Now by the Property of the Ellipse $\overline{SI} + \overline{I s}$, is always equal to $\overline{SH} + \overline{H s}$, being always equal to AP; it must therefore follow, that $\overline{SI} + \overline{I s}$ is greater than $\overline{SM} + \overline{M s}$; whence it appears, that the Point I is out of the Circle *SHM s*, and therefore the Angle *SI s* is less than *SM s*, or its equal *SH s*. Now by the Property of the Ellipse demonstrated in all the Conick Writers (*vide Elementa Conicorum Lud. Trevigar Lib. 2. Prop. 10.*) The Angles *sIg* and *SI t* are equal, and *sHG* is equal to *SHT*; therefore, since *sIS* is less than *sHS*, it must follow, that *SI t*, is greater than *SHT*, and what has been prov'd of I may be prov'd of the Point F, or any other Point in the Semi-ellipse *AHP*; therefore the Truth of the Proposition advanced is demonstrated.

8. [42. Page 354. *Comets — are — Planets moving in very excentrick* Annotat. *Orbits, &c.*] One may have a full Account of Comets, and their Mo- Lect. V. tions, in Dr. *Halley's* Treatise of Comets, which was first printed by it self, and then reprinted at the End of the *English* Translation of Dr. *Gregory's* Astronomy; as also in Sir *Isaac Newton's* 3^d Book of his *Principia*, from the Beginning of the 39th Prop. to the End of the 42^d, where his general *Scholium* begins. The Reader may be fully satisfied about Comets by consulting those Authors. However, it may not be improper to give a short Account of them here, by way of Abstract.

COMETS are a Sort of excentrick Planets, which move in very long Ellipses about the Sun in one of the *Foci* of those Ellipses, whose periodical Revolutions take up such a long Space of Time, that the same Man has never yet seen the same Comet twice. So that the Astronomy of Comets is but lately brought to any tolerable Perfection, for want of a sufficient Number of Observations: And the Accounts we have of them in former Ages, are from Historians, and not Astronomers; which last, while they took Pains to observe the Planets and fix'd Stars, neglected such Comets as appear'd in their Times, being carried away with the Opinion, that they were only Meteors near the Earth, and not to be reckoned among the celestial Bodies. But in the last Age, Comets were first found to be higher than the Moon, and consequently to belong to the heavenly Bodies; and for about 70 Years last past, Astronomers have been very diligent in observing such as have appear'd. Sir *Isaac Newton*, from his own Observations, and those of others, has given a Method how, from only three Observations of a Comet, made at different Times, to find out all that Part of its Orbit, in which it will continue visible to us; and Dr. *Halley* has first shewn, that the same Comet returns again after a certain Number of Years, and that the whole Orbit is elliptical. From Astronomers and Historians, we are inform'd of 25 Appearances of Comets within these last 400 Years, tho' the Comets belonging to our System may be fewer in Number; because the same Comets have been seen several times; but were not known to be the same, till Dr. *Halley* shew'd it, and settled the Orbits of three of them, making out, that the Comet seen in 1661 had been seen before, and observ'd to go the same Way in the Year 1532; which makes its Period to be of 129 Years. The Comet seen in the Year 1680 and 1681, appear'd before in the Year 1106, so that its Period is about 575 Years. Lastly, The Comet of 1682 appear'd in the Year 11607 and 1531, which shews its Period to consist of 75 Years. So that the Number of the Comets which have been seen will be reduced to 22. And if, according to Mr. *Whiston's* Conjectures, the Comet of 1682, appear'd also in 1456; and that of 1681 in 532, as also 44 Years before the Christian *Æra*, as Sir *Isaac Newton* and Dr. *Halley* have since discover'd, the Number will not be increas'd, since those Periods are before the Time of the first of the 25. But yet we cannot say, there are but 22, because several of them may have been at such a Distance from the Earth,

Annotat. Earth, as to have escap'd Observation; as the last might have done, if
 Lect. V. the Right Honourable the Lord *Pairley* had not discovered the last by
 chance in the Year 1723, and given Notice of it to other Astronomers,
 that they might also observe it.

The Comets are reckon'd not to be less than the Moon, nor much bigger than Venus; and those are the least, which come nearest to the Sun, lest they should agitate the Sun too much, as they come by it at their *Perihelion*, and likewise disturb those Planets which are near the Sun. That they are not very large appears, because they are not visible till they come nearer to us than Jupiter's Orbit; for as they are opaque Bodies, that reflect Light from the Sun, they must have been seen as far as Saturn, if they had been as big.

Whereas all the Planets have the Planes of their Orbits in, or near, the Plane of the Ecliptick, the most inclin'd making an Angle but of few Degrees; the Comets have their Orbits inclin'd to the Ecliptick in very great Angles, moving towards all Parts of the Heavens; and by that Means are less liable to disturb the Planets, because they come but twice in one Revolution into the Plane of the Ecliptick, and when they are at their *Aphelia*, very distant from the Sun, they are also very distant from each other, and thereby not so subject to attract each other with sufficient Force to disturb one another's Orbits sensibly; for when they move very slowly at their *Aphelia*, they are easily drawn out of their usual Course.

Comets are considerably rarer than the Planets; for if they were as solid, they would not have those Tails, which are observable, as they come near enough to be warm'd by the Sun, with a Degree of Heat, something less than that of the Planet Mars: Those Tails being Vapours rais'd from their *Nucleus*, or Globe, so as to make a very large Atmosphere, whose Particles being thrown off on that Side, which is opposite to the Sun, give that Appearance, which is call'd a *Tail*, when we see it at right Angles, or any great Angle; a *Beard*, when we see it obliquely, and a *Coma* or *Hairiness*, when we see it with so great an Obliquity, as to be almost in the Line of its Axis, so that the Vapour appears to encompass the Comet like a *Periwig*, or *Head of Hair*: And these three Appearances are owing to the different Positions of the Earth, in respect of the Comet; for a Tail of the same Bigness varies optically according to the Position of the Eye.

But then the Tail it self is continually changing; for often the Comet begins to appear without any sensible Tail, and by degrees the Tail is generated, and increases continually, as the Comet comes nearer to the Sun, and is the biggest of all when the Comet is just past the *Perihelion*, having then receiv'd its greatest Degree of Heat.

If the celestial Spaces, in which the Comets move, were not void of Matter, we could not see their Tails, because they are abundantly rarer than our Air, since one may plainly see the fix'd Stars through them. As we see the Light of the Sun reflected from those Particles that float about in the Air, when a Sun-beam of 1 or 2 Inches wide is let into a dark Room; so is the Light of the Sun-beams reflected to us from the Particles

cles of the Tails of Comets, because there is no other sensible Matter near them: For if there was a celestial Matter about them, tho' 50000 times rarer than our Air, it would reflect Light enough to confound that reflected from the Tail of the Comet in one undistinguish'd Glare: So that Comets would not appear to have any Tails at all; or, at most, but very small ones. Annotat. Lect. V.

The Particles or Vapours, that form the Tail, seem to be at first darted every way from the Comet; but on that Side, which is towards the Sun, they turn back again in parabolick Lines, so as to go on the contrary Way, and to help to increase the Tail on the Side opposite to the Sun, as if the Sun's Rays gave them an Impulse, and drove them that Way. The 12th Figure of Plate 28, represents a Comet, and the lower Part of its Tail, as Dr. Hook carefully observ'd it through a Telescope near its Perihelion, with the purer Part of its Atmosphere, winding it self into the Tail, and the cloudy Part of it plac'd round about the central Solid. But, as the Comet of 1680 and 1681 was the most remarkable, and most nicely observ'd by Sir Isaac Newton, and many other Astronomers; it will be sufficient to give an Account of that, and the Figure of that Part of its Orbit, wherein it was visible to us, such as Sir Isaac Newton has given it in the last Edition of his *Principia*. And, as I can't give his Sense better than in his own Words, I shall take the Liberty to make use of them.

" I have given (*Pl. 28. Fig. 13.*) a true Representation of the Orbit, Pl. 28. F. 13.
 " which this Comet describ'd, and of the Tail which it emitted in several Places, in the annexed Figure; protracted in the Plane of the Trajectory. In this Scheme ABC represents the Trajectory of the Comet, D the Sun, DE the Axis of the Trajectory, DF the Line of the Nodes, GH the Intersection of the Sphere of the Orbit of the Earth, with the Plane of the Trajectory, I the Place of the Comet, *Nov. 4. Anno 1680*, K the Place of the same, *Nov. 11.* L the Place of the same, *Nov. 19.* M its Place, *Dec. 12.* N its Place, *Dec. 21.* O its Place, *Dec. 29.* P its Place, *Jan. 5.* following. Q its Place, *Jan. 25.* R its Place, *Feb. 5.* S its Place, *Feb. 25.* T its Place, *March 5.* And V its Place, *March 9.* In determining the Length of the Tail, I made the following Observation.

" *Nov. 4. and 6.* the Tail did not appear; *Nov. 11.* the Tail just begun to shew it self, but did not appear above half a deg. long, through a 10 Foot Telescope; *Nov. 17.* the Tail was seen by Ponthæus more than 15 deg. long, &c.

Then Sir Isaac proceeds, in giving an Account of several Persons, as well as himself, who observ'd the Tail, and all its Changes, in Increase and Decrease, as it appear'd to them in several Countries; and at last he goes on thus.

" *Feb. 25.*

Annotat.
Lect. V.



" Feb. 25. the Comet was without a Tail, and so it continued till it disappear'd.

" Now, if one reflects upon the Orbit describ'd, and duly considers the other Appearances of this Comet, he will be easily satisfied, that the Bodies of Comets are solid, compact, fix'd, and durable, like the Bodies of the Planets. For, if they were nothing else, but the Vapours of Exhalations of the Earth, of the Sun, and other Planets, this Comet in its Passage by the Neighbourhood of the Sun, would have been immediately dissipated. For the Heat of the Sun is as the Density of its Rays, that is, reciprocally as the Square of the Distances of the Places from the Sun. Therefore, since on Dec. 8. when the Comet was in its *Perihelion*, the Distance thereof from the Center of the Sun, was to the Distance of the Earth from the same, as about 6 to 1000; the Sun's Heat on the Comet was at that time to the Heat of the Summer's Sun with us, as 1000000 to 36, or as 28000 to 1. But the Heat of boiling Water is about 3 times greater than the Heat, which dry Earth acquires from the Summer-Sun, as I have try'd; and the Heat of red hot Iron (if my Conjecture is right) is about 3 or 4 times greater than the Heat of boiling Water. And therefore the Heat which dry Earth on the Comet, while in its *Perihelion*, might have conceived from the Rays of the Sun, was about 2000 times greater than the Heat of red hot Iron. But by so fierce a Heat, Vapours, and Exhalations, and every volatile Matter, must have been immediately consum'd and dissipated.

" This Comet, therefore, must have conceiv'd an immense Heat from the Sun, and retain that Heat for an exceeding long Time. For a Globe of Iron of an Inch Diameter, expos'd red-hot to the open Air, will scarcely lose its Heat in an Hour's time; but a greater Globe would retain its Heat longer in Proportion of its Diameter, because the Surface (in Proportion to which it is cool'd by the Contact of the ambient Air) is in this Proposition less, in respect of the Quantity of the included hot Matter. And therefore a Globe of red hot Iron, equal to our Earth, that is about 40000000 Feet in Diameter, would scarcely cool in an equal Number of Days, or in above 50000 Years. But I suspect, that the Duration of Heat, may on account of some latent Causes, increase in a yet less Proportion, than that of the Diameter; and I should be glad, that the true Proportion was investigated by Experiments.

" It is further to be observ'd, that the Comet in the Month of *December*, just after it had been heated by the Sun, did emit a much longer Tail, and much more splendid, than in the Month of *November* before, when it had not yet arriv'd at its *Perihelion*. And universally, the greatest and most fulgent Tails always arise from Comets, immediately after their passing by the Neighbourhood of the Sun. Therefore the Heat received by the Comet, conduces to the greatness of the Tail. From whence, I think, I may infer, that the Tail is nothing
" else


“ else but a very fine Vapour, which the Head, or *Nucleus* of the Comet
 “ emits by its Heat, &c. Annotat.
Lect. V.

After this, our incomparable Author gives an Account of three several
 Opinions concerning the Tails of Comets ; and having refuted two of
 them, goes on to prove the third (which is also his own) in these
 Words. —

“ That the Tails of Comets do arise from their Heads, and tend to-
 “ wards the Parts opposite to the Sun, is further confirm'd from the
 “ Laws, which the Tails observe. As, that lying in the Planes of the
 “ Comets Orbits, which pass through the Sun, they constantly deviate
 “ from the Opposition of the Sun towards the Parts, which the Comets
 “ Heads in their Progress along these Orbits have left. That to a Spec-
 “ tator, placed in those Planes, they appear in the Parts directly opposite
 “ to the Sun ; but as the Spectator recedes from these Planes, their De-
 “ viation begins to appear, and daily becomes greater. That the Devi-
 “ ation, *ceteris paribus*, appears less, when the Tail is more oblique to
 “ the Orbit of the Comet, as well as when the Head of the Comet ap-
 “ proaches nearer to the Sun ; especially, if the Angle of Deviation is
 “ estimated nearer the Head of the Comet. That the Tails, which have
 “ no Deviation, appear strait, but the Tails which deviate, are likewise
 “ bent into a certain Curvature. That the Curvature is greater, when
 “ the Deviation is greater ; and is more sensible, when the Tail, *ceteris*
 “ *paribus*, is longer : For in the shorter Tails, the Curvature is hardly to
 “ be perceiv'd. That the Angle of Deviation is less near the Comet's
 “ Head, but greater towards the other End of the Tail ; and that, be-
 “ cause the convex Side of the Tail regards the Parts, from which the De-
 “ viation is made, and which lye in a right Line, drawn out infinitely
 “ from the Sun, through the Comet's Head. And that the Tails, that
 “ are long and broad, and shine with a stronger Light, appear more re-
 “ splendent, and more exactly defin'd on the convex than on the concave
 “ Side. Upon which Accounts, it is plain, that the *Phænomena* of the
 “ Tails of Comets depend upon the Motions of their Heads, and by no
 “ means upon the Places in the Heavens in which their Heads are seen ;
 “ and, that therefore the Tails of Comets do not proceed from the Re-
 “ fraction of the Heavens, but from their own Heads, which furnish the
 “ Matter that forms the Tail. For as, in our Air, the Smoak of a heated
 “ Body ascends, either perpendicularly, if the Body is at Rest, or oblique-
 “ ly, if the Body is moved obliquely : So in the Heavens, where all Bodies
 “ gravitate towards the Sun, Smoak and Vapour must (as we have already
 “ said) ascend from the Sun, and either rise perpendicularly, if the smoak-
 “ ing Body is at Rest ; or obliquely, if the Body in all the Progress of
 “ its Motion, is always leaving those Places, from which the upper or
 “ higher Parts of the Vapour had risen before. And that Obliquity will
 “ be least, where the Vapour ascends with most Velocity, to wit, near the
 “ H h h “ smoking

Annotat. "fmoaking Body, when that is near the Sun. But because the Obliquity
 Lect. V. "varies, the Column of Vapour will be incurvated; and, because the Va-
 "pour in the preceding Side, is something more recent, that is, has as-
 "cended something more late from the Body, it will therefore be some-
 "thing more dense on that Side, and must on that account reflect more
 "Light, as well as be better defin'd.

Then Sir *Isaac*, shewing how very rare our Air must be at the Distance of one Semi-diameter of the Earth from its Surface; makes Use of it, as an Argument, to shew, that a very small Quantity of Air and Vapour is abundantly sufficient to produce all the Appearances of the Tails of Comets. Then goes on shewing—"That we may pretty nearly determine the Time spent during the Ascent of the Vapour from the Comet's Head, to the Extremity of the Tail, by drawing a right Line from the Extremity of the Tail to the Sun, and marking the Place where that right Line intersects the Comet's Orbit. For the Vapour, that is now in the Extremity of the Tail, if it has ascended in a right Line from the Sun, must have begun to rise from the Head, at the Time the Head was in the Point of Intersection. It is true, the Vapour does not rise in a right Line from the Sun, but retaining the Motion, which it had from the Comet before its Ascent, and compounding that Motion with its Motion of Ascent, arises obliquely. And, therefore, the Solution of the Problem will be more exact, if we draw the Line, which intersects the Orbit, parallel to the Length of the Tail; or rather (because of the curvilinear Motion of the Comet) diverging a little from the Line or Length of the Tail. And by means of this Principle, I found, that the Vapour, which *Jan. 25.* was in the Extremity of the Tail, had begun to rise from the Head before *Dec. 11.* and therefore had spent in its whole Ascent 45 Days; but that the whole Tail, which appear'd on *Dec. 10.* had finish'd its Ascent in the Space of the two Days then elapsed from the Time of the Comet's being in its *Perihelion*. The Vapour, therefore, about the Beginning, and in the Neighbourhood of the Sun, rose with the greatest Velocity, and afterwards continu'd to ascend with a Motion constantly retarded by its own Gravity; and the higher it ascended, the more it added to the Length of the Tail. And while the Tail continued to be seen, it was made up of almost all that Vapour, which had risen since the Time of the Comet's being in its *Perihelion*; nor did that Part of the Vapour, which had risen first, and which form'd that Extremity of the Tail, cease to appear, till its too great Distance, as well from the Sun, from which it receiv'd its Light, as from our Eyes, render'd it invisible. Whence also it is, that the Tails of other Comets, which are short, do not rise from their Heads, with a swift and continual Motion, and soon after disappear; but are permanent and lasting Columns of Vapours and Exhalations; which ascending from the Heads with a slow Motion of many Days, and partaking of the Motion of the Heads, which they had
 " from

“ from the Beginning, continue to go along together with them through
 “ the Heavens. From whence again, we have another Argument, prov-
 “ ing the celestial Spaces to be free and without Resistance; since in them, 
 “ not only the solid Bodies of the Planets and Comets, but also the ex-
 “ tremely rare Vapours of Comets Tails, maintain their rapid Motions
 “ with great Freedom, and for an exceeding long Time.

Then after having shewn, why the Tails of Comets, when they are ve-
 ry near the Sun, rise to such prodigious Heights from their Heads,
 and yet do not quit the Heads in the Motion of the Comets; he
 goes on.

“ The Tails therefore, that rise in the *Perihelion* Positions of the Co-
 “ mets, will go along with their Heads into far remote Parts, and toge-
 “ ther with the Heads will either return again from thence to us, after
 “ a long Course of Years; or rather, will be there rarified, and by de-
 “ grees quite vanish away. For afterwards, in the Descent of the Heads
 “ towards the Sun, new short Tails will be emitted from the Heads with
 “ a slow Motion; and those Tails by degrees will be augmented im-
 “ mensly, especially in such Comets as in their *Perihelion* Distances desc-
 “ end as low as the Sun's Atmosphere. For all Vapour in those free
 “ Spaces, is in a perpetual State of Rarefaction and Dilatation. And
 “ from hence it is, that the Tails of all Comets are broader at their up-
 “ per Extremity, than near their Heads. And it is not unlikely, but
 “ that the Vapour, thus perpetually rarify'd and dilated, may be at last
 “ dissipated, and scatter'd through the whole Heavens, and by little and
 “ little be attracted towards the Planets, by its Gravity, and mix'd with
 “ their Atmosphere. For, as the Seas are absolutely necessary to the Con-
 “ stitution of our Earth, that from them the Sun, by its Heat, may ex-
 “ hale a sufficient Quantity of Vapours, which being gathered together
 “ into Clouds, may drop down in Rain, for watering of the Earth, and
 “ for the Production and Nourishment of Vegetables; or being conden-
 “ sed with Cold on the Tops of Mountains (as some Philosophers with
 “ Reason judge) may run down in Springs and Rivers; so for the Conser-
 “ vation of the Seas, and Fluids of the Planets, Comets seem to be re-
 “ quir'd, that from their Exhalations and Vapours condens'd, the Wastes
 “ of the Planetary Fluids, spent upon Vegetation and Putrefaction, and
 “ converted into dry Earth, may be continually supply'd and made up.
 “ For all Vegetables entirely derive their Growth from Fluids, and after-
 “ wards in great Measure are turn'd into dry Earth by Putrefaction, and
 “ a Sort of Slime is always found to settle at the Bottom of putrified
 “ Fluids. And hence it is, that the Bulk of the solid Earth is conti-
 “ nually increased, and the Fluids, if they are not supply'd from without,
 “ must be in a continual Decrease, and quite fail at last. I suspect more-
 “ over, that 'tis chiefly from the Comets that Spirit comes, which is in-

Annotat. " deed the smallest, but the most subtil and useful Part of our Air, and
 Lect. V. " so much requir'd to sustain the Life of all things with us.

" The Atmospheres of Comets, in their Descent towards the Sun, by
 " running out into the Tails, are spent and diminish'd, and become nar-
 " rower, at least on that Side, which regards the Sun; and in receding
 " from the Sun, when they less run out into the Tails, they are again
 " enlarged, if *Hevelius* has justly mark'd their Appearances. But they
 " are seen least of all, just after they have been most heated by the Sun, and
 " on that account then emit the longest and most resplendent Tails, and
 " perhaps at the same time the *Nuclei* are environ'd with a denser and
 " blacker Smoak, in the lowermost Parts of their Atmosphere. For
 " Smoak, that is rais'd by a great and intense Heat, is commonly the
 " denser and blacker; thus the Head of that Comet, which we have
 " been describing, at equal Distances both from the Sun and from the
 " Earth, appear'd darker after it had pass'd by its *Perihelion*, than it did
 " before. For in the Month of *December* it was commonly compar'd
 " with the Stars of the third Magnitude; but in *November*, with those
 " of the first or second. And such as saw both Appearances, have de-
 " scrib'd the first, as of another and greater Comet than the second.

Sir *Isaac* confirms this, by the Observations of several Astronomers; then gives a Problem concerning a Comet's Trajectory (to which, and all that we have omitted of his on that Subject, I refer the mathematical Reader) and several Tables of Observations of Comets Places; and at last he speaks in general of Comets in the following Words.

" Because of the great Number of Comets, of the great Distance of
 " their *Aphelia* from the Sun, and of the Slowness of Motions in the
 " *Aphelia*, they will by their mutual Gravitations, disturb each other: So
 " that their Excentricities, and the Times of their Revolutions will be
 " sometimes a little increased, and sometimes diminished. Therefore we
 " are not to expect, that the same Comet will return exactly in the same
 " Orbit, and in the same periodick Times. It will be sufficient, if we
 " find the Changes no greater, than may arise from the Causes just spo-
 " ken of.

" And hence a Reason may be assign'd, why Comets are not compre-
 " hended within the Limits of a Zodiack as the Planets are; but being
 " confin'd to no Bounds, are with various Motions dispers'd all over the
 " Heavens; namely, to this Purpose, that in their *Aphelia*, where their
 " Motions are exceeding slow, receding to greater Distances one from
 " another, they may suffer less Disturbance from their mutual Gravitati-
 " ons. And hence it is, that the Comets, which descend the lowest,
 " and therefore move the slowest in their *Aphelia*, ought also to ascend
 " the highest.

" The

“ The Comet, which appear’d in the Year 1680, was in its *Perihelion*, Annotat.
 “ less distant from the Sun, than by a sixth Part of the Sun’s Diameter; *Lect. V.*
 “ and because of its extreme Velocity in that Proximity to the Sun, and
 “ some Density of the Sun’s Atmosphere, it must have suffer’d some Re-
 “ sistance and Retardation; and therefore, being attracted something near-
 “ er to the Sun in every Revolution, will at last fall down upon the Body
 “ of the Sun. Nay, in its *Aphelion*, where it moves the slowest, it may
 “ sometimes happen to be yet farther retarded by the Attractions of other
 “ Comets, and in consequence of this Retardation descend to the Sun. So
 “ fixed Stars, that have been gradually wasted by the Light and Vapours
 “ emitted from them for a long time, may be recruited by Comets that
 “ fall upon them, and from this fresh Supply of new Fuel, those old
 “ Stars acquiring new Splendour, may pass for new Stars. Of this kind are
 “ such fix’d Stars as appear on a sudden and shine with a wonderful Bright-
 “ nefs at first, and afterwards vanish by little and little.

9. [46. Page 357.—*Action and Reaction are equal and contrary, as well in the Repulsions as Attractions.*] So the Gravitation betwixt the Earth and its Parts is equal; for if we suppose the Earth FEHIK * to be divided *Pl. 28. F. 14. into two equal Parts, it is evident, that those Parts would come together with equal Forces. But if they be divided into two unequal Parts, viz. EFG and EIG by a Plane as EG; they will then also gravitate, or press towards one another with equal Forces (see *Lect. 1. No. 11.*) for if L. 1. No. 11. we cut off from the Part EIG, the smaller Part HIK equal to EFG; then will the greater Part EGHK, become the Obstacle, upon which the two equal Quantities of Matter EFG and HIK do press, whilst it self remains in *Equilibrio* between them. If we were to suppose EFG to be separated from EIG, in their Tendency towards each other by their Gravity, the little one would move so much faster towards the great one, as it has less Matter than the great one; so that the *Momentum* being equal in both, they will at meeting come to Rest, destroying each other’s Motion. For if the great one had more Force, it would go *in infinitum*, carrying EFG before it in the Direction IF.

The better to illustrate the Law of Action and Reaction, I think it may not be improper, to give my Reader the following Problem, communicated to me by a Friend.

P R O B L E M.

To find what is the Force with which the Breech of a Cannon is acted upon, as it shoots out a 24 lb Ball.

According to *Mersennus*, a 24 Pounder, weighing 6400 lb, it gives its Ball an uniform Velocity of 600 Paris Feet in a Second, which makes 640 English Feet. Now, let the Weight of the Cannon be called = c = 6400 lb.

Annotat. 6400 lb, the Weight of the Ball = b = lb, and the uniform Velocity, Lect. V. which the Powder gives it = V = 640 Feet in a Second. The *Momenta* of the Cannon, and the Ball given by the same Force of Powder, will be equal; therefore $cu = bV$; consequently, $6400 : 24 :: 640 : 2,4 = u$ = the Velocity of the Cannon, if it recoil'd on an horizontal Plane perfectly smooth. But if the Cannon cannot recoil, the Force of the Powder acting but by Way of Pressure, will give the Cannon a Shock, with a Force as 15630. Therefore, when the Cannon recoils, Part of the Force of the Powder is employ'd in giving it a Velocity of 2,4 Feet in a Second, and the remaining Part acts upon the Cannon by Way of Pressure. If then we would find how much the whole Pressure 15630 is diminish'd, when the Recoil is of 2,4 Feet in a Second; since the whole Pressure is to the partial Pressure, as the whole Velocity is to the partial Velocity, the two Parts will be, as $640 - 2,4 : 2,4$; or as $637,6 : 2,4$; or as $796 : 3$. Therefore we must divide 15630 in the same Proportion, making $797 : 3 :: 15630 - x : x$; consequently $797 \div 3$ (800) $3 : 15630 : x = 57,6$; which Quantity being taken from 15630, leaves a Pressure as 15572,4 when the Cannon recoils freely. But the Platform of the Battery not being perfectly smooth, and commonly rising backwards, the Recoil of the Cannon, which in such a Case is oblig'd to rise, will be so little, that one may consider very near the whole Force of 25630 as acting on the Cannon, and whatever stops it from recoiling.

Now, as it is not sufficient, only to compare the Weights and Velocities of the Ball and Cannon (because the Force, which is express'd by the Number 15630, might have been express'd by any other Number, as it would have been by 184320, if we had taken 7680 Inches, instead of 640 Feet in a Second for the Velocity of the Ball) we must shew, what Weight express'd in Number of Pounds pressing upon the Cannon, as a Weight laid on; is equal to the Force with which the Powder presses the Gun from within, as it throws out the Ball; which will be thus found. Let us suppose the Length of the Cannon within to be of 12 Feet, and that the Powder, as it expands in firing, drives out the Ball with a Velocity uniformly accelerated, so that the Ball, when out of the Cannon, shall by an uniform Velocity, move 24 Feet in the same time, that it mov'd 12 Feet in the Cannon by the accelerated Velocity. Therefore as the Ball, by that uniform Velocity, moves 640 Feet in a Second, it spends but $\frac{1}{26}$ and

$\frac{2}{3}$ of a Second in moving 24 Feet, and consequently spends but $\frac{1}{26\frac{2}{3}}$ in running the Length of the Inside of the Cannon. Now, since by accelerated Velocities the Spaces gone through, are as the Squares of the Times, then will $\left[\frac{1}{26\frac{2}{3}}\right]^2$ be to $[1 \text{ Second}]^2$; or, $1 : 26\frac{2}{3}^2 (= 711\frac{1}{9}) ::$ as 12 Feet:

to $8533\frac{1}{3}$ Feet, which the Ball would go in a Second with the accelerated Velocity, which it has in the Cannon; if therefore Gravity, with an accelerated

celerated Force of 16 Feet in a Second, gives the Ball a Force of 24 $\frac{1}{2}$ Annotat.
 Weight; the Action of the Powder, which is capable of giving it in the Lect. V.
 same time a Velocity of $8533\frac{1}{4}$ Feet in a Second, must give it a *Momen-*
tum, or Force against the Cannon, equal to a Weight of 12800 $\frac{1}{2}$; }
 because 16 Feet : are to 24 $\frac{1}{2}$:: as $8533\frac{1}{4}$ Feet : to 12800 $\frac{1}{2}$, since *Action*
 and *Reaction* are equal.

If we allow the Cannon any sensible Recoil; the Effect of the Pressure of the Powder must be diminish'd by the same Analogy, as we diminish'd the *Momentum* in the former Consideration.

— *A Pendulum, whose Length is 39 Inches and 2 Tenths English Measure, will perform one Vibration in one Second of Time.*] That Length (according to a Measure, that has been taken at a Mean from two Standard Measures of an *English Yard*, one of which is repositd at *Guildhall*, and the other at the *Exchequer*) is too much, 39,13 Inches, being the Length of a Second-Pendulum, which has been found, by comparing together many Experiments, none of which gave the Length above 39,133, nor less than 39,125. But I made use of 39,2 Inches, because it has been commonly reckon'd the true Length, till more accurate Experiments have been by Mr. *George Graham*, and others, within these 15 or 20 Years.

10. [68. Page 377. *The whole Curve so describ'd is call'd a Cycloid.*] Let the Line BC* represent Part of the Earth's Surface, on which a Wheel* Pl. 29. F. 1. or generating Circle rolling, does with a Point of its Circumference describe a Cycloid, as we have shewn; it is plain, that so small a Part of the Earth's Surface, which a Wheel runs over in one Revolution, may be consider'd as a Plane; and also, that if the generating Circle should run along the under Side of the Line CB, it would by a Point of its Circumference, first apply'd to one End as C, describe the same Sort of Curve, or Cycloid CGB, as if it had ran above the Line, which Curve would terminate at B, the Base of the Cycloid CB, being equal to the Circumference of the generating Circle, and the Curve CGB equal to 4 times the Diameter. But if the generating Circle should run along the inner Surface of a Sphere, as from the Point c to b, instead of running from e to K along a Tangent of equal Length with the Curve cAb, the Cycloid cGb will be shorter and less curve, than if its Base cAb had been strait, as one may see by comparing it with the Cycloid CGB, produced by the same generating Circle. This is call'd an Epicycloid. Now, if we suppose a large Wheel, or generating Circle, to be so big, that its Diameter shall bear a sensible Proportion to the Diameter of the Earth, its Revolution under the Earth's Surface would generate such a Cyloid. And if the generating Circle should be half as big as a Diameter of the Earth, as the Circle $\kappa\gamma$, the Point κ by the rolling of the generating Circle along the Hemisphere, or Semi-circle $\kappa A \beta$, would describe $\kappa \beta$ a Diameter of the Earth; so

* We have taken 16 Feet instead of $16,1$ to avoid Fractions.

that

Annotat. that the Cycloid would degenerate into a strait Line. Now, in all these
 Lect. V. Epicycloids, the Vibrations of a Pendulum, or of an heavy Body, oscil-
 lating on either Side of the middle Point G or γ , &c. will be *isochronal*.

Pl. 29. F. 1. For tho' we have shewn, that the Reason of a Body's falling faster in a
 Cycloid, than in an Arc of a Circle, or any other oblique Line, was be-
 cause the Body set out at first in the Direction in which Gravity acts; and
 here the Epicycloids being less curve than the common Cycloid, one
 would imagine, that they would lose that Advantage of Steepness at
 first: Yet we can shew, that in all the Epicycloids suppos'd within the
 Earth, as cGb , or even in that which becomes a strait Line, as $\kappa\beta$, Bo-
 dies begin their Motion in the same Direction as Gravity, whose Action
 tends towards the Center γ ; for wherever the generating Circle begins its
 Revolution, a Line as γc , drawn from the Center of the Earth (in which
 Gravity acts) passes through the Center of the generating Circle, and the
 describing Point: Whereas in the common Cycloid we build our Demon-
 strations upon the Supposition, that the Lines, in which Gravity acts, are
 parallel to one another, which Supposition is very proper, considering the
 great Distance of the Center of the Earth, whereto those Lines con-
 verge.

L. 1. An. 11. We have shewn in the 11th Annotation of *Lect. 1. Page 34 and 35*, as
 a Consequence of the Earth attracting according to its Quantity of Mat-
 ter; that the Force of the Action of Gravity is greatest at the Earth's
 Surface, decreasing as the Squares of the Distances from its Center in-
 crease, when Bodies are remov'd outwards farther and farther from the
 Earth's Surface; and that if Bodies are suppos'd to come nearer to the
 Center, within the Earth, the Force of Gravity acting upon them, will
 decrease directly as the Distance from the Center. If therefore (suppo-
 sing the Earth penetrable) several Bodies, as for Example 4, were placed,
 one at the Surface at β (about 4000 Miles from the Center γ) another at G
 (2500 Miles from it) another at H (2000 Miles from the Center) and the
 fourth at L (but 1000 Miles from it) the Force of Gravity acting upon
 those several Bodies to drive them to the Center, would be respectively as
 4000, 2500, 2000, and 1000; that is, as their Distances from γ ; and
 consequently, if they should set out at the same Moment of Time, they
 would also arrive at the Center at the same Instant. So, that if there
 was an Hole through the Earth along the Diameter of it as $\beta\kappa$, a Body
 falling from β , which, according to Mr. *Whiston's* Calculation, would come
 from the Surface to the Center in 21 Minutes and 9 Seconds, would
 go from β to κ in double the time, and so vibrate backwards and forwards
 in the Diameter $\beta\kappa$, like a Pendulum: And if it began its Motion at H ,
 it would take up as much time in going to I , vibrating backwards and for-
 wards (or rather up and down) in the Line HI . Now therefore, if the
*isochronal Vibration in a Diameter of the Earth, which follows from a Sup-
 position of the Attraction of the Earth, being proportional to its Quantity of
 Matter, be also a Consequence of the isochronal Vibrations of Pendulums in a
 Cycloid (which has been mathematically and experimentally demonstrated) it*
 must

must follow that that Supposition is true—tho' there are many other ways of proving it.

Annotat.
Lect. V.

11. [70. Page 373. *The Way to measure the least Alteration of Dimensions in Metals, so as to make it sensible, &c.*] Dr. Petrus van Muschenbroek, the ingenious Professor of Astronomy, &c. at Utrecht, contriv'd a Machine, which he calls a *Pyrometer*, for this Purpose, whose Description in his own Words, is as follows.

The 2d Figure of Plate 29, represents the whole Machine, with all its Parts together as it is us'd.

" A A A is an Iron turn'd up perpendicularly at one End, which Re- turn is $1\frac{8}{10}$ Inches high. The other End, which is distant from it $4\frac{3}{4}$ Inches, is also turn'd up, but then it is turn'd back again, so as to make a broader horizontal square Plate, the Side of which Square is 2 Inches. The Iron it self is 1 Inch wide and $\frac{3}{8}$ of an Inch thick, which Thickness was requir'd, lest it should easily or soon grow hot, which would hinder the Accuracy of the Experiments.

" Upon the Iron Plate stands a Brass Machine, which is drawn by it self in Fig. 3. where it is represented larger, and seen from another Side, the better to discover its Parts, which for that Purpose are mark'd with the same Letters as in Fig. 2. This is fix'd to the Iron by two Screws X, X, which are its Legs. D is a circular Plate of $2\frac{1}{10}$ Inches Diameter divided into 300 equal Parts, which we have not number'd in the Figure, by reason of their Smallness. This divided Plate stands upon four Pillars E, E, E, E, which join it to the lower Brass Plate: Between these two Plates there is a perpendicular Steel Arbor or Axis F, which has on its lower Part a Pinion of 6 Leaves or Teeth, and on its upper a Wheel of 60 Teeth mark'd G: There is also another Axis I H, supported by a Cock that comes down from the upper Plate, and which Axis with its upper End passes through the said upper Plate, so as to receive the Hand or Index I K; having at its lower End a Pinion of 6 Leaves to take the Teeth of the Wheel G. The Hand by one Turn of the Pinion H is carried round to all the Divisions. There is besides, a little Rack or streight Piece L with Teeth, which take the Leaves of the Pinion F, while the Rack slides along under two small Cocks P, P, where it is pressed towards the Pinion F by Help of two small Screws M, M, or drawn from it as there is Occasion, that the Teeth may take properly without sticking or being too loose. The Teeth of the Rack are 25 in an Inch, and as it moves forward and backward, the Pinion F is carried round, and consequently the Wheel G, which carries round the Pinion H, together with the Hand I K. Let us suppose the Rack to have run the Length of 1 Inch; then F and G will have turn'd round 4 times and $\frac{1}{8}$; and consequently the Pinion H, will have gone round $10 \times 4\frac{1}{8}$, that is, $41\frac{5}{8}$ times, because H turns round 10 times for G once:

I i i

" So

Annotat. " So that the Index I K, will have mov'd round $41\frac{2}{3}$ times, and because
 Lect. V. " the upper Plate is divided into 300 Parts, the Hand must have run over
 " 300 Divisions $41\frac{2}{3}$ times; that is, 12500 Parts. Therefore, when the
 " Hand goes but from one Division to another, the Rack moves but the
 " $\frac{1}{12500}$ th Part of an Inch. Now, since the Divisions are still large e-
 " nough to observe the Motion of the Hand to the Half of one of them, one
 " may perceive when the Rack has mov'd $\frac{1}{25000}$ Part of an Inch. Care
 " has been taken, that the Teeth and Pivots, should have but just Play
 " enough to go free, without which, one can make no accurate Experi-
 " ment. There is a female Screw goes through the Head of the Rack.

Pl. 29. F. 4. " The 4th Figure represents a square Bar, or Parallelopiped of Metal,
 " upon which the Experiment is made, $5\frac{8}{10}$ Inches long, and $\frac{3}{10}$ of an Inch
 " thick. One End of it O has a small Tail, that it may communicate no
 " sensible Degree of Heat to the Iron A A; it is received in a Notch
 " near B, and fix'd by the Screw C. Its other End N has an Hole in it,
 " through which Hole goes the Screw Q, which makes it fast to the
 " Rack L; but that End is also made small (as you see in the Figure)
 " lest the Heat should be communicated to the Rack.

" The Bar being thus fix'd, cannot become longer, without shooting
 " forward the Rack L, and thereby moving round the Hand I K, by
 " means of the Wheel and Pinions F, G, H; so likewise, when it becomes
 " shorter, they must move the contrary Way. Now, lest the Weight of
 " the Bar, should hinder the Motion of the Rack, I have thrust in a
 " Piece of a Watch Spring between the square Iron and Brass Plates E A,
 " exactly of such a Strength as to support the Bar as much as its Gravity
 " would depress it; so that the Rack would then move with all possible
 " Freedom. However insignificant this Particular may seem, yet it is
 " very necessary to be nice in it; for in making Experiments with Bars
 " of different Metals, I was oblig'd to draw out, or thrust in my Spring
 " more or less, according to their different Weights. The Hand, or In-
 " dex I K moves so freely, that you will perceive it to move when the Bar
 " of Tin is made fast to the Machine, if you but apply your warm
 " Hand to the Bar; for Tin requires but a very small Degree of Heat
 " to alter its Dimensions, as this Experiment proves. But to apply more
 " Heat conveniently, especially the Flame of burning Spirits, there is a

Pl. 29. F. 5. " Box made of Brass (see Fig. 5.) $3\frac{1}{2}$ Inches long, $1\frac{1}{10}$ Inch wide, and $\frac{4}{10}$
 " Inch deep, which is cover'd at Top with a Piece of blue Stone (such as
 " will bear the Fire, call'd in Dutch, *Lye*) which is represented in the Fi-
 " gure the underside upwards. It was necessary to make this Cover of
 " Stone, lest it should grow too hot, and set Fire to the Spirit within the
 " Lamp, as will happen when the Cover is made of Metal. It has a long
 " Hole cut through the Middle, into which is let in a Brass Plate T,
 " which has 5 small equidistant Holes $\frac{6}{100}$ Parts of an Inch asunder, and
 " $\frac{7}{100}$ Inch in Diameter, to transmit the Wicks of the Lamp. This
 " Lamp has 4 Feet, which closely take in the Iron A between them,
 " that

“ that in every Experiment the Flame may exactly come against the Mid- Annotat.
 “ dle of the Bar ; but the Bottom of the Lamp must not touch the Lect. V.
 “ Iron, which in such a Case would be heated, and by its lengthning
 “ would disturb the Experiment: And even thus, in some Experiments I
 “ have found it a little warm’d. The Distance between the Bottom of
 “ the Bar and the upper Surface of the Lamp, must be half an Inch,
 “ that the Cotton Wicks that stand up $\frac{3}{10}$ of an Inch, may give an exact
 “ and equal Flame. The Cotton Threads must be very fine and even,
 “ and 5 of them twisted together must make a Wick of about $\frac{5}{100}$ of an
 “ Inch in Diameter. I took particular Care of every Circumstance in making
 “ Experiments with this Machine; because the least Omission might
 “ lead one into great Errors: For if the Wick is drawn up too high thro’
 “ the Cover of the Lamp, the Flame will be too large, which shews the
 “ Necessity of keeping it to the same Height; Likewise in Experiments,
 “ that are made with highly rectified Spirit of Wine, if there was not
 “ always an equal Quantity put into the Lamp, the Flame would be
 “ supply’d too fast or too slow, and so the Heat would vary; therefore
 “ I took Care always to make my Experiments with the Lamp half full
 “ of Spirits, whereby I had the best Flames, which were of a cylindrick
 “ Form, from the Lamp to the Metal which they heated; only they
 “ widen’d a little at Top—the Diameter of their Base was $\frac{13}{100}$ Parts of
 “ an Inch. And to prevent the Motion of the Air, or the Breath of
 “ one’s Mouth from moving the Flames, I cover’d the whole Instrument
 “ with a Glass; except just the Dial Plate, which came up above the
 “ Glass the better to observe the Index.

“ This being all ready, I proceeded to examine how much, Iron,
 “ Steel, Copper, Brass, Tin, and Lead were expanded by one Flame;
 “ then how much they were expanded by Two; then by Three; then by
 “ Four; and lastly, by all the five Flames: Then also, whether there
 “ was any Difference when two Flames burn’d next to each other, or two
 “ more distant Flames.

“ One Day when it began to freeze, and Farenheit’s Thermometer was
 “ at 32 Degrees, the Wind in the West, the Sky clouded, and the Mer-
 “ cury in the Barometer stood at 29 $\frac{4}{12}$, I laid the Metals above-mentio-
 “ ned, close to one another, upon a Stone, that they might be cool’d
 “ equally: Then I apply’d them successively to the *Pyrometer*, and having
 “ first lighted one Flame, I observ’d their Expansion: Then having taken
 “ them off from the *Pyrometer*, I set them a cooling till they became as cold
 “ as at first, and try’d them on the *Pyrometer* with two Flames; and so
 “ on till I tried the Effect of all the five Flames upon them. For Brevi-
 “ ty’s sake, I have digested into the following Table what those Effects
 “ were. The Degrees of Expansion are mark’d in Parts, of which each
 “ is the $\frac{1}{12,000}$ th Part of an Inch. It is to be observ’d of Tin, that it
 “ will easily melt, when heated by two of the Flames burning next to
 “ one another, therefore one cannot try it with more Flames than two.

Annotat. "Lead commonly melts with three Flames, which are next to one another, provided they burn long.
 Lect. V. "

" A T A B L E ;

" *Shewing how Metals are expanded by the Heat of Flames*
 " *of Spirit of Wine, different in Number, but equal in*
 " *Bignefs.*

| <i>Expansion by one Flame in the Middle.</i> | Iron | Steel | Copper | Brass | Tin | Lead |
|--|------|-------|--------|-------|-----|------|
| | 80 | 85 | 89 | 110 | 153 | 155 |
| <i>By two Flames in the Middle next to one another.</i> | 117 | 123 | 115 | 220 | | 274 |
| <i>By two Flames $2\frac{1}{2}$ distant from one another.</i> | 109 | 94 | 92 | 141 | 219 | 263 |
| <i>By three Flames in the Middle next to one another.</i> | 142 | 168 | 193 | 275 | | |
| <i>By four Flames in the Middle next to one another.</i> | 211 | 270 | 270 | 361 | | |
| <i>By all the five Flames.</i> | 230 | 310 | 310 | 377 | | |

" These Experiments were made several times over ; and tho' they did
 " not always agree, yet they never differ'd so much as five Degrees,
 " which is so little as not to be worth Notice; and this may be owing
 " to the Play of the Teeth, which cannot be avoided in these Experi-
 " ments. But I always took the Mean of a great Number of Experi-
 " ments.

" From

“ From these Experiments it appears at first View, that Iron is the
 “ least rarefied of any of these Metals, whether it be heated with one or
 “ more Flames: And therefore is the most proper for making such Machines
 “ or Instruments, which we would have to be least alter’d by Heat and
 “ Cold. Annotat.
Lect. V.

“ Therefore it is very proper to make the Rods of Pendulums for
 “ Clocks, of Iron: They are not so good of Steel; and much worse of
 “ Brass; yet sometimes they are made of Copper, because it is not so
 “ liable to rust; but yet, that is wrong. So likewise the Measures of
 “ Yards or Feet should generally be made of Iron, that their Length may
 “ be as near as possible the same, Summer and Winter.

“ 2dly, The Expansion of Lead and Tin by only one Flame, is pretty
 “ near the same.

“ 3dly, The same Flame expands Tin and Lead almost the double of
 “ Iron; for their Expansions are as 155 to 80; that is, nearly as 2
 “ to 1.

“ 4thly, The Flames which are next to one another, and act upon the
 “ metalline Rods, cause a greater Rarefaction, than when they have a
 “ sensible Interval between them: For two neighbouring Flames expand-
 “ ed Iron 117 Degrees, and two farther asunder, but 109; and the same
 “ was observable in all the other Metals.

“ Now, this happens, because all the Parts of the Fire do not rise di-
 “ rectly up and apply themselves to the Rods over them; but some fly
 “ off sidewise from every Part of the Flame. Now, when two Flames
 “ are at a Distance from one another, the lateral fiery Parts fly off from
 “ the Flames, and do not act upon the Metals; whereas, when they are
 “ near together, those Parts which fly off from the Side of one Flame
 “ towards the other, are reflected back from that Flame, as from a *Spe-*
 “ *culum*, and coming back into their own Flame again, are driven up-
 “ wards, and enter into the Metal, which thereby receiving a greater
 “ Quantity of Fire, must consequently swell the more.

“ 5thly, Now, let us compare together the Expansions of the same
 “ Metal, produc’d by one, two, three, or more Flames: Two Flames
 “ did not give double the Expansion of one; nor three Flames three
 “ times the Expansion, but less; and the Expansions differ so much the
 “ more from the *Ratio* of the Number of the Flames, as there are more
 “ Flames acting at the same time. I shall hereafter give the Demonstrati-
 “ on of that *Phænomenon*; now I shall only give the Proportions of the
 “ Expansions observ’d, in the following short Table.

| Annotat. | In Iron. | In Steel. | In Copper. | In Brass. | In Lead. |
|----------|---|---|---|--|--|
| Lect. V. | 80:117 :: 1:1 $\frac{1}{8}$ $\frac{7}{10}$ | 85:123 :: 1:1 $\frac{3}{8}$ $\frac{3}{5}$ | 89:155 :: 1:1 $\frac{5}{8}$ $\frac{9}{10}$ | 110:200 :: 1:1 $\frac{2}{1}$ $\frac{0}{10}$ | 153:263 :: 1:1 $\frac{1}{1}$ $\frac{1}{10}$ $\frac{3}{10}$ |
| | 80:142 :: 1:1 $\frac{1}{8}$ $\frac{1}{10}$ $\frac{1}{10}$ | 85:168 :: 1:1 $\frac{1}{8}$ $\frac{3}{10}$ $\frac{3}{10}$ | 89:193 :: 1:2 $\frac{1}{8}$ $\frac{3}{10}$ $\frac{3}{10}$ | 110:275 :: 1:2 $\frac{1}{1}$ $\frac{1}{10}$ $\frac{5}{10}$ | |
| | 80:211 :: 1:2 $\frac{1}{8}$ $\frac{1}{10}$ $\frac{1}{10}$ | 85:270 :: 1:3 $\frac{1}{8}$ $\frac{1}{10}$ $\frac{1}{10}$ | 89:270 :: 1:3 $\frac{1}{8}$ $\frac{1}{10}$ $\frac{1}{10}$ | 110:361 :: 1:3 $\frac{1}{1}$ $\frac{1}{10}$ $\frac{1}{10}$ | |
| | 80:230 :: 1:2 $\frac{3}{8}$ $\frac{0}{10}$ $\frac{0}{10}$ | 85:310 :: 1:3 $\frac{1}{8}$ $\frac{1}{10}$ $\frac{1}{10}$ | 89:310 :: 1:3 $\frac{1}{8}$ $\frac{1}{10}$ $\frac{1}{10}$ | 110:377 :: 1:3 $\frac{1}{1}$ $\frac{4}{10}$ $\frac{7}{10}$ | |

“ 6thly, Before Metals are, from the same Degree of Cold, brought to melt, they are not equally expanded; but some are more expanded, and some less. For Tin began to run when it was rarefied 219 Degrees: But Brass, which was expanded 377 Degrees, was far from being red hot, and consequently very far from melting; and Copper was expanded 310 Degrees, whereas perhaps it must have been expanded twice as much before it could melt.

So much translated from Dr. Muschenbroek, * serves to shew some of the many Uses of the *Pyrometer*; which I have improv'd by the following Changes.

1st, Instead of square Rods of Metal, to make the Experiments upon, I use cylindrick ones; because I can more certainly have them of the same Size, by drawing them through the same Hole of a strong Steel Plate such as Wire-drawers use: And by that means, I am sure, that the Bulks of the Metals compar'd are equal.

2dly, Instead of the Pinion F, I have a small Roller of Steel truly round, but not polish'd, and filed on the Surface in the Direction of its Axis, so that it may be consider'd as a little Wheel with an infinite Number of Teeth: Then the Wheel G upon the same Axis has no Teeth, but only a Groove to receive a fine Watch-chain (or even an Horse-hair) whereby it carries round a Roller at H with a little Groove, upon the Axis of which Roller the Index is fix'd at the upper End I.

3dly, That the Chain whereby G carries H, may be neither too slack, nor too tight, the whole Dial-Plate (which in my *Pyrometer* is square) and the Cock and Pinion H are driven towards, or from the Wheel G, by a Screw fasten'd to the upper Frame-Plate, which Plate receives the Top of the Pillars and Arbor of G, whilst the Dial-Plate slides upon it. N. B. There is no such Plate under the Dial-Plate in Dr. Muschenbroek's *Pyrometer*.

Pl. 29. F. 2. 4thly, Instead of the Rack N L (Fig. 2.) I have a long thin Plate of Steel about $\frac{1}{100}$ Parts of an Inch broad, rough filed, so as to move the first Roller F by rubbing against it. It is temper'd of a Spring Temper, and a little curv'd, so as to have its convex Side bear against F; but

* See the Notes of his Latin Translation of the *Experiments of the Academia del Cimento*, Part the 2d, from Page 12 to Page 18, &c.

when

when it is fasten'd to the Rod at N, there is a Spring fix'd to the lower Annotat. Brass Plate, which draws it streight and tight by its End L in the Direction N L, and instead of the Cocks P, P (*Fig. 3.*) there are two Pullies placed horizontally, whose broad vertical Grooves receive and direct the Steel Plate or Ruler, that supplies the Place of the Rack. Leçt. V.
 Pl. 29. F. 3.

5thly, Instead of the Watch Spring to support the Bars, I have a small Brass Roller of $\frac{1}{10}$ of an Inch Diameter, whose Axis is horizontal; and this Roller is rais'd up from its Pedestal, that is fix'd to the Iron at W by a Screw, so as to come under every Rod of Metal, and support it at its End N.

The 6th Figure represents the lower Frame Plate, with my Alterations. Pl. 29. F. 6.

BN is the round Rod of Metal to be tried, with the Steel Plate made fast at N by the Pin Q. This Steel Plate, whose natural Situation is represented by the pointed Line N L, is here kept in the strait Line N L, by means of the Spring S L, which hooks in the End of it at L, drawing it in the Direction N L; and it is directed by the Grooves of the Pullies P, P, so that its Side, that is inclin'd to be convex when left at Liberty, now presses on the Roller H, which it turns by its Friction, as it accedes towards, or recedes from L. The pointed Circle G g represents the Wheel above on the same Axis as H: g i G the Watch-Chain, carrying round the last Roller i, and Index u i k, as in Dr. *Muschenbroek's Pyrometer*. The rest is easily comprehended from what has been said, and a Sight of the 7th Figure, which represents the support-Pl. 29. F. 7. ing Brass Roller, rais'd and depress'd, by turning round the screw'd Plate p p.

N. B. *There is no Inconvenience in using the Horse-hair if you take care to place its Knot at G; for as the Motion is made in the Direction G m g, the Knot will never go so far as g, and therefore in all its Motion bear equally in the Groove of the Wheel G.*

By using Rollers, the Shake of Teeth is wholly avoided in my *Pyrometer*; so that the Index begins to move the very Moment, that Heat is apply'd to the metalline Rods; and if you blow out the Flame, that very Instant the Index begins to return, which cannot happen where there are Teeth.

I mean the quick Return when the Heat is remov'd. I must own, that if we consider the Motion of the Index only one Way, when once the Teeth take, and it begins to move, it will go on regularly; therefore Dr. *Muschenbroek's Experiments* may be depended upon. However, I shall try them again with my *Pyrometer* the first Leisure. I must also do the Doctor justice, to own, that he has signified to me in one of his Letters,

Annotat. Letters, that if he had not lost his Workman, he would have made a
Lect. V. new *Pyrometer* without Teeth.



12. [73. *Who assert, that the Earth is an oblong Spheroid (like an Egg) higher at the Poles than the Æquator, &c.*] Monsieur *Cassini* says, that the Earth is an oblong Spheroid, higher at the Poles than the Æquator, making the Axis longer than a Diameter of the Æquator about 13 *French Leagues*, which he deduces from comparing his Father's Measures of the Meridian from *Paris* to the *Pyrenean Mountains* with those of *Monf. Picard*, of which an Account may be seen in the *Memoirs* of the R. Academy for 1713; then having afterwards continued the Meridian, which is drawn through *France*, from *Paris* to *Dunkirk*, he still draws Consequences to prove the Earth an oblong Spheroid; but then makes the Axis exceed the æquatorial Diameter $3\frac{1}{4}$ Leagues.

Now, in order to prove this, M. *Cassini*, taking the Measures above-mentioned to be exact enough, not only to determine the Magnitude of a Degree of the Earth, corresponding with a Degree of a great Circle of the Heavens; but also to shew, even the Difference in the Degrees of the Earth (reckoning those, that were measur'd in the South of *France*, to exceed those towards the North, by a certain Number of Fathoms and Feet) demonstrates, that if the Degrees of the Earth are longer towards the Æquator than the Poles, the Plane of the Meridian must be an Ellipse, whose long Axis is that of the Earth.

Monsieur *Cassini's* Demonstration is certainly right; but the Measures taken could not, in the Nature of Things, determine any thing about the different Length of Degrees so near to one another; because as that Difference in one of the Suppositions is but of 11 or 12 Fathoms, and in another of about 31 at most, the Latitude must be taken to an Exactness, exceeding to the Nature of any Astronomical Instruments yet made, and much more than that of the Instruments made Use of by the Gentlemen who carried on the Meridian.

For in the first place, the Instrument, with which they took the Observations for the Latitude at the two Ends of the Meridian, was a 10 Foot Sector, where the 200th Part of an Inch answers to 8 Seconds of a Degree. Now the 200th Part of an Inch, being one of the least visible Parts that we can see in a divided Line, they could not take an Angle nearer than that; nay, their Instrument, according to their own Description of it, was divided but to every 20 Seconds. They allow, that 16 Toises (or Fathoms) upon the Surface of the Earth, answer to 1 Second in the Heavens; and they don't pretend to have taken an Observation nearer than to about 3 Seconds, which therefore cannot determine a Difference less than 48 Toises; whereas the Degrees are only suppos'd to decrease (at most) 31 Toises each from *Collioure* (the Southern Part of their Meridian) to *Dunkirk*; but an Error of 8 Seconds would make a Difference of 128 Toises on the Surface of the Earth, above 10 times greater than the Difference of the Degrees in the first Supposition, and
4 times

PLA. I.










Plate II.








Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.




Fig. 8.




Fig. 7.




Fig. 5.




Fig. 6.














Fig. 9.























Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.




Fig. 6.




Fig. 7.




Fig. 8.




Fig. 9.




Fig. 10.




Fig. 1.



Fig. 2.




Fig. 3.




Fig. 4.




Fig. 5.




Fig. 6.




Fig. 7.




Fig. 8.




Fig. 9.



Fig. 10.




Fig. 11.




Fig. 12.



Fig. 13.




Fig. 14.




Fig. 15.



Fig. 16.



Fig. 17.



Fig. 18.




Fig. 19.



Fig. 20.



Fig. 21.



Fig. 22.



Fig. 23.



Fig. 24.



Fig. 25.



Fig. 26.



Fig. 27.



Fig. 28.




Fig. 29.




Fig. 30.




Fig. 31.




Fig. 32.




Fig. 33.




Fig. 34.




Fig. 35.




Fig. 36.




Fig. 37.




Fig. 38.



Fig. 39.




Fig. 40.



Fig. 41.



Fig. 42.



Fig. 43.



Fig. 44.



Fig. 45.



Fig. 46.



Fig. 47.



Fig. 48.



Fig. 49.



Fig. 50.



Fig. 51.




Fig. 52.




Fig. 53.



Fig. 54.




Fig. 55.




Fig. 56.




Fig. 57.



Fig. 58.




Fig. 59.













Fig. 60.
















Fig. 1.




Fig. 2



Fig. 7 Page 454



Fig. 5.




Fig. 3.




Fig. 8.
Page 454




Fig. 6.





Fig. 9.
Page 455.



Fig. 4.






Fig. 2




Fig. 1.




Fig. 4.




Fig. 3.




Fig. 6.




Fig. 5.




Fig. 8.




Fig. 7.




Fig. 13.




Fig. 11.




Fig. 12.





Fig. 10.






Fig. 2.




Fig. 6.




Fig. 1.




Fig. 8.




Fig. 10.




Fig. 4.




Fig. 5.




Fig. 3.




Fig. 7.




Fig. 11.




Fig. 12.




Fig. 13.




Fig. 14.




Fig. 15.




Fig. 16.




Fig. 17.




Fig. 18.




Fig. 19.




Fig. 20.




Fig. 21.




Fig. 22.




Fig. 23.




Fig. 24.




Fig. 25.




Fig. 26.




Fig. 27.




Fig. 28.




Fig. 29.




Fig. 30.




Fig. 31.




Fig. 32.




Fig. 33.




Fig. 34.




Fig. 35.




Fig. 36.




Fig. 37.




Fig. 38.




Fig. 39.




Fig. 40.




Fig. 41.




Fig. 42.




Fig. 43.




Fig. 44.




Fig. 45.




Fig. 46.




Fig. 47.




Fig. 48.




Fig. 49.




Fig. 50.




Fig. 51.




Fig. 52.




Fig. 53.




Fig. 54.




Fig. 55.




Fig. 56.




Fig. 57.




Fig. 58.




Fig. 59.




Fig. 60.




Fig. 61.




Fig. 62.




Fig. 63.




Fig. 64.




Fig. 65.




Fig. 66.




Fig. 67.




Fig. 68.




Fig. 69.




Fig. 70.




Fig. 71.




Fig. 72.




Fig. 73.




Fig. 74.




Fig. 75.




Fig. 76.




Fig. 77.




Fig. 78.




Fig. 79.




Fig. 80.




Fig. 81.




Fig. 82.




Fig. 83.




Fig. 84.




Fig. 85.




Fig. 86.




Fig. 87.




Fig. 88.




Fig. 89.




Fig. 90.




Fig. 91.




Fig. 92.




Fig. 93.




Fig. 94.




Fig. 95.




Fig. 96.




Fig. 97.




Fig. 98.




Fig. 99.




Fig. 100.

Fig. 4.




Fig. 3.




Fig. 1.




Fig. 5.




Fig. 6.




Fig. 9.




Fig. 7.




Fig. 8.




Fig. 2.




Fig. 12.




Fig. 10.




Fig. 11.





Fig. 13.








Fig. 2.





Fig. 7.




Fig. 8.




Fig. 9.





Fig. 10.






Fig. 4.









Fig. 7.



4 times greater than that Difference in the last. Besides, the Latitude was not observ'd in the intermediate Places between *Paris* and *Collioure*, with the above-mention'd Instrument of 10 Foot Radius; but they made Use of a Quadrant, whose Radius was only 39 Inches, and sometimes an Octant of 3 Foot Radius. Nay, they say themselves in their Account, that it is not the Observations made at the Ends of the Meridian, that we are to deduce the Difference of the Length of a Degree from, but the Altitudes taken at several Places between the Extremes; and if we grant that they can take an Angle very well to 4 or 5 Seconds with the greater Instrument, they cannot come nearer than 12 or 15 Seconds with the Quadrant or Octant, which we must depend upon for the Difference of the Measure of Degrees. So that upon the whole, we are to determine a Length of 31 Toises, by an Instrument, which is liable to err above 200.

And indeed, upon a strict Examination of the Account of the Measures of Degrees, I found many Errors and Inconsistencies; particularly in the measuring of the Height of Mountains, where 9 Observations of the Barometer for measuring the Height of them in the South of *France*, which are said to confirm their trigonometrical Measures, so entirely clash with one another, as to determine nothing at all. Nay, the Conclusion, that Degrees on the Earth increase in going towards the *Æquator*, is only drawn from some pick'd Observations; for if they are all compar'd together, the Degrees will appear from thence to increase in going Northwards.

Now, abstracting from Observations, the ingenious *Mons. Mairan* has endeavour'd to prove, that in an oblong Spheroid, Pendulums must be made shorter to swing Seconds at the *Æquator*, than in greater Latitudes; but the Principles he has assum'd, are not to be allow'd him; and if they were, it would follow from them, that a Pendulum swinging Seconds at *Paris*, must be shorten'd an Inch to make it swing Seconds at the *Æquator*; whereas the Fact is, that such Pendulums carried from *Paris* to the *Æquator*, are only shorten'd $\frac{1}{10}$ of an Inch. Therefore, *M. Mairan*, by proving a great deal too much, proves nothing in this Case. Those that are curious to know all, that relates to this Dispute, may consult the *Philosophical Transactions*, No. 386, 387, 388† and 389, in which, I think, I have fully prov'd the oblate spheroidal Figure of the Earth, according to *Sir Isaac Newton*, and shewn the oblong Spheroid to be impossible.

† *N. B.* There are some *Errata* in my second Dissertation, which are corrected in the last Abridgement of the *Transactions*, by *Messieurs Reid* and *Gray*.

F I N I S.

K k k



At the Desire of several Friends, I have here subjoin'd a short Description of my Planetarium, an Instrument, which I made lately, to shew the Motion of the heavenly Bodies.

Description of the PLANETARIUM.

MACHINES and Movements for representing the Motions and Appearances of the heavenly Bodies, have been justly esteem'd in all Ages; especially, since the *Copernican* System has been generally receiv'd; not only as the most probable Hypothesis; but as Sir *Isaac Newton* has, from the Laws of Gravity, prov'd it the true System of the World* — at least, a System of so much of the Universe as relates to us Inhabitants of the Earth; that is, so far as we can discover, not only by our Eyes, but by means of Telescopes.

By Help of such Machines, a great many Persons, who have not time to apply themselves to the Study of Astronomy, and yet are desirous to be acquainted with the celestial Appearances, in a few Days may get a competent Knowledge of several Phenomena, and especially, be cur'd of the common Prejudices against the Motion of the Earth.

When such Astronomers, as are skill'd in Mechanics, have made or contriv'd these kind of Movements, they have answer'd the Ends propos'd, by shewing so much as the Machine was intended to represent; but, when Instrument-makers, without a competent Knowledge of Astronomy, or proper Instructions from Persons of Skill, have made improper Additions to such Machines as had been contriv'd by Astronomers (under pretence of great Improvements) merely to make them pompous and costly; the true Intention of the first Inventors has been destroy'd, and the Buyers have paid dear for false Notions of Astronomy.

Mr. *George Graham* (if I am rightly inform'd) was the first Person in England, who made a Movement to shew the Motion of the Moon round


* We must not forget, that the Rev. and ingenious Mr. *James Bradley*, Savilian Professor of Astronomy at Oxford, has demonstrated it also of late from Astronomical Observations.

Diameters & Quantity of Matter
767

Plate 30.

Sollt

'Periods' and Distances:



Saturn's Ring is distant on every side from Saturn's body 21,000 Miles, and is as much in Breadth.

I Sener sculpt!

the Earth, and of the Earth and Moon round the Sun, about 25 or 30 Years ago. In this Machine every thing, that was shewn, was well and properly executed——as the Phænomena of Day and Night, and their gradual Increase and Decrease, according to the Seasons, the Places of the Earth where the Sun is successively vertical, and seems to describe its Parallels, the real annual Motion of the Earth, which gives the Sun an apparent annual Motion, the Rotation of the Sun about its Axis, the periodical and synodical Month, the solar and sydereal Days, the successive Illumination of all the Parts of the Moon, &c. This Machine being in the Hands of an Instrument-maker, to be sent with some of his own Instruments to Prince *Eugene*, he copied it, and made the first for the late Earl of *Orrery*, and then several others, with Additions of his own. Sir *Richard Steele*, who knew nothing of Mr. *Graham's* Machine, in one of his Lucubrations, thinking to do justice to the first Encourager, as well as to the Inventor, of such a curious Instrument, call'd it an *Orrery*, and gave Mr. *J. Rowley* the Praise due to Mr. *Graham*.

Since that Time, the *Orreries* have been much in vogue, and executed in the most ornamental Manner, first by Mr. *Rowley*, then by other Instrument-makers; but the Addition of the rest of the Planets and the Satellites, reckon'd by the Ignorant to be an Improvement, does but give confus'd Ideas, and false, as to Distances and Proportions of Bigness; which will always be so, whilst the Orbit of the Moon about the Earth, is fix'd to the same Machine, that has the primary and secondary Planets. The general System (in which our Earth has but a small Part) ought to be by it self——The Sun, Moon, and Earth, ought also to be shewn by themselves; — and a System of a primary Planet (as Jupiter, for Example) with its Satellites in their true Proportion of Magnitudes and Distances, should also be represented by it self.

These Considerations (and the Desire of giving a true Notion of the celestial *Phænomena*, in the plainest and most expeditious manner, to such Persons as do me the Honour to frequent my Courses of Experimental Philosophy) induc'd me to contrive and make a Movement with all the Requisites above-mentioned, that might shew in true Proportion what can be so represented in a celestial Machine.

I have made it to consist of several Parts to be put on and taken off successively; and have call'd it a PLANETARIUM, whose Description I am now going to give: But before I enter upon it,

I must desire the Reader to cast his Eye upon the 30th Plate, which is Mr. *Whiston's* Scheme of the *Solar System* epitomiz'd, whereby he will make himself acquainted with the mean Distances, Magnitudes, Periods, Quantities of Matter, Revolutions about the Axes, Densities, and mean Velocities in their Orbits, of all the primary and secondary Planets; and the Orbits of several Comets, the Periods of three of them. Only I must observe some Things relating to this Scheme, which (if not consider'd) might lead into Error.

1. Tho' we see the Distance of the Planets from the Sun set down in Miles, we are not to imagine, that we can measure their Distances to a few Miles, as we really can the Moon's Distance; because the Semi-diameter of the Earth (whose Quantity is known in Miles) being about $\frac{1}{60}$ th Part of the Moon's Distance, is a sufficient Measure for the Moon's Distance; whereas it is scarce a twenty thousandth Part of the Sun's Distance, therefore very insufficient for a Measure of that Distance. There are indeed other Methods of coming at the Sun's Distance, as by the Parallax of Mars, but none of them can ascertain it to be a Million of Miles. We must stay till the next Transit of Venus across the Sun's Disk in the Year 1761, May the 26th, when we may, by observing that Transit, find out the Sun's Distance within a 500th Part of the whole Distance, as Dr. Halley has shewn in the *Philosophical Transactions*, No. 348. p. 454. and even that Advantage will bring us but within 15 or 16000 Miles of the Truth. However, we know the proportionable Bigness and Distances of the Bodies revolving round the Sun; that is, if the Sun's Distance from the Earth be, as here set down, 81 Millions of Miles, all the other Distances and Diameters are right: We know those Distances in Diameters of the Sun, which is the Way that most Astronomers express them; for whenever that comes to be known, all the rest will be known too; and therefore their respective Proportions may be represented by Schemes and Machines: But not exactly what Proportion they bear to the Moon and Earth.

2. Venus is here set down, as turning about her Axis in 23 Hours; but since Mr. Whiston publish'd his Scheme, Signior Bianchini has observ'd, that she is 24 Days and 8 Hours revolving round her Axis, which Axis is inclin'd to the Plane of her Orbit in an Angle of 15 Degrees.

3. The Moon's Velocity in her Orbit, which is the only Planet, whose real Velocity we know, has been omitted here. She moves 2300 Miles in an Hour. The other Planets only move in an Hour the Miles set down, supposing the Sun at the Distance of 81 Millions of Miles from the Earth.

To give a more easy Idea of the Distances and Diameters of the Bodies, we will take them at a Mean in round Numbers; and supposing the Distance from the Earth to the Sun, to be divided into 10 Parts, Mercury will be distant from the Sun 4 of those Parts, Venus 7, Mars 15, Jupiter 52, and Saturn 95. If we call the Sun's Diameter 100; that of $\frac{1}{2}$ will be $7\frac{1}{10}$; of $\frac{1}{4}$ 10; of $\frac{1}{8}$ $12\frac{1}{2}$; of $\frac{1}{16}$ 15; of $\frac{1}{32}$ 18; of $\frac{1}{64}$ 21; and of the Moon $28\frac{1}{2}$; and the Satellites of $\frac{1}{4}$ and $\frac{1}{2}$ are conjectur'd to be about as big as our Earth.

4. Tho'



4. Tho' the different Curvature of the Parts of the Ellipses, in which the Planets move, is not sensible enough to be express'd even in a larger Scheme; yet the Excentricities might have been express'd; but that it was omitted to avoid Confusion among the Comets Orbits. The Excentricities of the Planets may be express'd by supposing the Line of the mean Distance of each, divided into 1000 equal Parts, and then the Excentricities will be the following Numbers of those Parts.

| | | Equal Parts | | | | | |
|---------------------------------|-----------|--------------|------|-----------------------|-----|----------------|---|
| Imagine the mean
Distance of | Saturn | divided into | 1000 | their Excentricity is | 55 | of those Parts | The Moon's
Excentricity
is $\frac{111}{1000}$ Parts
of its Distance
from the Earth. |
| | Jupiter | | 1000 | | 48 | | |
| | Mars | | 1000 | | 93 | | |
| | the Earth | | 1000 | | 17 | | |
| | Venus | | 1000 | | 7 | | |
| | Mercury | | 1000 | | 210 | | |

The PLANETARIUM, (Plate 31.)

Is fix'd in a Frame of Ebony about 6 Inches High, and 3 Foot in Diameter, contain'd by twelve vertical Planes, on which are represented the 12 Signs of the Zodiac. The upper Surface is flat of polish'd Brass, on whose outward Circumference are screw'd in six Brass Pillars, which support a large flat silver'd Ring representing the Ecliptick, with several Circles drawn upon it. The three innermost are divided into 12 Parts for the Signs of the Zodiac, each of which is divided into 30 Degrees; and among those Degrees are grav'd in their proper Places, the *Nodes*, *Aphelia*, and greatest North and South Latitudes of the Planets. Between the next two Circles are mark'd the Cardinal Points. The next three Circles have the Months and Days of the Month, according to the *Julian* Account; and the three last have them likewise engrav'd, according to the *Gregorian* Account. But in the 30th Plate I have omitted the *Julian* Account, as I have omitted the *Gregorian* in the 31st Plate. Upon the Brass Surface of the Machine, are graduated Silver Circles, which carry the Planets (represented by silver'd Balls) upon Arbors or Stems, that raise them up to the Height of the Plane of the Ecliptick; and by turning about the Handle or Winch of the *Planetarium*, all the Planets move at their proportional Distances from a little gilt Ball in the middle, which represents the Sun, and perform their Revolutions according to their periodical Times. There are fix'd Indices of blued Steel, which shew the Longitudes of the Planets, by pointing to the Divisions of the silver'd Rings or Circles, as they move round. But as these Circles, being concentrick, give only the mean Distances; the true Orbits, according to their Excentricities, are grav'd on the Outside of each Circle, with the periodical Times taken from the Tables, to shew what the Revolutions are nearer than can be perform'd.

perform'd by any Machine. The *Nodes* and *Aphelia*, with the Places of greatest North and South Latitudes, are also mark'd on those Orbits. As the Distances are in their true Proportions to each other; so likewise are the Bodies of the Planets in their just Proportions to one another. But it cannot be expected, that the Diameters of the Planets should be in Proportion to the Diameters of the Orbits; because, taking Jupiter under 3 Inches Diameter, and the Earth a little more than a quarter of an Inch, as in my Machine, it would require the System to be of the Bigness of a Mile and $\frac{1}{4}$, the Orbit of Saturn about 9000 Feet in Diameter, and so of the rest, which would make the Machine 3000 times bigger than it is. And if we would suit the Bodies to the Machine as it is, the Bodies must be 3000 times less, which would render them all invisible, but the Sun; and that would be less than $\frac{1}{1000}$ th Part of an Inch. For this Reason, I could not put in a Ball big enough to represent the Sun in respect of my Planets; but we are to suppose the Sun (in respect of them) as big as the inner Circle of the Silver Ring, which represents the *Ecliptick*.

Now in the ORRERIES (as they call them) there is shewn no Proportion of the Orbits to one another; nor of the Bodies to one another.

As the Orbit of the Moon, and the Orbits of Jupiter and Saturn, are quite lost in this Proportion of the Orbits of the primary Planets, much more are the *Satellites* themselves; therefore the *Satellites* are not put on in this Position of the Machine. But Saturn's Ring is join'd to Saturn's Body, according to its Proportion, and the Inclination of its Plane to the Plane of Saturn's Orbit: And as the Planet is carried round, the Ring always moves parallel to it self, as it does in the Heavens. Thereby we see why the Inhabitants of the Earth, in one Revolution of Saturn, see the Ring twice in the most open Situation of the *Anse*, as in *Figure 1*,* and twice, as if it had no Ring, that is, when the Edge of the Ring is towards the Earth (the Plane of the Ring going through the Observer's Eye) and the successive increasing and decreasing of the visible Bigness of the *Anse*.

* Pl. 31. F. 1.

The 2d Figure represents Jupiter with his Belts, and the Spots whereby his Revolution has been observ'd.

When the Spectators have look'd long enough upon the Machine, to have a compleat Idea of the proportionable Bigness of the Planets; Jupiter and Saturn are taken off, and then there are put on another Jupiter and another Saturn 3 times less than the former, in order to put *Satellites* about them (and at the same time the Moon is joined to the Earth) and shew how the *Satellites* accompany their primary Planet in its Course round the Sun. These *Satellites*, which are Pearls upon crooked Stems, do not turn by Clock-work round their Primaries (as has been done in some of the large *Orreries*) but are only set by the Hand; because, to do it, would be only a needless Expence, to give a false Notion of their Bigness, Distances, and Inclination of their Orbits, in respect of their Primaries.

ries. But to give a right Notion of Jupiter and his Satellites, and Saturn and his Satellites; there is shewn for each of these Planets a System a-part, where the Distances from the Primary, and the Bigness of the Satellites, are express'd: And in this System, tho' Jupiter is but of about an Inch Diameter, the outermost Satellite is as far distant from Jupiter's Center, as Saturn is from the Sun in the Machine, which shews the Inconsistency and Disproportion of making the Satellites to move round Jupiter in an *Orrery*. Saturn's Satellites are still more improperly put in; because four of them move in Orbits very much inclin'd to Saturn's Ecliptick, (*viz.* in an Angle of above 30 Degrees) and the 5th has its Orbit almost in the same Plane as Saturn's Ecliptick, with a Diameter greater than the Diameter of the whole *Planetarium*, even when Saturn is 3 times less than the Saturn of the *Planetarium*.

The next Thing put on, is a Contrivance to shew, that all the Confusion of the Planets Motions in the *Ptolemaick Hypothesis* (call'd their Stations and Retrogradations) is not really, but apparently so, in the *Copernican*, or true System of the World. And this is done by two Steel Indices, one of which being always applied to the Sun, and successively to the Top of the Stem of the Planet to be examin'd, whilst the other is apply'd to the Earth (as a Center) and the said Planet; by turning the Handle of the Machine, the *Heliocentrick* * and *Geocentrick* Places of the Planet are seen on the Ecliptick at the same Time; shewing, why the Planets seem to go backwards and forwards when view'd from the Earth; tho' they go all the while regularly from West to East, as they would be seen from the Sun.

Then to shew the true Inclination of the Orbits of the Planets; having taken off all the silver'd Balls, which represented the Planets before, and their Stems, six Orbits of gilt Brass Wire are put on, by means of Pillars set upon the Orbits, that are grav'd upon the great Plane of the Machine, so high, as to have one Part above, and the other below, the Plane of the above describ'd Ecliptick; whilst the Nodes, in two opposite Parts of the Wire Circles are exactly in the Plane of the Ecliptick, and the *Aphelia* and *Perihelia*, according to their right Distances from the Sun, directly over the Letter A and P in each Orbit grav'd below. For Example, in Jupiter's Orbit, two Pillars of equal Height (and the Height of the Plane of the Ecliptick above the great Plate) stand upon the Holes at Q and q, to sustain that Part of the Orbit, which is in the Node; a Pillar a little higher stands at N L, and one a little lower at S L. The Heads of these Pillars being round, represent Jupiter at each Node, and at the greatest North, and the greatest South Latitudes. Besides these four Balls, there are two others moveable upon the Wire, to be plac'd, one at the *Aphelion*, and other at the *Perihelion*, or in any other Places to shew the Longitude or Latitude of the Planet at any time; for one of the Steel Indices above-mentioned, laid over the

* Places seen from the Sun, or from the Earth.

Center of the Sun and the Planet, shews its *Heliocentrick* Longitude, at the same time, that a moveable Arc of a Circle vertically standing upon the Degree of the Ecliptick, which the Index is over or under, shews the *Heliocentrick* Latitude: The *Geocentrick* Longitude and Latitude, being shewn in the same Manner, when the Index comes from the Earth. *The other five Wire Orbits are fix'd the same Way.* There is also a double Index opening like a Pair of Compasses, the Hole of whose Center being plac'd on the Sun's Stem, the Legs taking in the Orbits of Mercury or Venus, the greatest Elongation of those Planets is shewn by a graduated Arc, that measures the angular Distance of those Legs.

N.B. *The greatest Elongation of ♄ is 28 Degrees, and of ♀ 48 Degrees.*

There is also a Wire bent into a parabolick Figure, and supported by three Pillars, to shew the lower Part of a Comet's Orbit; that is, so far as we can observe of it. The Head of each Pillar represents the Comet at that Place; and the fine Silver Wires extended from the Comet in its *Perihelion*, shew the Tail of the Comet, when it is biggest.

As the Observations of the Eclipses of Jupiter's Satellites, are of great Use in Astronomy; I have made a particular System of Jupiter and his Satellites applicable to the Machine, whereby the Distances, Bignesses, and true Periods of the Satellites, in respect of Jupiter, are so truly shewn; that when we have taken off the Sun and the Wheel-work, which carries ♄, ♀, ☉, and ♂, and put in the Wheel-work belonging to the System of Jupiter (each of these being put on or off in one Piece) some of the Wheels, that were before in the *Planetarium*, help to move Jupiter and his Satellites in such manner; that if they be once set by the Tables, you will by turning the Winch, see what Eclipses of any of the Satellites will appear at any Time. There is a Dial-plate with three Hands or Indices in one Corner of the *Planetarium*, where the Handle for this Purpose is applied, one of which Hands shews the Hours of Jupiter's Rotation about its Axis in our Hours, and the Hours of Jupiter (that is, dividing his Rotation once about his Axis into 24 Parts), the next Index shews the Hours of the Motions of any of the Satellites, and the third Hand shews the Days.

By looking at Jupiter and the Satellites, as they are mov'd by turning the Handle; together with the Use of the Lantern, to represent the Sun; we visibly see the following things, as they really happen.

1. The Immersion; that is, when a Satellite enters into Jupiter's Shadow.
2. The Emerfion of a Satellite out of the Shadow.
3. When the Immersions, or when the Emerfions are visible.
4. When

4. When a Satellite is hidden by the Body, of Ψ , before it comes into, or after it is gone out of the Shodaw.

5. When a Satellite may be seen to cross the Body of Ψ .

6. When a Satellite makes a Solar Eclipse in Ψ , by throwing its Shadow upon some Part of its Surface.

7. When Satellites eclipse one another.

So much is shewn by the PLANETARIUM in respect of the general System: As for the Representation of the Phenomena of the Sun, Moon, and Earth in respect to each other, there ought to be another Machine; but for Conveniency of Carriage, I have contriv'd to change the above-mentioned Machine into one for this Purpose, tho' there be some Trouble in making the necessary Changes.

Plate 32. represents the Planetarium with only the Sun, Moon, and Earth.

1. The Ecliptick being set upon other Pillars, is raised about 2 Inches higher than it was before.

2. The Surface of the Machine within those Pillars, is a Plate painted Blue, instead of the polish'd Brass before; because the blue Colour of the Sky depends wholly upon the Earth's Atmosphere, and not a blue Arch or Canopy at an immense Distance.

The Clock-work under this Plate, applying it self to the Clock-work within the Machine, and the Winch being fix'd to another Part of the Machine, the following Motions will be perform'd.

3. The Sun (which is here a gilt Brass Ball of 2 Inches Diameter) turns round his Axis from West to East once in 25 Days 6 Hours; every Turn of the Handle being now a Day, which was a Month in the general System, and the Hours being mark'd by an Index, and divided Plate, at the Bottom of the Axis of the Earth.

4. The annual Motion being stopp'd, the Earth (whose Center, as well as that of the Sun, is in the Plane of the Ecliptick) turns in 24 Solar Hours round an Axis, inclin'd to the Plane of the Ecliptick in an Angle of $66\frac{1}{2}$ Degrees from West to East, not moving out of its Place. But when the annual Motion is not lock'd, then it turns about its Axis once in 23 Hours and 56 Minutes, and at the same Time is carried round in its annual Orbit, from West to East also; its Axis is all the while remaining parallel to it self.

An Account of the Phænomena explain'd only by the Rotation of the Earth about its Axis.

Pl. 32.


The Earth is here represented by a silver'd Ball of an Inch Diameter, on which are drawn several Circles, as is done upon the common Globes, viz. the two Polar Circles, the Tropicks, the *Æquator*, the Ecliptick, and the Parallel of *London*; likewise 24 Meridians, one of which grav'd in thicker than the rest, represents the Meridian of *London*, which Place is express'd by a strong Spot or Point in $51\frac{1}{2}$ Degrees of North Latitude. From a rais'd and bent Wire-Arch *fkg* hangs a Plate like a Crescent, wanting 47 Degrees of a whole Circle at Bottom, that it may not fall foul upon the Axis of the Earth. The Plane of this Plate does always pass through the Center of the Earth, and facing the Sun, is always perpendicular to a Ray coming from the Sun to the Earth, and consequently divides the enlighten'd from the darken'd Part of the Earth: Therefore I call it a *Solar Horizon*, and W, E mark'd upon it, shew the West and the East.

In a Line suppos'd to be drawn from the Center of the Sun to the Center of the Earth, is fix'd, upon two Props, the Solar Ray or Wire *mn*, to shew on what Part of the Earth the Sun shines perpendicularly at any Day, Hour, or Minute of the whole Year.

There is an Index *b*, which shews the Place of the Earth on the Ecliptick, and likewise opposite to it, there is another Index (with a little Sun grav'd upon it) to shew the Sun's Place in the opposite Part of the Ecliptick.

Taking hold of the blue Plate, the Earth must be brought to the first Degree of Capricorn, which will be shewn by its Index; then the Axis of the Earth (always making an Angle of $66\frac{1}{2}$ Degrees, with the Plane of the Ecliptick) must be so set, that the North Pole may be towards the Sun, and a Plane passing through the Axis of the Earth, and Number XII in the Dial-Plate D, may also pass through the Sun's Center. This gives the Situation of the Earth for the longest Day in our Northern Hemisphere, the Solar Index, which points to the first Degree of Cancer, pointing also to the 10th Day of *June*: Set the Meridian of *London* to the Solar Ray, and the Index of the Dial-Plate D at Bottom to XII, and you will have the Situation of *London* at Noon for the longest Day of the Year, where an Inhabitant being suppos'd to stand with his Face towards the South, will see the Sun before him at its greatest Altitude.

Now, by gently turning the Handle of the Machine, the Point representing *London*, or an Inhabitant of it, will (by the Rotation of the Earth) be carried away towards the East, while the Sun seems to move Westward, or towards his right Hand; and when *London* is brought to the Eastern Part of the Solar Horizon, you will see the Hour of Sun-set shewn



shewn by the Index on the Earth's Axis and Dial-Plate D. Turn on, and *London* will move in the shaded Part of the Earth on the other Side of the Solar Horizon in respect of the Sun. When the Index is come to the other Number XII on the Dial-Plate, *London* is come to Midnight: As you turn on, *London* will come out from under the Solar Horizon at W, the Index below then pointing to the Hour of Sun-rise, an Inhabitant perceiving the Sun as coming out of his Horizon in the East, or on his left Hand. You may then also observe, that in the whole Turn of the Earth about its Axis, the Solar Ray only points to the Tropick of Cancer; no other Countries, but those that are under that Tropick, having the Sun vertical. And this would happen all the Year round, if the Earth had no annual Motion; neither would there be any Change of Seasons, or Alteration of the Length of Days. Here by observing, how the Solar Horizon cuts the several Parallels, one may perceive, that all of them (except the *Æquator*) between the North and South Polar Circles, are unequally divided into diurnal and nocturnal Arcs, the former being greatest on the North-side of the *Æquator*, and the latter (that is, such as are behind the Solar Horizon) on the South-side of it. The Northern Polar Circle in this Situation, is always enlighten'd, being on the Sunny-side of the Solar Horizon, whilst all the Regions contain'd within the South Polar Circle continue in the Dark, notwithstanding the Earth's diurnal Revolution; the annual only being capable to change their uncomfortable Situation of perpetual Darkness and Winter, to Day, Spring, Summer, &c.

Now, let the Movement for the annual Motion be unlock'd, and join'd to the diurnal; then by turning the Handle, the Ball representing the Earth, will move truly as the Earth does, and represent all the *Phænomena* depending upon the two Motions.

First then, whilst it continues to turn upon its own Axis, once in a Day, it goes forward in the *Magnus Orbis* from West to East, according to the Order of the Signs (or *in consequentia*, as Astronomers call it) as is shewn by its Index, pointing successively to all the Signs and Degrees of the Ecliptick, whilst the Sun, round which it moves, seems to describe the same Orbit, also *in consequentia*, or from West to East, at the Distance of six Signs; that is, seeming to set out from the first Degree of Cancer, just as the Earth does really set out from the first of Capricorn, as the Solar Index *i* (over *Taurus* in Plate 32.) plainly shews.

Now, as in the annual Motion of the Earth, the Axis always remains parallel to it self, the Situation of it, in respect of the Sun, must be continually chang'd; and a Plane passing through the said Axis and the Number XII upon the Dial-Plate D, will not pass through the Sun's Center again, when once it has left it, till at the Distance of half a Year, and then the North Pole of the Earth (which was nearest to the Sun before) will be farthest from the Sun. N. B. *The Situation of the Earth, in respect of the Pole Star, to which the Axis points, never sensibly changes;*

because the whole Diameter of the Earth's Orbit bears no Proportion to the Distance of the fix'd Stars, which are 4000000 times farther from us than the Sun, and therefore their Parallax can't amount to one Second Minute of a Degree.

Secondly, Observe, as the Earth moves on in the Ecliptick, that the North Polar Circle does gradually get under the Solar Horizon, which cuts it into two equal Parts, as the Earth enters into *Aries* (and the Sun seemingly into *Libra*) as it does also every parallel Circle all over the Earth, making the Diurnal and Nocturnal Arcs equal; and this is call'd the *Æquinox*, the Days and Nights being then of 12 Hours all over the Earth, as is also shewn by the horary Index at D: And both Poles being equally enlighten'd have the Sun in their Horizon; likewise the Solar Ray, after having successively pointed to all the Parallels suppos'd between the Tropick of Cancer and the *Æquator*, now points to the *Æquator*, whose every Part (for this Day, or one Turn of the Handle) successively receives the perpendicular Light and Heat of the Sun; which is what the *Ptolemaicks* call the Sun being in (or describing) the *Æquator*.

By continuing to turn the Handle, the Earth goes on gradually *in consequentia*, and the Solar Horizon shews how the Days continue to decrease as the diurnal Arcs shorten; and by degrees the whole Space contain'd by the North Polar Circle hides it self from the Sun, till it is quite behind the Solar Horizon, when the Earth comes to the first Degree of Cancer; which shews, that all the Countries within that Space have no Day at all, whilst the Countries within the South Polar Circle, being now on the Sunny-side of the Solar Horizon, have no Night at all. Now, the Solar Ray points to the Tropick of Capricorn, which by the Rotation of the Earth, the Sun seems (or is said) to describe. The Sun's Index also being in Capricorn, and likewise pointing to the 11th Day of *December*, shews, that we have the shortest Day at *London*, whose Measure is shewn by the horary Index. The Earth being still carried on, the Solar Ray will describe the Parallels over again, being at the *Æquator*, when the Earth enters *Libra* (and the Sun seemingly *Aries*) and the Solar Horizon divides all the Parallels into equal diurnal and nocturnal Arcs, the Poles being now in the Plane of the Solar Horizon. Then as the Earth goes on, more of the North Polar Circle comes out towards the Sun, the diurnal Arcs, and consequently the Days, in our Northern Hemisphere increase, while those beyond the *Æquator* decrease, till at last the Earth comes to the first Degree of Capricorn, from whence it mov'd at first: Then also is the Axis of the Earth return'd to the same Situation in respect to the Sun, that it had at first.

Thirdly, Take off the Earth and its Solar Horizon, and put on another Earth of 3 Inches Diameter, whose Center will still be in the Plane of the Ecliptick. This Earth having a graduated Meridian, and Pins (representing Inhabitants) perpendicularly stuck into the Tropicks, the *Æquator*,

tor, and some Parallels between; the Solar Ray (shorten'd a little to serve for this Earth) will more plainly shew, when the Sun is perpendicular to any of these Inhabitants, and how much it is more or less oblique to those without the Tropick in different Seasons, Pins being also fixed in those Places. The Solar Ray will likewise exactly shew the Degrees of the Sun's Declination upon the Degrees of the Meridian of this Globe.

Fourthly, Take off this Globe, and put on another, which is only of 2 Inches Diameter, of Ivory, having a Map of the World grav'd upon it, and a Silver Horizon to set easily for any Place. By this the Globe with its Horizon differently set, and the Solar Ray, are shewn the Phenomena of the *right* Sphere, of the *oblique* Sphere, and of the *parallel* Sphere. Having set the Horizon for any Place, you see easily by the Solar Ray, when the Sun is above or below the Horizon of that Place, and at what Point of the Compass it rises in that Place all the Year round: Whether it has Amplitude or not, whether Northern or Southern, and how much more or less in one Latitude than another. That is, how much, and when, the Sun rises and sets North or South of the Points of due East and West.

Fifthly, Put the little Earth on again, and having brought it to the first Degree of Capricorn, set the Hour-Index at XII towards the Sun, as also the Meridian of *London*. Then we are to consider, that a Plane passing along the Axis of the Earth, and the Pole Star (at an immense Distance) does also pass through the Meridian of *London*, and the Center of the Sun. Now, having unlock'd the annual Motion, that the Earth may turn about its Axis without moving out of its Place, every Turn of the Handle will carry round the Meridian of *London* till it comes back into its Place again, and *London* faces the Pole Star as before; that is, returns into the Plane, out of which it was carried; and this is call'd a *Sydereal Day*. Now, as the Sun is in the same Plane as the Axis of the Earth and the Pole Star, the *Sydereal Day* and *Solar Day* will be the same, because the Meridian of *London* comes to the Sun's Center at the same Time that it comes to the Pole Star. But as this depends upon the Immobility of the Earth's Center, it is not true in Fact; therefore to know what does really happen, we must put on the annual Motion. Then one Turn of the Handle not only gives the Earth one Turn about its Axis, but makes it go forward almost one Degree in the *Ecliptick*. Now, if you observe the Meridian of *London*, you will find, that it is not yet come to the Solar Ray; because tho' it is come into the Plane, that passes through the Center of the Earth and the Pole Star, this Plane no longer passes through the Center of the Sun, because the Center of the Earth has quitted its Place, and mov'd in *consequentia*; so that the Handle must be mov'd a little farther, before the Meridian of *London* (which has gone quite round to make a *Sydereal Day*) comes to the Solar Ray to make a *Solar Day*, which is longer than the *Sydereal Day*, the Space of four Minutes in Time. Therefore the

Earth

Earth turns about its Axis in 23 Hours and 56 Minutes; but any Country which has Noon, or the Sun in the South, must make more than one Revolution about the Axis to come to Noon again. This Difference between the Solar and Sydereal Day is so little, that it cannot easily be perceiv'd in one single Revolution; but if we turn the Handle 15 Times, those 15 Differences making an Hour (taken all together) you will see, that after just 15 Turns, the Plane passing through the Earth, and the Pole Star has so far left the Sun, that the Solar Ray instead of pointing to the Meridian of *London*, only points to that, which is immediately before it, which is an Hour more easterly, and the Hour Hand at Bottom, which points to XII, only shews the Sydereal Day, but must by means of the Handle be brought to the Number I, before it points to the Sun's Center. If the Earth be carried on one Month in its annual Course, the Hour Index, when at XII, will want 2 Sydereal Hours of the Solar Noon: After 3 Months it will want 6 Hours: After 6 Months it will want 12 Hours, or be at the Solar Midnight, when it is at the Sydereal Noon; and this is the Reason, why the Stars near the Sun (which any Day at Noon would be visible about the Sun, if it was totally eclipsed, but are invisible, because their Light is swallow'd up in his) after 6 Months will be visible at Midnight. After 9 Months, the Hour-Index will want 18 Hours of pointing to the Sun; and after 12 Months, or one annual Revolution of the Earth, it wants 24 Hours, or nothing; that is, one Day is lost, which the Sydereal Year wants of the Solar Year; this Day being made up of all the Differences, or 360 times 4 Minutes; but more exactly, $365\frac{1}{4}$ times 3 Minutes, and 56 Seconds.

Sixthly, Bring back the Earth to the first Degree of Capricorn, take off the Solar Ray, and stick a little Index into the Sun in the Hole, which it has to represent one of the Spots, which have shewn us the Time of its Revolution round its Axis. Unlock the annual Motion, and set this new Solar Index towards the Meridian of *London*, or the Hour-Index at XII. After 25 Turns and $\frac{1}{4}$, the Solar Index will be come about and point to the Center of the Earth; but the Meridian of *London* and Hour-Index, which is in the Plane of it, will be got beyond XII quite to VI. This shews the absolute Time of the Sun's Revolution round its Axis. Place every thing as before, then put on the annual Motion, observing, that as the Solar Index points to the Center of the Earth, it points also to the first Degree of Capricorn. Turn the Handle $25\frac{1}{4}$ times, and you will find, that the Solar Index does not yet point to the Center of the Earth (tho' by pointing to the 1st Degree of Capricorn, which the Earth has now left, the Sun appears to have made one entire Revolution about its Axis) but it requires near two Turns of the Handle to make it do so; this is the Reason, why Astronomers, who did not believe the Motion of the Earth, in observing a Solar Spot, reckon'd one Revolution of the Sun round its Axis, when the Sun was seen from the Earth in the same Part of the Sun's Disk, as it had been observ'd at first; and so affirm'd, that
the

the Sun was about 27 Days turning on its Axis; when that Revolution is but apparent, the real Revolution being ended two Days before.

Seventhly, Take out the Pin or Index from the Sun's Surface, bring back the Earth to the first Degree of Capricorn, and unlock the annual Motion. Then on the Moon's Orbit *a b c* fix the Moon's Stem, and put on the Moon *e d*, which is represented by a little Silver Ball, whose Diameter is $\frac{2}{7}$ ths of that of the Ball representing the Earth (according to the true Proportion of those Bodies in Nature) having one Side plain to represent that Side of the Moon, which we always see, and the other Side graved to represent that Side of the Moon, which is never seen from the Earth. Set the plain Side of the Moon towards the Earth, the Body of it being placed exactly between the Sun and the Earth, as in the time of the New-moon; then turn the Handle. You will observe; first, that 27 Turns $\frac{1}{3}$, will carry the Moon quite round its Orbit, and bring it back to the same Place again, which shews the Moon's Period is 27 Days and near 8 Hours; secondly, that the same Side of the Moon is always turn'd towards the Earth, the grav'd Hemisphere of the Ball being always averse from it; thirdly, that the Moon turns only once about its Axis, whilst it makes one Revolution round the Earth; and fourthly, that the *Periodical* Month would be the same as the *Synodical*, if the Earth had no annual Motion; that is, the Revolution of the Moon, counting its Beginning from the New or Full-moon, would not end till the next New or Full-moon: Whereas in Nature, the Period of the Moon (or periodical Month) is perform'd in something less than 27 Days and $\frac{1}{2}$; but the synodical Month, or Time from New-moon to the next New-moon, or from Full-moon to the next Full-moon, takes up the Time of about 29 $\frac{1}{2}$ Days: Because, in the Time of one Revolution, the Earth is carried on so far in the *Ecliptick*, that when the Moon has compleated its Period round the Earth, it is no longer in a Line with the Center of the Earth and Sun; but must move on still in its Orbit above 2 Days to be in a Line with the Sun and Earth. This is easily shewn by the Machine. For if you bring the Earth to the first Degree of Capricorn, and place the Moon between the Earth and Sun; and, having put on the annual Motion, turn the Handle 27 $\frac{1}{3}$ times, the Earth will be got on so far in the *Ecliptick*, that the Moon (tho' it has compleated its Period) is not yet in a Line with the Sun; but if you hold a String over the Center of the Moon and Earth, and another over the Center of the Sun and the first Degree of Capricorn (which Line the Earth and Moon did set out from) you will find those two Strings to be parallel, which shews that the Moon has really finish'd its Period; tho' it does not seem to have done so in respect of the Sun. But if you give the Handle a little more than two Turns, you will finish the synodical Month, by bringing the Moon again between the Sun and Earth: And the Earth's Place on the *Ecliptick*, will shew that it has advanc'd 29 Days and an half.

Pl. 32.

Eighthly,

Eighthly, Bring back the Earth to the first Degree of Cancer, and having put on the black Cap, which belongs to the Moon, with its convex Side opposite to the Sun as at *e*, bring the Moon between the Earth and Sun; that is, to the Place of New-moon: Then turning the Handle, all the Phases of the Moon will be given successively, as they happen, whilst the Moon going round the Earth in its Orbit, is also carried round the Sun, together with the Earth. For, first, the black Cap representing the shadowed Part of the Earth, does wholly cover all the white Hemisphere of the Moon, which is turn'd towards the Earth, thereby shewing, how the New-moon comes to be invisible to us; then after a few Turns of the Handle, you may perceive, that a small Part of the white Hemisphere is come from under the Cap (that is, out of the Shadow) and makes a Crescent, and as you go on, you see the first Quarter; the Half-moon after 7 Turns, and still more and more comes from under the Cap, till 14 Turns and $\frac{1}{2}$ makes the *Phænomenon* of Full-moon, when the white Hemisphere is come quite out of the Cap, and the Earth is between the Sun and Moon. As you turn on, the Phases of the Moon gradually decrease till you come to have New-moon again. And now all the above-mention'd *Phænomena*, may be seen at the same Time, as it happens in the Heavens.

Ninthly, Take off the Sun, and in the Place of it, put the Lantern belonging to the *Planetarium*, which has two convex Glasses (without which we could not have the Rays parallel, converging, or diverging, with sufficient Splendor) to shew how the Earth and Moon are enlighten'd by the Sun, together with the *Phænomena* of Eclipses. Let the Room be made dark, and then you will see the *Phænomena* of Day and Night, and the Seasons and Phases of the Moon above-mention'd, very plainly, without either the solar Horizon, or Moon's black Cap, which were only made use of, when we could not compare Light and Darkness.

By making Use of the bigger Earth, with Pins properly fix'd for Inhabitants, you may plainly see what Geographers say concerning the Inhabitants of the Earth, in respect of their Shadows, as they call them *Ascii*, *Amphiscii*, and *Heteroschii*; you will also the better compare together the Days and Seasons of the *Antæci*, *Periæci*, and *Antipodes*; as likewise (when you use the Ivory Earth, with its Silver Horizon) see plainly how the Sun appears to rise above our Horizon and set below it, with or without Amplitude, real Rays of Light now performing what the Solar Ray of Brass Wire did before but imitate. There is also a Contrivance of a Silver Plate, to put upon any Place of the Surface of the Earth, parallel to the moveable Horizon above-mention'd, and which turns along with it, to shew the Distinction between the sensible and rational Horizon of a Place; but at the same time to make it appear, that the Sun's immense Distance makes them so nearly co-incide, that the rising and setting is the same at the same moment in respect to both.

The

The Moon's secondary Light (that is, the Light, which being reflected from the Earth, gives the dark Part of the Moon's Hemisphere, which is turn'd towards the Earth, such a faint Illumination as we receive from the Moon) whereby in serene Weather, at the time of the Moon's Crescent or Wane, we see the remaining Part of the Moon's Disk of a dun Colour (nay and sometimes see the whole Moon it self at the very Time of the New-moon) is very well shewn here by the Light of the Lantern; as also how it is, that the Earth in its Turn, becomes a Moon to the Moon.

In order to shew the *Phænomena* of Eclipses; first, put on the little Earth, the Moon being on its Stem, and the Lantern in its Place, and you will observe, how the Moon in passing between the Sun and Earth, throws its Shadow upon some Part of the Earth, whose Inhabitants have a total Eclipse of the Sun in those Places, which the Shadow passes over, and a partial one in the *Penumbra*, or near the Shadow. When the Moon being Full, passes on the other Side of the Earth, so as to lose her Light by the Interposition of the Earth's Body, that is, as she goes through the Shadow of the Earth, there is a total Eclipse of the Moon, which all the Time of its Duration is seen by all the Inhabitants of that Hemisphere of the Earth, which is towards the Moon at that Time. Now, tho' this Position of the *Planetarium*, does shew what a Solar and a Lunar Eclipse is; it does not represent what really happens in regard to the Time of Eclipses; for it shews an Eclipse every New and Full-moon, whereas we seldom have more than four Eclipses in a whole Year, and sometimes not so many. The Reason is, that tho' the artificial Earth and Moon on the Machine, do bear a just Proportion to one another, the Orbit of the Moon is 10 times less than it ought to be in Proportion to the Bodies: Besides, the Orbit of the Moon is not in the same Plane with the Ecliptick, but makes an Angle of about 5 Degrees with it; so that unless the Moon be in, or near, the Nodes (or the Places where the Moon's Orbit cuts the Plane of the Ecliptick) at the Time of Full or New-moon, there will be no Eclipse; the Shadow of the Moon passing northerly over, or southerly under the Globe of the Earth at New-moon; and the Moon, at Full, passing northerly over, or southerly under the Shadow of the Earth at Full-moon. Some of the *Orrery*-makers indeed have made the Moon rise and fall in its Course, in an Orbit properly inclin'd, and besides, have made the Nodes of that Orbit move backwards, or *in antecedentia* (that is, from East to West) one Revolution in 19 Years, as it happens in Nature, but neither increasing the Moon's Orbit, nor putting on smaller Balls for the Moon and Earth, this Inconveniency of too frequent Eclipses has not been avoided. Others, to avoid it, have given the Moon's Orbit 3 or 4 times a greater Inclination than it really has, destroying one Truth to explain another.

Now, in my *Planetarium*, I shew the true Inclination of the Moon's Orbit, the Motion of the Nodes, and true Magnitude of the Orbit, in

M m m

respect

respect of the Bodies, and consequently the true *Phænomena*, relating to the Time and Quantity of the Eclipses, by making the following Changes.

I take off the Moon and Earth, and upon the silver'd Circle *np* (which is divided in 19 Parts, corresponding with 19 Years to shew the Motion of the Nodes) I fix upon Pillars an Orbit of fine gilt Wire, properly inclin'd, with Nodes upon it, and two small Moons, which, as well as a small Earth, now to be put on, bear a just Proportion to the Bigness of the Orbit: And now by means of a little Piece of white Paper, held behind the Earth at the Time of New-moon, the Shadows of the Earth and Moon, shew whether there will be an Eclipse of the Sun or not; if an Eclipse, whether partial or total, of long or short Duration, whether Northerly or Southerly; likewise by holding a Paper behind the Moon at the Time of the Full-moon, you may see whether there will be an Eclipse of the Moon or not; and if an Eclipse, of what kind, &c

Tenthly, There are Parts to fix to the *Planetarium*, to shew what *Parallaxes* are, and how they are to be observ'd; as the annual, menstrual, and diurnal; making it very evident, how the Distances of the Sun, and that of the fix'd Stars, have been attempted, but not sufficiently settled by those Methods, but that of the Moon exactly discover'd. There is also a Contrivance to shew what is meant by the *Longitude* at Sea, which shews how it must be at last found by the Appulses of the Moon to the fix'd Stars, when the Theory of the Moon comes to be perfectly settled.

I have also contriv'd an additional Part to shew a *Celidography* upon the *Planetarium*; that is, the *Phænomena* of the Spots of Venus, of Day and Night, and the Seasons in that Planet, which (according to *Signior Bianchini's Discoveries*) does in 24 Days and 8 Hours, turn round an Axis inclin'd in an Angle of 15 Degrees, to the Plane of the Orbit of the Planet, that Axis always continuing parallel to it self. There is also a Wire, to represent a perpendicular Solar Ray, and a Solar Horizon, to shew how sensible a Spiral the Sun must daily seem to move in to an Inhabitant of Venus. The Lantern also may be put on instead of the Sun, &c. But I forbear fixing this additional Machine to my *Planetarium*, till other Astronomers Observations have confirm'd *Signior Bianchini's* new and great Discoveries.

The Machine to shew mechanically, how Planets and Comets, by a Ray drawn from the Sun, describe Areas proportionable to the Times, which I promis'd in the sixth Annotation to Lect. 5. (Page 406.) properly belongs to the Planetarium; therefore I choose to describe it here.

See Plate 29. Fig. 7, and 8.

The 7th Figure represents the upper Surface of a Wooden-Frame, or Case of Clock-Work, held together by four Screws. *ETFV* is a silver'd circular Plate, divided into 88 Parts (88 being the Number of Days of Mercury's periodical Time) with an Handle or Winch *HG*, and Index *GE*. *IKPLM* is a silver'd oval Plate, divided into the same Number of Parts as the circular Plate, with a Channel or Groove in it, for the little Brass Ball or Planet *P* to move in. *S* represents the Sun in one of its *Foci*, *L* the *Perihelion*, and *I* the *Aphelion*. *SO* is an Index of Steel, which going through the Planet *P*, carries it in the elliptick Channel, with a Velocity reciprocally proportional to its Distance from *S*. The Part *SP* of this Index, which is intercepted between the Sun and Planet, or *S* and *P*, represents the *Radius Vector* (already describ'd) which continually becomes shorter, as the Planet goes from the *Aphelion* to the *Perihelion*, and lengthens gradually as the Planet goes from the *Perihelion* to the *Aphelion*: The Part *PO* of the Index, which goes beyond the Planet, and whose End *O* does here describe the Circle *OQRIN*, is of no Consequence in our Astronomical Consideration. If a Planet (as for Example, Mercury) moved in a circular Orbit, concentrick with the Sun, it would not only describe equal *Areas* in equal Times, but also equal *Arcs*; then by uniformly turning the Handle *H*, the End of the Index at *E* would truly represent the Motion of such a Planet. But if the Planet moves in an Ellipse in one of whose *Foci* the Sun is, and the *Perihelion* be 6 times nearer than the *Aphelion*; the Planet setting out from the *Perihelion* *I*, will move in an accelerated Motion in the Direction *IK*, *PL*, and from *L* will come by *M* to *I* with a retarded Motion, and so in all its Revolutions continually change its Velocity, describing equal *Areas* in equal Times, whose Equality is shewn, and prov'd in the following Manner. The Planet *P* with its Index through it, being brought to *I*, set the Index *GE* at *E*; then by means of the Handle *H*, carry on the Index 10 Degrees, or from *E* to *T*, and you will perceive at the same time, that the Planet in the Ellipse will only go from *I* to *K* $3\frac{1}{4}$ Divisions, the *Radius Vector* having moved from *SI* to *SK*, whereby the *Area SIK* will be describ'd. Then if the Planet be set at the *Perihelion* *L* (to which Place it will be brought by turning the Handle *H* till the Index comes to *F*) the *Radius Vector* on the Ellipse will then be *SL*: And if the Index of the circular Plate, which represents the Time of the Motion, be carried on from *F* (44 Degrees) to *V* (54 Degrees) the Planet will go from *L* the whole Arc *LM*, or 20 Degrees, setting out 6 times faster than it did at first from *I*, the *Radius Vector* describing the *Area SLM*, so much broader than *SIK*, as *SL* is shorter than *SI*, which makes this last *Area SLM* equal to the former *Area SIK*, and the Velocity of the Planet reciprocally as the Distance. If any intermediate Place be taken in the Curve of the Ellipse; as

M m m 2

for

for Example, the Point *p*, whilst the Index on the circular Plate moves 10 Degrees, the Planet will go through the elliptical Arc *pP*, the *Radius Vector* describing the *Area pSP* equal to the former *Areas*.

Now, to shew the Contrivance of the Machine, let us take off the upper Board of the Frame, as in *Figure 8*, where the circular and oval Plates, the Planet-Handle and the Indices, are only represented in prick'd Lines to shew their Places. Under the Bar or Bridge *VG*, are two Ovals *QT* and *ON*, moving each other about by means of a Cat-gut in a Groove on their Edge, which Gut crosses at *K*. These Ovals have their Centers of Motion on their alternate *Foci* *S* and *I*. The Sun and Center of Motion of the *Radius Vector* is fix'd over *S*. Now if the Handle be placed at *I* the Center of the other Ellipse *NO*, that Part of the Circumference of the Ellipse *QT*, which touches the Circumference of the Ellipse *NO* is carried round with Velocities proportionable to the Distances from *I*, or to the Length of the Lines *IK*, *I₄*, *I₃*, *I₂*, *I₁*, &c. which represent unequal Leavers. Now, in the present Situation represented by the Draught, the Velocity of the Planet *P* in *Perihelio* is greatest of all; for *PS* being equal to *SK*, *P* has the Velocity of the Point *K*, which is led by the longest Leaver *IK*. But when the Ellipse *NO* has made half a Turn, the Ellipse *QT* changes its Situation, its Point *V* coming to *s*, where it is led by the Leaver *I₁*, six times shorter than before, and the Planet having slipp'd along its *Radius Vector*, quite to its End, is over the Point *I* at its *Aphelion*, where it has 3 times less Velocity. The Rest in Relation to the different Situation of the Ovals may be easily conceiv'd by carefully observing the Figure. As for the two Wheels, whose Numbers of Teeth are equal (no matter what the Numbers are) they are only made Use of to keep the Circle and Oval on the Surface of the Machine (*Fig. 7.*) at a proper Distance, and to give the Planet and the Index of the circular Plate of the same Direction.

N. B. *The Ovals of Wood, and the oval-silver'd Plate on the Machine, must be similar; but they may be any kind of Ellipses. I have taken one here, more excentrick than any Planet's Orbit, only to make the Phænomena the more sensible; tho' not so excentrick as a Comet's Orbit, which would have exceeded the Bounds of the Machine.*

The END of the DESCRIPTION of the PLANETARIUM.

ADDENDA.



A D D E N D A.

Betwixt No. 28, and No. 29 in the First Lecture, add what follows.

IF two leaden Balls of about an Inch Diameter be pared with a Knife, so as to cut off a Segment of about $\frac{1}{4}$ of an Inch Diameter, and they be press'd together (giving them a little Twist) pretty hard, they will stick together with a great Force; sometimes so great as to hold above 100 lb. See the 6th Volume of the *Philosophical Transactions*, abridg'd by Messrs. Reid and Gray, Part 2d Page 2 and 3.

Betwixt the 20th and 21st Annotation to Lect. 1. this is to be added.

Dr. Petrus van Muschenbroek, the ingenious Professor of *Astronomy*, &c. at *Utrecht*, has with indefatigable Pains and Application, made Experiments of the Attractions and Repulsions of Load-stones, in respect to Iron, and to each other; but could never find any regular Proportion in the Increase of Attraction in their Approach to, or Decrease of Attraction in their Recess from, one another — only, that the Force of the magnetick Virtue did increase in the Approach to, and diminish in the Recess from, the Stone; but not exactly as the Distance, nor as the Square, or Cube of the Distance, nor as the Square, or Cube of the Distance reciprocally — nor in any Proportion reducible to Numbers; and therefore he very reasonably conjectures, that the Repulsions and Attractions disturb one another, so as to confound the Proportion; nor are we to hope for any Rule concerning this Matter, till a Way be found (if ever it can be) of separating the attracting from the repelling Parts. I refer the curious Reader to his Dissertation concerning the Magnet, which is very worth every inquisitive Philosopher's Perusal. It is printed at *Leyden* in *Quarto*, with several other valuable Dissertations of the same Author.

After the 23 Annotation to Lect. 1. what follows concerning Electricity, may be added.

Mr. *Stephen Gray*, who has made greater Variety of electrical Experiments, than all the Philosophers of this and the last Age; has, since I began this Book, found out several new *Phænomena* in Electricity, the Particulars of which may be seen in the *Philosophical Transactions*, No. 417.

But for the Sake of such, as may not have an Opportunity of seeing the *Transactions*, I will give a general Account of his Discoveries.

1. That all such Substances as cannot be made electrical by rubbing, may yet receive an electrical Virtue (as is visible by their attracting and repelling Leaf-Gold) from a rubb'd Glas Cane (whether solid or hollow) not only in contact with the Tube, but by the intermediation of a String to very great Distances, as for Example, of above 800 Feet. Nay, if the Tube does not touch, but is only rubb'd near the String, the electrical Virtue will run along. And what is very remarkable, as the Virtue runs along an hempen String, that String must be supported by Hair, or Silk Lines; cross Strings of Hemp or Flax, or even Wire, stopping the Propagation of the Virtue to a Distance, by receiving it laterally.

2. Very large Surfaces, as Maps, Table-Cloaths, &c. are impregnated with electrical *Effluvia*.

3. A Load-stone, and Iron hanging at it, will receive the electrical Virtue, which therefore is not disturb'd by magnetick *Effluvia*.

4. The Electricity will be carried several Ways at the same Time, without touching the String of Communication with the Tube; as to two Balls of Ivory at the Ends of a very long String by rubbing the Tube near the Middle of the Spring.

5. The electrical Attraction receiv'd, is not proportionable to the Quantity of Matter in Bodies, a solid and an hollow Cube of Wood attracting the one as much as the other.

6. The *Effluvia* are also carried round in a Circle, and communicated from one Circle to another.

7. This Virtue will be communicated to Leaves of Trees; nay even to Fluids, as to Bubbles of soap'd Water.

8. Animals also will receive this Virtue. A Man suspended horizontally by two Hair Lines, attracts and repels Leaf Gold with his Face and Hands, if the Tube be rubb'd near his Feet; nay, at the End of a long Fishing Rod held in his Hands, a Ball will attract and repel. It is remarkable, that the Virtue is strongest at that Part of the impregnated Body,

Body, which is farthest from the Tube ; for if the rubb'd Tube be held over the Head, not the Head, but the Face will then attract, and so *vice versa*. If the Man but just touch the Floor with a small Cane or Wire, the Virtue runs to the Floor, and the Man is no longer electrical. So hanging by an hempen Rope, carries away the Virtue to the Cieling.

9. If two Boys be by Hair-Lines suspended as above, and at a considerable Distance from one another, the Virtue may be communicated from the one to the other by a String, which both hold, or tied to their Cloaths, the Electricity being always strongest in him, who is at the greatest Distance from the Tube.

10. Mr. *Stephen Gray* has found since, that if a Man stands upon a Cake of Rosin, or of Glass, or of some other Substances, which are of themselves electrical, or become so by rubbing ; the Effect will be the same as if he hung from a Hair-Line. So that if a whole Regiment of Soldiers were in a Line, and each Man standing upon a Cake of Rosin held the next by the Hand, or only communicated with him by a String, I do not doubt, but they might all be so impregnated with electrical Virtue ; that when the Tube should be rubb'd near the first Man, the last would with his Hand, Face, or Cloaths, attract and repel Leaf-Gold.

11. It has been also found, that Electricity will be communicated thro' dense and large Bodies.

After the End of No. 8. Lect. 2. add—

Care must be taken not to draw Consequences from this Experiment ; because the Success of it will be different according to Circumstances. If the cylindrick Weights * be plac'd quite close to the Spring when it is bent, the Weight K (of 8 Ounces) will be shot farther than half the Distance that L (of 4 Ounces) is shot to ; and that will happen for the Reasons given in the 4th Annotation of *Lect. 5. Page 397 and 398.* But if both the Weights be set at some Distance from the bent Spring, so that the Spring may give a sudden Blow, and not act long upon the Cylinders ; then the Experiment will succeed. *N. B.* That Distance must be found by Tryals.

At the End of the 12th Annotation to the 2d Lecture.

Mr. *Charles de Labelye*, having further consider'd the Motion of the loaded Cylinder (see *Plate 4. Fig. 15.*) as it rolls up on an inclin'd Plane, communicated to me several Propositions relating to that Motion, which I thought would not be unacceptable to the curious Reader.

PROP.

PROP. I. Plate 6. Fig. 3.

To find TS universally.

Let $MT = a$, $MS = b$, $MB = x$. Then we shall have $\sqrt{aa - xx} = BT$, and $\sqrt{bb - xx} = BS$. Now, since $BT - BS = TS$, it will follow, that $\sqrt{aa - xx} - \sqrt{bb - xx} = TS$, which consequently will be known by making $x =$ to the Sine of the Angle of Inclination of the Plane with the Horizon, supposing MT Radius.

Another Way, without extracting Roots.

As $MS (= b)$: is to the Sine of the Angle $MTS ::$ so is $MT (= a)$: to the Sine of the Angle $MS T$; which being known MS is also known: then say,

As the Sine of the Angle MTS : is to $MS (= b) ::$ so is the Sine of the Angle $MS T$: to TS required.

Now, supposing the Center of Gravity in C , it is evident, that the Cylinder will roll up till the Point V in the Circumference (having made $CV = CR = TS$) comes to v ; for then the Center of Gravity will be in the same Position, as when at S over T . And to find in what Point of the Line Ta the Point V will be applied, it is evident, that the Line Tv will be equal to the Arc TV : therefore supposing a Cycloid * whose generating Circle is equal to the Base of the Cylinder and Vertex at T and Axis upon TA , if you draw VO parallel to the Plane TF ; by a Property of the Cycloid, the Line VO , being part of an Ordinate, will be equal to the Arc TV . Therefore making Tv equal to VO , you will have v the Point required.

Pl. 6. F. 4.

Another Way.

Since the Ordinate HO is made up of VO equal to the Arc sought, and HV equal to the Sine of the Angle of the said Arc; by placing the same Cycloid, so as to have its Vertex at H , and Axis on HT , it will cut the Plane in Y , so that HO will be equal to TY , from which subtracting $Yv = HV$, the Remainder will be $Tv = VO =$ the Arc TV requir'd.

Another Way.

Draw through V , VH parallel to the Plane; and placing the Cycloid with its Vertex at W , and its Axis on WL parallel to HT , it will cut the Plane at v ; for $Lv = TY = HO$, and LT being equal to $WH = HV = vY$, there will remain $Tv = VO =$ the Arc TV required.

PROP.

P R O P. 2. Plate 6. Fig. 5.

It may be farther ask'd, *What Degree of Inclination to the Horizon, a Plane must have for the Cylinder to roll up that Plane the Length of any given Arc of the Circumference of its Base*; as for Example, suppose the Arc given to be $T V$.

To find this, we must consider, that when the Center of Gravity is over V , after the Cylinder has roll'd, the Line CV , drawn through that Center of Gravity V , will make the same Angle with the Line VM (drawn through the Point V and M , the Center of Magnitude) as the Line CT makes with TM , when the Center of Gravity is at S over T . Now, since the Angle CVI or CVM is equal to CTM , and $CIV =$ to its opposite MIT ; it must follow, that TMV given, is equal to TCV sought. Whence we conclude, that a Circle passing through T , M and V , will also pass through C . From whence we obtain this following Method—*The Arc TV being given, to find the Center of a Circle passing through T , M , V ; which Circle will cut a Circle describ'd round the Center of Magnitude M (with the Distance of the Center of Gravity from it) at C : then making $CR = CV$, draw RCT , and where it cuts the Circle of the Basis of the Cylinder at T , draw MT , making the Angle MTS ; and this Angle will be equal to νTZ , the required Angle of Inclination of the Plane $T\nu$, as is evident from what has been said.*

P R O P. 3. Plate 6. Fig. 5.

It may be also demanded, *How many Degrees of the Circumference, or what Part of an Arc, will the Cylinder roll upon, to describe a given Space on the inclin'd Plane*; that is, $T\nu$ being given, to find TV . This may be done two Ways; for having the Diameter, you must say—As 113 : is to 355 : so is the Diameter: to the Circumference, half of which will be the longest $T\nu$, which the Cylinder can describe. So by the Rule of Proportion—as half the Circumference: is to 180 Degrees:: so $T\nu$ (measur'd in Parts of the same Bigness, as those of the half Circumference): will be to the Number of Degrees in the Arc TV .

But if it be required to find the Point V by Construction; upon the Point T of the Line $T\nu^*$ or the Plane given, erect perpendicularly the ^{*Pl. 6. F. 6.} Line TA or Diameter of the Cylinder, and describe the Circle equal to the Basis of the Cylinder, then placing the Cycloid made use of before, so that the Axis being kept parallel to the Diameter of the Circle (which is here equal to the generating Circle) when the Vertex of the Cycloid is in the Circumference of the Circle, and the cycloidal Curve falls at the same Time on the Point ν ; you will have given you, by the Vertex of the Cycloid,

cloid, the Point W; then taking T V equal to T W, it will be also equal to T u from what has been said before, &c.

As for finding the true Maximum of the Rise of the Cylinder, it cannot be obtain'd without finding some algebraick Formula, or Expression, for the Rise of the Center of Magnitude, as well as another Formula, without any more unknown Quantities, than in the former Way of expressing the Descent of the Center of Gravity of the Cylinder whilst it is rolling. Now from a due Consideration of the Figure, it appears, that these Formulæ require the Rectification of the Circumference of a Circle, or of some Arc thereof; which, not being possible to be done without infinite Series's, has made me leave the Subject.

After No. 45. (Lect. 3.) Page 104, add—

There is another very curious Contrivance, whereby a Power, whose Intensity is continually diminishing, does yet produce an Effect continually increasing, which is a Sort of a mechanical Paradox; but as it is in common Use, we are apt to overlook it. I mean the Application of the main Spring in a Gun-Lock, to drive the Cock, which carries the Flint against the Hammer (or Steel-Plate, to strike Fire) with an accelerated Motion, tho' the Spring is all the while unbending it self. FSP* is the Spring bent up to the highest Degree, when the Gun is cock'd; but its natural Situation when wholly unbent, is FSp. ACT is the Tumbler, whose Axis of Motion is at C, which is the middle of an Axel or Arbor, going through the Plate of the Lock, on the End of which the Cock (being the Weight to be mov'd by the Power) is fix'd upon a Square. Now, the Tumbler is in Effect an *Axis in Peritrochio*, with several Wheels, where the Power is successively applied from the least to the greatest, as in the Fuzee of a Watch. I have represented only three of them, according to the three most equal Positions of the Spring, by the pointed Circles $\alpha \beta$, $a b$, and A B. Now, tho' the greatest part of these Wheels is cut away when the Tumbler is reduc'd to its proper Figure A $\alpha \beta$ T t, there is enough left of each of them for the Hook of the Spring P to apply it self successively to them all, as it presses on the Arm A α , with a sliding Motion from α to A, which is the whole Range of the Spring in its Action. When the Gun is cock'd, as in this Figure, the End of the Spring being applied at α , acts upon the Tumbler at the Distance α C, by the Wheel $\alpha \beta$, so as to press upon it at the greatest Disadvantage, at the same time, that the Spring is most bent, or has the greatest Intensity. This is felt by applying the Hand to the Cock, which then makes the least Resistance.

Pl. 22. F. 8. In the 8th Figure, the Gun being at Half-Cock, the Spring acts upon the Tumbler with more Advantage, pressing in the Tangent of the Circle, or Wheel

Wheel *a b*, at the Distance *a C*, which gives the Cock more Force to descend, as may be felt by the Hand: For tho' the Spring is a little unbent, and so somewhat weaker; the Diminution of Intensity in the Spring, is not so great as the mechanical Advantage gain'd by the Distance being increased from αC to *a C*.

In the 9th Figure, the Spring is still more unbent; but yet the Cock is Pl. 22. F. 9. press'd down with a much greater Force, because the End of the Spring *P* now acting in the Tangent of the Circle *AB*, has *AC* for its Distance of Power, about 3 times greater than αC (the Distance of Power when the Gun is cock'd) whereas it is not unbent above $\frac{1}{3}$, as may be seen by comparing the 7th Figure with the 9th.

At the End of the 8th Annotation to Lecture 3. Page 167, Insert what follows—

The late Mr. *Peter Daudé*, from whom I had this Proposition concerning the Pressure upon the Center-Pin of an *Axis in Peritrochio*, not having at first consider'd, that the rising and the falling Weight in the turning the *Axis in Peritrochio*, partake of the Nature of a Pendulum (because, as they have different Velocities, they mutually accelerate and retard one another) gave me afterwards the following Solution, wherein that Error is corrected. But, as it depends upon the Doctrine of Pendulums, which I have only consider'd towards the End of this first Volume, I thought proper not to give it the Reader till after I had treated of Pendulums. But, before I enter upon Mr. *Daudé's* Solution, it is proper to premise something concerning the Center of Oscillation.

1st, The Center of Gravity of two Bodies join'd together by a mathematical Line, is that Point in the Line, where each of those two Bodies acts with its whole Weight in a State of Rest; and this happens when their Momenta, which are in that Case the Product of their Quantities of Matter, by their Leavers, or Distances from that Center of Gravity, are equal. So that that Center feels the Weight of both Bodies.

2dly, The Center of Oscillation of two such Bodies, is a Point in that Line, upon which the two Bodies in Motion act with all their Momenta; and this happens, where the Products of their Momenta, by their Distance from that Center of Oscillation are equal; in which Point their Forces being equal, and their Motions not being contrary to one another, they act, or fall, so that this Point moves with the natural Velocity of falling Bodies, and they strike an Obstacle with their whole Weight.

3dly, If we suppose these two Bodies revolving round the Center of Suspension, the Centers of Gravity and Oscillation, will both be on that Side, where the greatest Momentum is.

4thly, If the Center of Suspension be between the two Bodies, the Center of Oscillation will be on the Outside of the two Bodies. But if the Center of Suspension is on the Outside of the two Bodies, the Center of Oscillation will then be between the two Bodies.

Now, the Proposition, according to Mr. *Daudé's* last Solution, will be thus express'd.

PROPOSITION.

To find the Pressure, which two Bodies p and q in their Motion, with contrary Directions, exert upon the Center-Pin of an Axis in Peritrochio or double Pulley made of one Piece.

Let us suppose p greater than q , then the Center of Gravity of the two Bodies p, q being beyond their Center of Motion c towards the Body q , the Body q will descend and cause the Body p to rise.

Therefore to find their Center of Gravity s , we must make Use of this Analogy.

$$q : p :: b + a - z : z, \text{ and } qz = pb + pa - pz, \text{ which gives } z = \frac{pb + pa}{q - p}; \text{ and taking that Value from } a, \text{ we have } a - \frac{pb + pa}{q - p} = \frac{qa - bp}{q + p} = cs.$$

Now, considering the Motion of the two Bodies p, q , as that of a double Pendulum, whose Weight p, q , hang one below the Center of Suspension, and the other above it, we must find their Center of Oscillation o thus,

$$qa : pb :: b + a + x : x (qo) \text{ and } qa - pb : pb :: b + a : x = qo = \frac{pb + pba}{qa - pb}, \text{ and } qo + qc = co = a + \frac{pb + pba}{qa - pb} = \frac{qaa + pbb}{qa - pb} = co.$$

Therefore since the Center of Oscillation falls with the natural Velocity of heavy Bodies, which we will call $V=1$, let us make this Analogy $co \left(\frac{qaa + pbb}{qa - pb} \right) : cs \left(\frac{qa - pb}{q + p} \right) :: V(1) : sg$ the Velocity of the Fall of the Center of Gravity s . Then reducing the two first Terms of this Analogy to the same Denominator, we shall have this, $co(qqaa + qpbb + pqa + ppb) : cs(qqaa - 2qba + pbb) :: 1 :$

$\frac{qqaa - 2qpba + pbb}{qqaa + qpbb + pqa + ppb} = sg$, the Velocity of the Fall of the Center of Gravity. Therefore multiplying that Velocity of their Center of Gravity by the Sum of the two Bodies $p + q$, we shall have the *Momentum* of their Fall (or so much of their whole Weight as actually falls) equal

equal to $\frac{qqaa - 2qpba + ppbb}{qaa + pbb}$, which being subtracted from their whole natural *Momentum*, namely from $qV + pV$, there remains $\frac{qpbb + 2qpba + qpaa - qp \times b + a^2}{qaa + pbb} = \frac{qp \times b + a^2}{qaa + pbb}$, which is so much of their Weight as presses upon c their Center (or rather Axis) of Motion.

From the last *Æquation* but one, may be drawn the following *Corollaries*.

C O R O L L A R Y 1.

If $b = a$, the Pressure on c the Axis is $\frac{4pq}{p+q}$ as in the single Pulley.

C O R O L L A R Y 2.

If q be infinite, the Pressure upon the Axis will be $\frac{p \times a + b^2}{a}$, whence if $a = b$ the Pressure will be $4p$ as in the single Pulley.

C O R O L L A R Y 3.

If b be infinite, the Pressure upon the Axis will be q , because then the Leaver a is nothing in respect of the Infinite b .

C O R O L L A R Y 4.

If a be infinite, the Pressure upon the Axis will be p , because then the Leaver b is nothing in respect to the Infinite a .

S C H O L I U M.

It has been found above, that the Pressure upon the Center of the Axis in *Peritrochio*, is denoted universally by this Expression, $\frac{qpbb + 2qpba + qpaa}{qaa + pbb}$; but when $qa = pb$ there is an *Æquilibrium*, and then the Axis supports the two Weights $p + q$, and in that Case $\frac{qpbb + 2qpba + qpaa}{qaa}$ must $= q + p$, which (supposing it so) gives $qpbb + 2qpba + qpaa = qqa + qpbb + pqa + ppbb$; therefore by Reduction, $2qpba = qqa + ppbb$, and $qqa - 2qba + ppbb = 0$; and by extracting the Square Root $qa - pb = 0$, and consequently $qa = pb$.
Therefore

Therefore it is true, that when $qa = pb$, one may equally deduce from it the Truth of our general Expression, namely, that the Pressure upon the Axis is $q + p$. Q. E. D.

The Weight, which presses upon c , the Axis of the double Pulley cannot be increas'd by the Acceleration of the Weight q , because what makes the Weight of Bodies, which do not fall, by reason of an Obstacle hindering them, is only every single Impulsion of Gravitation, and not the Sum of many of those Impulsions; because each particular Impulsion is destroy'd by the Obstacle as soon as it is given. But the Impulsions upon the descending Weight do, each of them, give it a Velocity, which is not destroy'd; therefore the given Velocities, being accumulated, cause an Acceleration of Descent in the Weight q , whilst the Weight, that presses on the Axis of the Machine or Pulley, is always equal and the same.

Page 189. After the Words — *All other Cases may be deduced from this*, insert —

Tho' this may serve for finding the Sum of the Frictions to direct our Practice; yet as I have in the Calculation omitted several Decimals, whereby the Solutions might have come nearer to the Truth; for the sake of the Curious, who may expect a Solution entirely exact, I give here a Method to find it.

Since the Weights, that are to be added to the Power, in order to overcome Friction, continually decrease, and always in the same Ratio, they may be consider'd as the Terms of a geometrical Progression, whose last Term is ϕ .

In order to find the Sum of the Terms of such a Progression, the two following lemmatory Propositions must be premis'd.

I.

In a geometrical Proportion or Progression, the Sum of the Antecedents: is to the Sum of the Consequents:: as any one of the Antecedents: is to its Consequent. (See Euclid.)

II.

In a geometrical Progression, the second Term Minus the First: is to the First:: as the Last Minus the First: is to the Sum of all the Terms preceding the Last.

Let the Progression be $\div \div a . b . c . d . f . g .$ we must prove, that $b - a : a :: g - a : a + b + c + d + f$.

The

The Progreſſion may be thus always expreſs'd, $a:b::b:c::c:d::d:f::f:g$; but we have by the laſt $a:b::a+b+c+d+f:b+c+d+f+g$.

And by Inverſion,

$$b:a::b+c+d+f+g:a+b+c+d+f;$$

By Diviſion,

$$b-a:a::b+c+d+f+g-a-b-c-d-f:a+b+c+d+f;$$

And taking away the Expreſſions with contrary Signs, it will be

$$b-a:a::g-a:a+b+c+d+f: \quad \text{Q. E. D.}$$

C O R O L L A R Y 1.

So that calling the Exponent of the Progreſſion univerſally n , the firſt Term a , the laſt Term z , and S the Sum of all the Terms, excepting z , we have by the ſecond Lemma $na-a:a::z-a:S$, which gives $\frac{za-a^2}{na-a}=S$, which being reduced gives $\frac{z-a}{n-1}=S$. That is in Words at length.

The laſt Term Minus the Firſt, divided by the Exponent of the Progreſſion, leſſened by the Number 1, is equal to the Sum of all the Terms but the laſt.

C O R O L L A R Y 2.

And in Progreſſions, whoſe Terms continually decreate (as thoſe of the Weights to overcome Frictions do) the laſt Term being equal to 0, inſtead of $\frac{z-a}{n-1}=S$, we ſhall have, ſuppoſing z equal to the firſt Term, $\frac{z-0}{n-1}=S=\frac{z}{n-1}=S$. So that dividing the firſt Term by the Quantity equal to the Friction, the Exponent of the Progreſſion will be had; and dividing the firſt Term by that Exponent Minus 1, the Quotient will give the total Sum of all the Additions to be made on Account of Frictions.

To give Examples of it.

In the 4th Experiment of *Lecture 4. Page 186*, the firſt Term of the Progreſſion is 18, the ſecond 6, whence the Exponent is 3, which Exponent being leſſen'd by 1 is 2, by which Number, dividing the firſt Term

18, the Quotient 9 is the Sum of all the Terms but the First, which is the Power.

So in the Example of *Rule* the 4th, *Lecture* 4th, *Page* 188, the first Term is 108, the second 12, whence the Exponent is 9 and $\frac{108}{9-1} = \frac{108}{8} = 13,5$, which is equal to all the Additions.

So in the other Example, *Page* 189, the first Term is 108, the second 14,333 &c. whence the Exponent is 7,535 &c. and $\frac{108}{6,535} = 15,526$ &c. which is the exact Sum of all the Super-additions.

In the Text I have only taken 7,5 for the Exponent, having neglected the two last Decimals, which gives $\frac{108}{6,5} = 16,61 +$ as in the Body of the *Lecture*, &c.

After these Words (*Page* 261) *That the Parts of Bodies struck, move towards the Blow, is a Consequence of a Law of Nature, which shall be explain'd in my next Lecture*—insert what follows.

Having forgot to take Notice of this in the fifth *Lecture*, I beg the Reader would please to receive the Explanation here.

The common Experiment of laying a Stick with its Ends upon two drinking Glasses full of Water, and striking the Stick downwards in the Middle, with an Iron Bar, so as to break the Stick without breaking the Glasses or spilling the Water, plainly proves the Fact; and the Observation of what happens in the Experiment, shews it to depend upon the first *Law of Nature* (*Page* 284.) For if the Glasses A, B, * have upon their Brims the Stick A B, whose Center of Gravity at C, is struck by a Bar, whose Section is seen at C, the Stick, by the first *Law*, endeavouring to remain in its Place of Rest, has only its Tenacity to resist the Blow of the Bar, which breaks the Stick by the Time it is got to c, and now the two Parts of the Stick must be in the Position ca and cb, and still rise, turning towards C; for M, m, the Centers of Gravity of the two Parts of the Stick, endeavouring to remain in their respective Places, become the Centers of Motion of those two Sticks, whence it happens, that as by the Continuance of the Stroke, they are both push'd downwards at c, their Ends a and b must by a circular Motion be carried upwards and towards C.

This is confirmed by making the Experiment the contrary way; for if you strike the Stick upwards in the Direction c C, you will break the Glasses,

Glasses, if the Blow be strong, and if the Blow be not very quick, yet you spill the Water.

There is also another Experiment, which makes this very evident.

Let the Circle or Hoop of Lead ABC* be hung up by the String*Pl. 16. F. 9. S G, and then struck by the Bar above-mentioned at the Point B in an horizontal Direction; the Effect will be, that the Ring will change its Figure from the Circle A B C D E F to the irregular Oval A b C b D e F g; so that not only the Parts of the Ring at A and C come towards the Blow, while the Bar B (represented by its square Section) moves from B to b, but also the farthest Part of the Ring E comes forward to e, the Points A, C, F, D, becoming Centers of Motion for the Arcs (or Quadrants) BG, BH, EG and EH, to be put into the Positions b g, b b, e g and e b, in which Case they must alter their Figure, because they are not separated from one another at the Points B, H, E, G, &c.

Page 322. line 25, after— odd Numbers, 1, 3, 5, 7, 9, &c. insert

As a great Part of my Auditors, tho' very curious, are unacquainted with mathematical Studies; I always found it very difficult to make them understand the Effect of Gravity in accelerating Bodies in their Fall, by Galileo's Scheme of Triangles, till I had prepar'd them by what I have now said in the 14th Paragraph; but as that Account is not strictly true, I hope this following will satisfy every body.

The Actions or Accelerations of any centripetal Force upon a Body, are (at the same, or equal, Distances from the central Body) always proportionable to the Times; that is, equal in equal Times: And such are those of Gravity (as to Sense) near the Surface of the Earth.

Suppose a Body free to fall in *vacuo*, receives the Impressions or Impulsions of Gravity during a certain Time, as for Example a whole Second; if that Time be divided into a great many small Parts or Intervals (which I call Moments or Instants) the Velocity of the falling Body will increase uniformly or equally at every such Instant, which being all suppos'd equal, the Spaces describ'd in those Instants by the falling Body, must be as the Velocities; that is, as the Terms of an Arithmetical Progression, such as the following.

| <i>Moments or Instants</i> | | | <i>Velocities or Spaces</i> | | |
|----------------------------|---|---|-----------------------------|---|--------|
| 1st | — | — | — | — | 1 |
| 2d | — | — | — | — | 2 |
| 3d | — | — | — | — | 3 |
| 4th | — | — | — | — | 4 |
| 5th | — | — | — | — | 5, &c. |
| O o o | | | | | |

The

The first Term of which is the Velocity of the falling Body, at the End of the first Instant acquir'd by the Sum of the Impulsions of Gravity during that Instant. The last Term will be the Velocity of the falling Body at the End of the last Instant of the Fall: And in the like manner the middle Term of the Progression will express the Velocity acquired at the End of Half the Time of the Fall, by the Sums of the Impulsions of Gravity during Half the Time of that Fall. And since the Spaces describ'd in the same Time, are as the Velocities, the Sum of the Terms of the Progression will express the whole Space describ'd during the whole Time of the Fall. Now by the Doctrine of Progressions, it is universally known, that the mean Time multiplied by the Number of Terms, gives a Product equal to the Sum of the Progression; that is, in the present Case, the mean Velocity (which is the Velocity, that the Body has acquir'd at the End of Half the Time of its Fall) multiplied by the whole Time of its Fall, will give a Space equal to the Spaces describ'd by the Body during that Time, with an uniformly accelerated Motion.

That is in other Words,

That, if the Body was to move uniformly during the whole Time of the Fall, with that mean Velocity it has, when it has fallen half that Time; it would describe a Space equal to that, which it describes during the whole Time, when it sets out from Rest, and has its Velocity accelerated all the while.

Now since we have already said, that the Velocities, or the Accelerations of the falling Body are always proportional to the Times; the ultimate Velocity at the End of the Fall, must be double that of the mean Velocity, acquir'd at the End of Half the Time of the Fall; and consequently, if that last, or greatest Velocity, be multiplied by the whole Time of the Fall, the Product will be the Space, which the Body would describe with that last Velocity in the same Time, and with an uniform Motion, which is double of the Space describ'd during the same Time, with the mean Velocity, or its Equal, viz. the Space describ'd during the same Time by an accelerated Motion.

'Tis therefore evident, that after a Body is fallen freely during a certain Time; it has acquir'd, by the equal and uniform Accelerations of Gravity, a Velocity, such as would make it describe (in a Time equal to the Time during which it fell) a Space, double of that Space it went through in the same Time, with a Motion uniformly accelerated.

From hence, it will plainly appear, how the Motion of falling Bodies is accelerated; and the Spaces they describe in any equal given Times, are as the odd Numbers 1, 3, 5, 7, &c. and the total Spaces as the Squares of the Times. For let a Body *in vacuo* be suppos'd to fall by Gravity during the Time of 4 Seconds; and if it be found, that at the End of the first
Second,

Second, it has describ'd a Space equal to one Rod; was Gravity to act no longer upon it, it has already acquir'd by uniform Accelerations, such a Velocity at the End of that 1st Second, as would make it describe a Space equal to two Rods during the next Second with an uniform Motion. But Gravity continuing to act during the second Second, the Sum of its Actions or Impulsions will be (near the Surface of the Earth, where the Distance from the Center is not sensibly alter'd) equal to that of the Impulsions in the first Second, and therefore the Body will describe three Rods in the next Second: Now to find the Velocity of the falling Body at the End of the second Second; we may consider, that since from Gravity it acquir'd an uniform Velocity of two Rods *per* Second, by the Actions or Impulsions receiv'd during one Second, it will acquire a double Velocity, *viz.* of four Rods *per* Second of Time, after it has fallen, and been acted upon by Gravity, during two Seconds of Time; so that in the third Second (instead of the four Rods it would describe, was Gravity to cease to act) it will describe five Rods, because during that third Second, the Impulsions of Gravity are also suppos'd equal to those it receiv'd in the first. In the same Manner, and for the same Reason, was Gravity to cease to act, after having acted three Seconds, the Body would have a Velocity to describe six Rods in one Second (triple of two Rods *per* Second, which is the Velocity, that Gravity gives in the Impulsions of a Second) but Gravity acting as before, during that fourth Second, the Body will, by the Addition, or Sum of all those joint Impulsions, describe seven Rods; so that the Times and Spaces will be as follows.

| <i>Number of each particular Second.</i> | <i>Rods.</i> | |
|--|--------------|---|
| 1 = 1 st - - - - - | 1 | } <i>or Spaces
describ'd
in each
of them.</i> |
| 1 = 2 ^d - - - - - | 3 | |
| 1 = 3 ^d - - - - - | 5 | |
| 1 = 4 th - - - - - | 7 | |
| <hr/> | | |
| 4 Seconds - - - - - | 16 | <i>Rods</i> |

And it evidently appears from this Progression, that the total Spaces from the Beginning of the Fall, are as the Squares of the Times spent, during which the Bodies fall; for in one Second the Spaces describ'd are 1 Rod; in 2 Seconds, $1 + 3 = 4$ Rods (2×2); in 3 Seconds, $1 + 3 + 5 = 9$ Rods (3×3); and in 4 Seconds, $1 + 3 + 5 + 7 = 16$ Rods (4×4). Therefore it is true, that the Spaces describ'd by falling Bodies, are in a duplicate *Ratio* of the Times or Velocities; and on the contrary, that the Times of the Fall or the Velocities acquir'd at the End of those Times, are in a sub-duplicate Proportion of the Spaces, or as the square Roots of those Spaces.



AN ALPHABETICAL

I N D E X

FOR THIS

FIRST VOLUME.

A

- | | | |
|---|---------------|---|
| A bsolute Force | Page 32 | Animalcula, more of them in the Milt of one |
| Acceleration of Bodies in Motion | | Cod-fish, than there are Inhabitants upon |
| — applied to the Fall of Bodies | 317 to 321 | the whole Earth |
| Accelerating Force | 322 | — how inconceivably small the Globules of |
| Action of Gravity on the Planets and Comets | 32 | their Blood |
| — of the Sun and Moon to move the Sea | 350, 351, 352 | — yet immensely bigger than the Particles |
| — of the same on Planets and Comets | 365, 366 | of Light |
| — equal and contrary to Re-action | 355 | Arm, the Force of the Muscles to move it |
| Æquilibrium | 90 | to 159 |
| Æquinoctial Tides | 367, 368 | <i>Archimedes</i> , his Problem for raising the Earth, |
| Air, its Resistance | 343 | with a Calculation of it |
| Angle of Application to a Leaver or Balance | 141 to 145 | Areas, astronomical, explain'd |
| — of Traction | 160 to 163 | — discover'd by <i>Kepler</i> to be proportionable |
| | | to the Times of Description |
| | | — prov'd from Sir <i>Isaac Newton</i> |
| | | 404, 405, 406 |
| | | Astronomical Observations, by the Rev. Mr. |
| | | <i>Bradley</i> , <i>Savilian Professor</i> at <i>Oxford</i> , |
| | | proving the Motion of the Earth |
| | | 383 to 388 |
| | | <i>Aphe-</i> |

An Alphabetical I N D E X.

- Aphelion, Planets and Comets move slower in it Page 348
- Atoms, what 4
- Attraction, mutual of Bodies 7
- Experiments to shew how it acts 7, 8
- the duplicate reciprocal Proportion of its Action 31
- shewn by a Candle's Light 9, 10
- Attraction of Cohesion 10
- strongest in Contact *ib.*
- very considerable in Chrystal Buttons and Balls of Lead 183, 449
- Experiment of it with Oil of Oranges, between two Glass Planes 11
- with Fluids rising in small Tubes of Glass, in Bread and Loaf-Sugar 12
- with colour'd Spirit of Wine rising between two Glass Plates in the Figure of an Hyperbola 13
- prov'd by Experiments of Quicksilver 14
- of Iron Wire lying at the Bottom of a Vessel of Quicksilver, and likewise Silver Wire, when its Surface is foul 15
- prov'd by the Practice of Soldering 16
- it makes Drops of Water globular *ib.*
- its Laws *ib.*
- Attraction, magnetick 16, 40
- electrick 17, 18, 19
- not an occult Quality 21
- Aurora borealis* 31
- Axel in the Wheel, or *Axis in Peritrochio* 102
- Experiment upon it 102 to 105
- a further Consideration of it 118, 128
- compounded 129
- join'd with a Screw 130
- reduc'd to a Leaver 132
- Friction of it 137, 186
- Monsr. Perrault's new *Axis in Peritrochio* worse than the common Sort, on account of the Friction of the Ropes 230, 231, 232
- the same further consider'd 253, 254
- Axel-trees of Carriages must be streight, and at right Angles to the Pole 203, 204, 205
- ## B
- B**alance, being either a Beam and Scales, or a Steelyard 91, 92
- deceitful or false 95
- reduc'd to a Leaver 96
- how to make a very exact Pair of Scales 135
- how to suspend its Beam 138, 139, 140
- may be false, even when the *Brachia* are equal 140
- how to estimate the Force of Weights acting obliquely upon it 141, 142, 143, 144
- a particular Make of one, on which equal Weights make an *Æquilibrium* at unequal Distances 146, 147
- Effect of an oblique Push of a Power against its Beam 147, 148, 149, 150
- Balls of Lead cohere very strongly 183, 449
- Battering Ram 44, 45
- the Stroke of it compar'd to the Stroke of a Cannon-Ball 63, 64
- its Action consider'd 332, 337
- Bigness of a Comet 355
- Body, when struck, its Parts move towards the Blow 460, 461
- Bodies, their Particles in some Cases repel one another 17
- their Weight on the Moon's Surface 33
- their Weight proportionably greater on the Surface of small Planets than large ones, considering the Quantity of Matter in those Planets 34
- inclin'd, do not fall when their Line of Direction goes through their Base 57, 58
- at Rest, or in Motion, do not fall if their Center of Gravity be supported 59, 60
- kept from falling by adding other Bodies to them 61
- rolling and sliding on inclin'd Planes 86
- Bones of Animals reduc'd to Leavers, and the Manner that the Muscles act upon them, taken from BORELLI 153 to 159
- Block, or Blocks 99
- Bow, or Spring, an Account of it 375 to 379
- Brachia* of a Balance or Leaver 47, 91
- Bucket-Engine for raising Water 76
- ## C
- C**annon, the Action of its Ball compar'd with that of a battering-Ram 63, 64
- the Force exerted by the Powder on the Breech of a 24 Pounder 417, 418
- Carriages, consider'd, and Experiments concerning them 190 to 229
- with four Wheels more advantageous than with two 209 to 212
- Experiments to prove it 213, 214, 216
- Corollaries from those Experiments 217 to 229
- a new Sort to carry great Weights of Stone down-Hill, made use of by Mr. Allen at Bath, describ'd 274 to 279
- Caution to Persons concern'd in Engines 68, 69
- Center

An Alphabetical I N D E X.

| | |
|--|------------------------------|
| Center of Motion | Page 49 |
| Center of Gravity | <i>ib.</i> |
| — a mechanical Way of finding it | 73 |
| — a mathematical Way of finding it | 74 |
| — where that of a Windmill, and that of a Crane must be plac'd | 77 |
| — common of two or more Bodies | 52, 53, 54 |
| — of the Sun and Planets | 83 |
| — of a Load of Hay, or a Load of Iron | 87 |
| — of Oscillation and of Percussion | 373 |
| — how to find the Center of Oscillation | 455 |
| Centrifugal Force, what | 302 |
| Centripetal Force, what | 301, 302 |
| Chronometer, contriv'd by Mr. George Graham, for measuring Time to the 16th Part of a Second | 373, 374 |
| Cold and Heat, their Effects on the Rods of Pendulums | 373 |
| Cohesion, Attraction of | 10 |
| Comet, retrograde destroys the Hypothesis of Vortices | 354 |
| — may at Length fall into the Sun | 355 |
| — about the Bigness of the Moon | <i>ib.</i> |
| Comets, a general Account of them | 409, |
| — Sir Isaac Newton's Account of them | 410, 411 |
| — to 417 | 411 |
| — by a Ray drawn from the Sun, describe Areas proportionable to the Times | 446 |
| — to 448 | 448 |
| Compound Engines | 117 |
| Compound Motion of a Body acted upon at the same Time by two or more Forces | 124, 125, 286 to 290 and 380 |
| Cone, double rolling upwards | 55, 57 |
| Cranes | 119, 120, 121 |
| — the Gibbets of them inconvenient in the common Way | 178 |
| — Mr. Allen's Cranes considerably improv'd | 179, 180, 181 |
| Cycloid, its Description | 377 |
| — some of its Properties | 377, 378 |
| — Swings of a Pendulum in it are isochronal | 378 |
| — Experiments with cycloidal Channels | 378, 379 |
| Cycloids with curv'd Bases, or Epicycloids, some of their Properties | 419, 420 |
| Cylinder loaded, so as to roll up an inclin'd Plane by the Descent of its Center of Gravity | 55, 77 to 83 |
| — a further Consideration of its Properties | 451, 452, 453 |

D

| | |
|---|-----------|
| D esire of Victory in Disputes, hinders the Improvement of natural Knowledge | Page 1, 2 |
| Diameters and Distances of the Planets | 432 |
| Direction, Line of | 433 |
| Divisibility, of Matter <i>in infinitum</i> , but Matter not actually divided <i>in infinitum</i> | 72 |
| — of Quantity, demonstrated to be <i>in infinitum</i> | 4 |
| Drop of Oil of Oranges rising between two Glass Planes | 25 |
| — fix Properties of it | 11, 12 |
| Drops of Water made globular by Attraction | 37 |
| Ductility of Gold | 36, 37 |
| | 28 |

E

| | |
|---|-----------------|
| E arth, higher at the Æquator than towards the Poles, and the Reason why it ought to be so | 373, 374 |
| — an Experiment to illustrate this | 374, 375 |
| — the Earth's annual Motion prov'd by the Observations of the Rev. Mr. Bradley, Savilian Professor of Astronomy at Oxford | 383, 386 to 388 |
| — the Earth's Figure not an oblong, but an oblate Spheroid | 428, 429 |
| Effluvia, those of <i>Asa fætida</i> inconceivably small | 28 |
| — of Comets and Planets in the celestial Spaces, exceeding rare | 354 |
| — very considerable in the Sun's Atmosphere | <i>ib.</i> |
| Electricity | 17, 18, 19 |
| — prov'd by a rubb'd Tube attracting and repelling Feathers and Leaf-Gold, both in the Air and <i>in Vacuo</i> | 20 |
| — prov'd by an exhausted Glass Globe giving a purple Light when rubb'd | 21 |
| — several electrical Experiments by Mr. Stephen Gray | 450, 451 |
| Engine, to raise Water by a gaining and a losing Bucket | 76 |
| — Mr. Richard Newsham's, very convenient and useful for extinguishing accidental Fires | 247 |
| — newly contriv'd to try and estimate the Strength of different Men, as also of different Limbs in the same Man | 282, 283 |

Faculties,

An Alphabetical INDEX.

F

Faculties mechanical Page 88
 Flames of Spirit of Wine us'd in the Pyrometer 423 to 426
 Fly, its Use and Application 245, 246
 — it adds nothing to the Force of the Power, tho' it helps the Motion 246
 — join'd to a Screw, for coining Money 334, 335
 — how to calculate the Force of such a Machine 339
 Figure of the Earth 373, 374, 428, 429
 — of a Portion of a Fluid cannot be alter'd by any Pressure from another external Fluid 30
 Force, centrifugal 301, 302
 — accelerating 32
 — centripetal 31, 302
 — Experiments to explain those Forces 302 to 305
 — of Inactivity 284, 286
 — moving 33, 43
 Force of Men best applied in rowing with Oars 247
 — very ill applied in rowing with an horizontal Axis, having vertical Oars like the Spokes of a Wheel *ib.*
 — very great, when exerted by the Muscles, that extend the Legs and Thighs 248
 — compar'd by a new Contrivance 282, 283
 — of five Men equal to that of one Horse 254
 — of Men and Horses consider'd and compar'd 241 to 248
 — the same by *Monf. de la Hire*; with Remarks upon his Dissertation 267 to 273
 — *Philosophical Transactions* quoted concerning the Dispute about the Force of moving Bodies 339
 — of the Action of the Powder on the Breech of a Cannon shooting out a 24 Pound Ball 417, 418
 Forces requir'd to bend Ropes, &c. 229 to 241
 Friction, makes Sealing-wax transparent 40
 — not sufficiently consider'd by the common Contrivers of Machines and Engines 133
 — unavoidable 133, 134
 — very small in a Leaver 134
 — how to be consider'd in a Balance 134
 — very great in Pullies 136, 137
 — not very great in the *Axis in Peritrochio* 137
 — not in the inclin'd Plane 137
 — yet very great in the Wedge and Screw 137, 138

— of mechanical Engines, and of Ropes, further consider'd Page 182 to 248
 — not according to the Quantity of the touching Surfaces, but of the Weights pressing them together in their Motion 184, 185
 — Resistance arising from it 185
 — its Quantity in a great Beam drawn along the Ground 185, 186
 — how to find it in Gudgeons or Pivots 186, 187
 — in compound Engines 187, 188, 189
 — a Table of the Friction of Sledges 193
 — of Carts and Waggons 213
 Friction of the Sheeves and Ropes of a Tackle of three 236 to 239
 — Quantity of Friction equal to about one third of the Weight 249
 — how to diminish it in Wheel Carriages 249 to 251
 — a Machine to shew how much it is on different Surfaces, and how to diminish it in the Axels of Wheels 251, 252, 253
 — a more exact Method of calculating its Quantity 458, 459, 460

G

Gibbet of a Crane, Inconveniencies of it in the common Way 178
 — remedied by *Mr. Padmore* of *Bristol* 179
 Gun-lock consider'd 454
 Gun-powder, a small Quantity of it will shake an Acre of Ground 403
 — its Action on the Breech of a 24 Pounder 417, 418
 Gravity, Gravitation or Attraction defin'd 6
 — not an occult Quality 21
 — Center of Gravity 49
 — a mechanical Way to find it 73
 — a mathematical Way 74, 75, 76
 — Center of it in a Crane, and in a Windmill 77
 — in a Load of Hay, or a Load of Iron 87
 Gravity, or Gravitation, from the Surface to the Center of a Planet, decreases directly as the Distance from the Center 35
 — its Action on Planets and Comets to make their Motions regularly accelerated and retarded 150, 151, 152
 — mutual between the Earth and its Parts 417
 Gray, *Mr. Stephen*, his electrical Experiments 450, 451

Hammer

An Alphabetical INDEX.

H

- H**ammer Page 33
 — in its Use it is better to double the Velocity with the same Quantity of Matter, than to double the Matter keeping the same Velocity, but the Reverse in the Use of the Ram 388 to 400
 Heat, its Effect on Pendulums 373
 — Quantity of its Effect on Metals to dilate them in their Dimensions 423 to 426
 Horses, a mechanical Consideration of their walking 84, 85
 — draw with great Disadvantage, walking in a small Circle 242
 — likewise going up Hill 243
 — best of all horizontally and in a freight Line *ib.*
 — Strength of one Horse equal to that of five Men 254
 Hyperbola, made with rising Water between two Glass Planes or Plates 13
 — but in inverted Order with Quicksilver 40

I

- J**ACK 122
 Inclined Bodies do not fall, when their Line of Direction goes through their Base 57, 58
 Inclined Plane 105, 130
 — Experiments upon it 106
 — reduced to a Leaver 131
 — its Manner of acting, and that of a Wedge solved by a bended Leaver 159, 160, 161
 Inclined Towers not falling 58
 Inertia or Inactivity, its Force 284, 285
 Infinites are not all equal 25, 26
 Instruments in the common Use of Life reduced to Leavers of several Kinds 150, 151
 Intensity of a Power 48, 49
 Jockey, a strong Kentish Man 255

L

- L**abourers, *English*, generally stronger than *French* or *Dutch* 254
 Lamp, rolling 57
 Laws of Motion from Sir *Isaac Newton* 284
 — 1st Law explain'd and illustrated 284 to 317

- 2d Law explain'd, &c. P. 317 to 356
 — 3d Law explain'd, &c. 356 to 359
 — applied to account for the Tides 350 to 368
 Leaver, its several Kinds 96, 97, 119
 — homodromous and heterodromous 150
 — several Experiments upon it 97, 98
 — bended 98
 Length of a Pendulum to swing Seconds 373
 Light, its Resistance 3
 — its Density decreases, as the Squares of the Distances of the enlighten'd Bodies increase 35
 — is repell'd by Glass and other Bodies 41
 Line, parabolical 292
 — of Direction 72
 Load of Iron, its Center of Gravity 87
 Load-stone, well consider'd by Dr. *Petrus van Muschenbroek* 449

M

- M**achine, to shew whither the Friction depends upon the Quantity of Surface rubbing, or the Quantity of the Weight pressing the Surfaces together, as also how to diminish it in the Axes of Wheels, &c. 251, 252, 253
 — call'd a Pyrometer 421, 422, 423
 — a new one to try the Strength of Men 282, 283
 — newly contriv'd by the Author, to shew mechanically, how Comets and Planets, by a Ray drawn from the Sun, describe Areas proportional to the Time 440 to 448
 — a curious one at *Darby* for winding Silk 69, 70
 Mass 43
 Matter, what 2
 — in what Sense understood by the *Cartesians* *ib.*
 — the same in all Bodies 3
 — divisible, but not actually divided, in infinitum 4
 — trac'd through the natural and artificial Changes of Bodies, and shewn to be the same in them all 22, 23
 — a small Quantity of it may fill the Sphere of Saturn 26, 27
 — is divided into inconceivably small Parts 28
 — less of it in the World than *Vacuum* 30, 31
 — Subtile of the *Cartesians*, no Proof of a *Plenum*

An Alphabetical I N D E X.

- Plenum*, tho' its Existence were granted — an Experiment to illustrate compound Motion further, and at the same Time remove an Objection made against the Motion of the Earth Page 291
- Mechanicks Page 65
- Mechanical Organs, Powers, or Faculties 89, 90
- Powers all united in one Machine 123
- Men, their walking, mechanically consider'd 84, 85
- have but small Force to draw a Barge, or to turn a Handle vertically, when it is at the Top, or at the Bottom of the Revolution 244
- are much help'd by a Fly in turning a Handle 246
- the Force of Five equal to that of one Horse 254
- the Force of Men and Horses consider'd according to the different Applications of it 241 to 248
- *Monf. de la Hire* Dissertation on that Subject, consider'd 267 to 273
- Mediums, their Resistance occasionally consider'd 340, 341
- Mercury, Resistance of it 24
- attracted a little by Glafs, but very much by it self 39, 40
- attracts it self into a Hyperbola between two Glafs Planes 40
- several Experiments of its Attraction made by *Dr. James Jurin* 38, 39, 43
- Molecule*, what 4
- Momentum*, Quantity of Motion, and moving Force, the same 43
- made up of the Mass, multiplied into the Velocity 327, 328
- of falling Bodies proportionable to their Velocities, but not to the Height from which they fall 327, 328
- new and old Opinion about it compar'd 393
- Moon, Weight of Bodies on its Surface 33
- raises the Waters of the Sea where it is vertical, and also at the Antipodes of that Place 360, 361
- the Action of it consider'd hydrostatically 362
- Objections to that Doctrine answer'd 362, 363
- Motion, distinguish'd from Velocity 44
- Center of 49
- defin'd in general and in particular 63
- Quantity of it 43
- compound, when a Body is acted upon by two, or more, Forces at once 124, 125, 286 to 290, 380
- perpetual, shewn to be impossible 175 to 178
- its *Laws* from *Sir Isaac Newton* 284
- of Projectiles applied to the Moon's Motion 300, 301
- Experiments to shew, that Gravity and the projectile Force, make the Planets continue their Motions in their Orbits 309 to 317
- Acceleration of Motion explain'd 317 to 331 and 457, 458
- of Bodies on inclin'd Planes, and the Doctrine of Pendulums deduc'd from it 329 to 331, and 369 to 375
- of its 1st Law explain'd 284 to 317
- 2d Law explain'd 317 to 356
- 3d Law explain'd 356 to 369
- Laws of Motion and Gravity applied to account for the Tides 359 to 368
- Moving Force 33
- Muscles, acting upon the Bones to move them 153 to 158
- of the Arm, *biceps* and *brachii*, their Force calculated 154 to 159
- extensors of the Legs and Thighs, how to apply them most usefully 266

N

- N**ap-Tides 365, 366
- New Terms should always be explain'd 2
- New and old Opinion about the Force of moving Bodies 393
- Proof of the New, *viz.* that the *Momentum* or Force is as the Mass multiplied into the Velocity, by Experiments of a flat Pendulum, &c. 394 to 296
- Experiments seeming to disprove, but really confirming the new Opinion 397
- Newsham*, his Engines very useful 247
- Newton*, *Sir Isaac*, his Way of demonstrating the Operations of all the mechanical Powers by one Scheme 124 to 126
- his Account of Comets 411 to 417

O

- O**bservations of several fix'd Stars compar'd, proving the Motion of the Earth, by the *Rev. Mr. Bradley*, *Savilian Professor of Astronomy at Oxford* 283 to 388
- Occult Qualities, not admitted in the *Newtonian*

An Alphabetical I N D E X.

nian Philosophy Page 21, 42
 Opinion, old and new about the *Momentum* 348
 Orange-Oil, a Drop of it moving between two
 two Glafs Plates or Planes 37
 — fix Properties of it, as it riles between
 thofe Planes 38
Orreries, Machines fo call'd, commonly very
 faulty, as they give falfe Ideas of feveral
 celeftial Phænomena 433
 Oscillation, Center of it, fometimes the fame
 as the Center of Percuffion 373
 — how to find it 455

P

Pail of Water is kept from falling by
 fupporting its Center of Gravity, tho'
 the Stick that fupports it would fall of it
 felf 62
 Parabola, the Line in which Bodies projected
 horizontally, or obliquely, move 292 to
 300
 Parabolical Motion of a Cannon-Ball 292, 300
 Parallax, of the fix'd Stars, not to be obferv'd
 accurately enough by Inſtruments 384
 Pendulums, Motions of them deduc'd from the
 Motion of Bodies on inclin'd Planes 229
 to 331, and 369 to 375
 — their Rods lengthen and fhorten by Heat
 and Cold 373
 — their true Length for Seconds 419
 — Percuffion, Center of it 373
 — why oblique and direct Percuffion are
 miftaken for one another 380 to 382
Peritrochio, Axis in 102
 — Monf. *Perrault's* new one worfe than in
 the common Way, on Account of the
 Friction of the Ropes 230 to 232
 — the fame further confider'd 253, 254
 Periods of the Planets—See Plate 30.
 Perpetual Motion, impoffible 176 to 178
 Perihelion 348
 Piles, Rammer to drive them 336, 337
 Plane, inclin'd 105
 Planets, their Motion in their Orbits, fwifter
 in the Perihelion, and flowier in the A-
 phelion 348
 — can't preferve their Motions in a refift-
 ing Medium 353
 — not carried round in corporeal Vortices
 388 to 390
Planetarium, a new Machine for fhewing the
 Motion and feveral Appearances of the
 celeftial Bodies 430 to 448
 — thofe Parts of it, which fhew the gene-
 ral Syſtem, and the particular Syſtems
 of Jupiter and his Satellites, and Saturn,

and his Satellites, defcrib'd Page 433
 to 437
 — thofe Parts of it, which fhew the Mo-
 tions of the Earth and Moon about the
 Sun; the Eclipfes and all the refpective
 Phænomena of thofe three Bodies, def-
 crib'd 437 to 446
 Porters, at the Custom-Houfe Key carrying
 great Weights 263 to 273
 — in *Turkey* carrying more Weight than in
England 254 to 262
 Power, defin'd 48
 — the Word improperly applied 127
 Powers, mechanical reduc'd to the Leaver
 127
 Prefſure, of two Bodies on the Axis of a fin-
 gle Pulley, and alfo on the Axis of a
 double Pulley, or *Axis in Peritrochia*
 166, 167, 455
 Projectiles, move in a Parabola 292 to 300
 — the Laws of their Motion apply'd to that
 of the Moon 300, 301
 — moving perpeudicularly or obliquely
 upwards 325 to 327
 Pulley 99
 Pullies, upper and lower 100, 101, 118
 — their Friction very great 136, 137
 — combin'd 168
 — Tackles of Four, Five, or Six 100
 — the Friction of a Tackle of Three, and
 one of Five calculated 236 to 241
Pyrometer, for meafuring the Effect of Heat
 and Cold upon Metals, tho' they ſhould
 only expand or contract in Length the
 the 20000 Part of an Inch 421, 422
 423

Q

Qualities, occult, not admitted by the
Newtonians 42
 Quantity, Demonſtration of its Diviſibility in
infinitum 25
 Quickſilver attracts Quickſilver and Glafs;
 but it attracts it felf more than it does
Glafs 39

R

RAM, battering; Stroke of it compar'd
 to that of a Cannon-Ball 45, 63,
 64
 — further confider'd 332, 337
 Rammer to drive Piles 336, 337
 Rat-tail'd Crane 122
 Repulſion in Bodies 17
 Reſiſtance, of Mercury, Water, Air, Light,
 and other Fluids 3

An Alphabetical I N D E X.

| | |
|---|---|
| <p>— arising from Friction Page 182 to 248</p> <p>— from the bending of Ropes, consider'd
and calculated 229 to 241</p> <p>— of Mediums occasionally consider'd 340,
341</p> <p>— of Water compar'd with that of Wood
401</p> <p>— greater than that of Lead in one Cafe
402</p> <p>— greater than that of Wood, when ac-
ted upon by Gun-powder 402, 403</p> <p>Rolling Bodies 60, 86</p> <p>— Lamp 57</p> <p>Ropes, their Friction 229 to 241</p> <p>— Experiments upon their Resistance 233,
234</p> <p>— running Rope 99</p> <p>— the easiest Way of breaking a Rope
260, 262</p> | <p>Steelyard Page 91, 92</p> <p>Strength, cannot be gain'd without Loss of
Time and Space 67, 68</p> <p>— of Men and Horses compar'd 241 to
248</p> <p>— pretended Feats of it wholly owing
to Art 255, 256 257</p> <p>— accounted for, as not exceeding the
Power of any Man of moderate Strength
257 to 262</p> <p>— real Feats of extraordinary Strength
perform'd by <i>Thomas Topham</i> 280,
281</p> <p>Suppositions necessary for understanding Me-
chanicks 89, 90</p> <p>System of the World consider'd 432, 433</p> |
|---|---|

T

| | |
|--|---|
| <p style="text-align: center;">S</p> <p>Scales, how to make a very exact Pair
135 to 241</p> <p>Screw 112</p> <p>— its Force calculated 113, 114</p> <p>— its great Friction 114</p> <p>— a particular Advantage of it 115</p> <p>— applied to Percussion 116</p> <p>— join'd with a Wheel 119</p> <p>— Thread of it, square and sharp, com-
par'd</p> <p>Scorpions (mechanical Engines) compar'd with
Cannons 70, 71, 174, 175</p> <p>Sealing Wax, made transparent by Friction
42</p> <p>Simple Machines 88</p> <p>Sledges, Experiments upon them 193 to
196</p> <p>Sliding Bodies on inclin'd Planes 86</p> <p>Sling 335, 336</p> <p>Solstitial Tides 367, 368</p> <p>Space of 189 291 996 583 668 465</p> <p>200 Miles, which a Power, equal to the
Strength of one Man, must go through
to raise the Earth, but one Inch, accord-
ing to <i>Archimedes's</i> Problem 67</p> <p>Space of Time for the Performance, if the
said Power mov'd as swift as a Cannon-
Ball, will be 26 978 123 942 460</p> <p>Years 66, 67</p> <p>Spokes, of Wheels must be inclin'd to the
Naves 203</p> <p>Spring and Neap Tides caus'd by the Action
of the Sun and Moon, according to
their different Positions 365, 366</p> | <p style="text-align: center;">T</p> <p>TABLE, of the Friction of Sledges
293</p> <p>— of the Friction of Carts and Waggons
213</p> <p>— of the Forces requir'd to bend Ropes of
different Diameters, stretch'd by different
Weights round the Rollers of different
Diameters 235</p> <p>— of the Effect of Heat and Flame on
Metals measured by the Pyrometer 424,
426</p> <p>Tackle, of Three, a Calculation of the
Friction of its Sheeves and Ropes 236,
237</p> <p>— of Four, of Five, and of Six 100</p> <p>— Experiments of the Friction of the Tac-
kle of Five 240, 241</p> <p><i>Testudo</i> of the Roman Soldiers; a Conjecture
about it 263</p> <p>Tides, why daily retarded 364</p> <p>Times, periodical of the Planets experimen-
tally compar'd with their Distances 391,
392</p> <p><i>Topham, Thomas</i>, real Feats of Strength per-
form'd by him 280, 281</p> <p>Towers, inclin'd not falling 58</p> <p>Traction, Angle of it consider'd 160 to
163</p> <p>Tube, of Glass rubb'd, a remarkable Pro-
perty of it 41</p> |
|--|---|

V

| | |
|---|--|
| <p style="text-align: center;">V</p> <p>Vacuum, prov'd by Experiment 5</p> <p>— much more of it than Matter
13</p> <p>— inter-</p> | |
|---|--|

An Alphabetical I N D E X.

| | | |
|--|---|------------|
| — interspers'd, prov'd from the different | Wedge, defin'd | 107 |
| Weight of Bodies of equal Bulk | — further consider'd, and Experiments of | |
| Velocity, what | its Manner of Action | 107 to 112 |
| — of a Weight consider'd differently from | Weight defin'd | 48 |
| that of a Power | — of Bodies, on the Moon's Surface | 33 |
| <i>Vis inertia</i> | — of Bodies, greatest on the Surface of small | |
| <i>Vortices, Cartesian</i> , disprov'd by Reason and | Planets in Proportion to their Quantities | |
| Experiments | of Matter | 34 |
| | Weights acting obliquely upon a Leaver | 143 |
| | | to 145 |
| | Wheel-Carriages consider'd | 201 to 229 |
| | Wheels, high and low, compared | 171 to |
| | | 174 |
| | — great, more advantageous than little, | |
| | on any Ground whatever | 208, 209 |

W

| | |
|-----------------------------------|------------|
| W Aggon-Way | 189 |
| — Mr. Allen's describ'd by Mr. De | |
| Labelye | 174 to 179 |

F I N I S.

N. B. **T**HE Course of Experimental Philosophy, of which this First Volume contains Half, is perform'd by the Author at his House in *Channel Row, Westminster* (where Catalogues of the Experiments may be had) on such Days, and at such Hours, as shall be agreed upon by the Majority of the Auditors. *N. B. Every Auditor is to pay Three Guineas, when the Number is not less than twelve Persons; but any Three or Four, may any one Person, may have a Course to themselves paying the Price of Twelve.*

A short but full Course of Astronomy, will also be perform'd by Means of the PLANETARIUM, to any Number of Persons, not less than Ten, at a Guinea each; or to any less Number, who are willing to pay 10 Guineas, upon their giving a Day's Notice.

A D V E R T I S E M E N T.

THERE is now made by *John Senex*, Fellow of the Royal Society, a large Pair of Globes, 28 Inches Diameter, fit to adorn the Libraries of the Curious. On the Terrestrial are inserted, all the Discoveries and Observations hitherto made; and on the Celestial are placed all the Stars in *Mr. Flamsteed's* Catalogue, as publish'd by *Dr. Halley, &c.* being above 2000 more than ever were inserted upon any Globe. The Asterisms are designed, so as to answer the Description of the Antients; and the Letters of Reference, made Use of by *Bayer* in his Tables, are inserted. Also Globes of 17, 12, and 9 Inches Diameter; *Dr. Halley's* Zodiack; *Mr. Whiston's* Solar System, and two Pair of large Hemispheres, one on the Plane of the Equinoctial, and the other on the Plane of the Ecliptick, containing all the Stars in *Mr. Flamsteed's* Catalogue. These, with all the best Maps of the World, and Quarters, and all the general Divisions of *Europe*, are Sold by him, at the *Globe*, over against *St. Dunstan's Church* in *Fleetstreet*.



*BOOKS printed for J. SENEX, W. INNYS and R. MANBY,
and J. OSBORN and T. LONGMAN.*

THE Philosophical Transactions (from the Year 1720 to the Year 1732) abridged and disposed under general Heads. By Mr. REID and JOHN GRAY, A. M. F. R. S. In two Vol. Being a Continuation of the Abridgment publish'd by LOWTHORP, JONES, or MOTTE.

This Abridgment by Mr. REID and Mr. GRAY, being begun by our Approbation, and encouraged by the Assistance of many Members of the ROYAL SOCIETY, who have revised and improved their own Papers; we believe the same from our Knowledge of the Abilities of the Compilers, to be well executed.

JOHN ARBUTHNOT, M. D. F. R. S.
ROBERT BARKER, M. D. F. R. S.
JAMES BRADLEY, Prof. Astron. Savil. Oxon.
F. R. S.
GEORGE CAMPBELL, F. R. S.
J. T. DESAGULIERS, L. L. D. F. R. S.
MARTIN FOLKES, V. P. R. S.
JOHN HADLEY, V. P. R. S.

EDMUND HALLEY, L. L. D. Reg. Astron.
Prof. Geom. Savil. Oxon. F. R. S.
WILLIAM JONES, F. R. S.
JOHN MACHIN, Prof. Astron. Gresh. Sch.
R. S.
COLIN MAC LAURIN, F. R. S.
ABRAHAM DE MOIVRE, F. R. S.
JAMES STIRLING, F. R. S.

The Philosophical Works of the Honourable ROBERT BOYLE, Esq; abridged, methodized, and disposed under the general Heads of Physick, Staticks, Pneumatics, Natural History, Chymistry, and Medicine. The whole illustrated with Notes, containing the Improvements made in the several Parts of Natural and Experimental Knowledge since his Time. In three Volumes, 4to. By PETER SHAW, M. D. 1725.

An Introduction to Natural Philosophy: Or, Philosophical Lectures read in the University of Oxford, Anno 1700. To which are added, The Demonstrations of Monsieur Huygens's Theorem, concerning the Centrifugal Force and Circular Motion. By JOHN KEILL, M. D. Savilian Professor of Astronomy, F. R. S. The third Edition. 1733.

Physico-Theology, or a Demonstration of the Being and Attributes of God, from his Works of Creation. Being the Substance of sixteen Sermons, preach'd at St. Mary-le-Bow Church, at the Honourable Mr. Boyle's Lectures in the Years 1711 and 1712, with large Notes, and many curious Observations. By WILLIAM DERHAM, D. D. Canon of Windfor, Rector of Upminster, and F. R. S. the Eighth Edition, 8vo. 1732.

Astro-Theology, or a Demonstration of the Being and Attributes of God, from a Survey of the Heavens. Illustrated with Copper Plates. By WILLIAM DERHAM, D. D. Canon of Windfor, and F. R. S. the Sixth Edition, 8vo. 1731.

The Religious Philosopher: Or, The Right Use of Contemplating the Works of the Creator; designed for the Conviction of Atheists and Infidels. Translated from the Low-Dutch. By JOHN CHAMBERLAIN, Esq; F. R. S. To which is prefixed, a Letter to the Translator. By the Rev. J. T. DESAGULIERS, L. L. D. F. R. S. The Fourth Edition corrected, adorned with Cuts in three Vols. 8vo. 1730.

Philosophical Conversations: Or, A New System of Physicks, by way of Dialogue. With Eighty-nine Copper Plates. Written in French by Father REGNAULT, of the Society of Jesus. Translated into English and illustrated with Notes, by THOMAS DALE, M. D. In three Vols. 8vo. 1731.

Philosophiæ Naturalis Principia Mathematica. Autore ISAACO NEWTONO, Editio Tertia Aucta & Emendata. 1726.

Opticks, or a Treatise of the Reflections, Refractions, Inflections and Colours of Light. The Fourth Edition corrected. By Sir ISAAC NEWTON, Knt. 8vo. 1730.

Optical

B O O K S printed for J. Senex, W. Innys and R. Manby, &c.

Optical Lectures read in the Publick Schools of the University of Cambridge, Anno Domini 1669. By the late Sir ISAAC NEWTON, then Lucasian Professor of Mathematicks, never before printed. Translated into English out of the original Latin.

Mathematical Elements of Natural Philosophy, confirmed by Experiments; or, an Introduction to Sir ISAAC NEWTON'S Philosophy; written in Latin by WILLIAM JAMES 'S GRAVESANDE, Doctor of Laws and Philosophy, &c. Translated into English by J. T. DESAGULIERS, L. L. D. F. R. S. The Fourth Edition. 2 Vols. 8vo.

An Essay on Perspective. Written in French by WILLIAM JAMES 'S GRAVESANDE, Doctor of Laws and Philosophy, &c. Now translated into English by E. STONE, F. R. S. 8vo. 1724.

The Construction and principal Uses of Mathematical Instruments. Translated from the French of M. BION, Chief Instrument-maker to the French King. To which are added, The Construction and Use of such Instruments as are omitted by M. BION; particularly those invented and improv'd by the English. By EDMUND STONE, F. R. S. The whole illustrated with Twenty-six Folio Copper Plates, containing the Figures, &c. of the several Instruments, *Fol.* 1723.

The Method of Fluxions, both Direct and Inverse. The former being a Translation from the celebrated Marquis DE L'HOSPITAL'S *Analyse des Infiniments Petits*: And the latter supply'd by the Translator, E. STONE, F. R. S. 1732.

The Elements of Euclid; with select Theorems out of Archimedes. By the learned ANDREW TACQUET. To which are added, Practical Corollaries, shewing the Uses of many of the Propositions. The whole abridg'd and publish'd in English by WILLIAM WHISTON, Mr. Lucas's Professor of the Mathematicks in the University of Cambridge. The Third Edition. 8vo. 1727.

An Analytick Treatise of Conick Sections, and their Use for resolving of Equations in determinate and indeterminate Problems: Being the Posthumous Works of the Marquis DE L'HOSPITAL, Honorary Fellow of the Academy of Royal Sciences. Made English by E. STONE, F. R. S. 1723. 4to.

Geometria Organica: Sive Descriptio Linearum Curvarum Universalis. Auctore MAC LAURIN, Matheseos in Collegio Novo Abredonensi Professore, & Reg. Soc. Soc. 4to 1720.

Dissertationes Physico-Mathematicae, partim antea Editae in Actis Philosophicis Londinensibus, jam Auctiores & Emendatiores, partim nunc primum impressae. Auctore JACOBO JURIN, M. D. 1732.

Pharmacopœia Officialis & Extemporanea, or a compleat English Dispensatory, in four Parts. Containing, I. The Theory of Pharmacy, and the several Processes therein. II. A Description of the officinal Simples, with their Virtues and Preparations, Galenical and Chymical. III. The officinal Compositions, according to the last Alteration of the College: Together with some other of uncommon Efficacy, taken from the most celebrated Authors. IV. Extemporaneous Prescriptions, distributed into Classes, suitable to their Intention in Cure. To which is added, An Account of the common Adulterations, both Simples and Compounds; with some Marks to detect them by. By JOHN QUINCY, M. D. The Ninth Edition, much enlarged and corrected. 8vo. 1733.

Statistical Essays: By STEPHEN HALES, B. D. F. R. S. Rector of Farringdon, Hampshire, and Minister of Teddington, Middlesex. In two Vols. 8vo. The first Vol. containing Vegetable Staticks; or, an Account of some Statical Experiments on the Sap in Vegetables. Being an Essay towards a Natural History of Vegetation: Of Use to those, who are curious in the Culture and Improvement of Gardening, &c. Also a Specimen of an Attempt to Analyse the Air, by a great Variety of Chymio-statical Experiments, which were read at several Meetings before the Royal Society. The second Vol. containing Hamastaticks; or an Account of some Hydraulic and Hydrostatical Experiments, made on the Blood and Blood-Vessels of Animals. Also an Account of some Experiments on Stones in the Kidneys and Bladder; with an Enquiry into the Nature of those anomalous Concretions. To which is added an Appendix, containing Observations and Experiments relating to several Subjects in the first Volume. The greatest Part of which were read at several Meetings before the Royal Society, with an Index to both Volumes.

Three Physico-Theological Discourses, concerning, I. The primitive Chaos and Creation of the World. The general Deluge, its Causes and Effects. III. The Dissolution of the World, and future Conflagration. Wherein are largely discuss'd the Production and Use of Mountains; the Original of Fountains, of formed Stones, and Sea-Fishes Bones and Shells found in the Earth; the Effects of particular Floods and Inundations of the Sea; the Eruptions of Volca-

nos; the Nature and Causes of Earthquakes. Also an Historical Account of those two late remarkable ones in Jamaica and England, with practical Inferences. By JOHN RAY, late Fellow of the Royal Society. The Fourth Edition. 8vo. 1732.

•*Cursus Medicinæ*; or, A compleat Theory of Physick: In Five Parts. I. The whole Doctrine of the Animal Oeconomy. II. The Nature, Difference, Causes and Symptoms of Diseases. III. The Diagnosticks, and Prognosticks of Distempers. IV. The Method of preventing Diseases. The Art of Healing; or the Manner of applying Remedies to particular Cases, with a preliminary Discourse of the Rise, Progress, Success, Nature and Principles of Medicine. Done principally from those admirable Institutions of the learned H. BOERHAAVE, Professor of Physick in the University of Leyden; and improv'd from others of the most eminent Authors. By JOHN CRAWFORD, M.D. 8vo. 1724.

Lectiones Pharmaceuticæ; or, A Course of Lectures in Pharmacy, Chymical and Galenical; explaining the whole Doctrine of that Art. By the late learned Dr. JOHN QUINCY. Published from his original Manuscript, with a Preface by P. SHAW, M.D. 4to. 1723.

The Gentleman's Steward and Tenants of Manors instructed. Containing rational, easy, and familiar Rules and Tables for finding the Value of Estates of Freehold, Copyhold, or Leasehold, as well on Lives as for Years absolute, &c. With an Enquiry into the Nature of the Annual Disbursements, Precariousness of the Tenure, and Casualties, that Estates in Fields or Houses, or both, are charged with, or liable to; and how they are to be accounted for in the Valuation. Wherein the Errors and Absurdities of all the common Methods of Valuation, and the Disadvantages thence accruing, sometimes to the Landlord and sometimes to the Tenant, are justly expos'd. The Tables for valuing Estates on Lives being founded on Dr. HALLEY's Hypothesis, and calculated by the Method laid down by Mr. ABR. DE MOIVRE, to 4, 5, 6, 7 and 8 per Cent. To which is added, An Appendix: Containing the Description and Use of an Instrument for discovering the Number of Feet contained in any Timber-Trees before they are cut down, by Inspection only. By JOHN RICHARDS of Exon. 8vo. 1730.

Philosophical Experiments and Observations of the late eminent Dr. ROBERT HOOKE, S. R. S. and Geom. Prof. Gresh. and eminent Virtuoso's in his Time. With Copper-Plates. Published by W. DERHAM, D.D. F. R. S. 8vo.

A Treatise of Algebra, in two Books: The first treating of the Arithmetical, and the second of the Geometrical Part. By PHILIP RONAYNE, Gent. The Second Edition. 8vo.

The Wisdom of God manifested in the Works of the Creation. In two Parts; viz. The Heavenly Bodies, Elements, Meteors, Fossils, Vegetables, Animals (Beasts, Birds, Fishes, and Insects) more particularly in the Body of the Earth; its Figure, Motion, and Consistency in the admirable Structure of the Bodies of Man, and other Animals; as also in their Generation, &c. With Answers to some Objections. By JOHN RAY, late Fellow of the Royal Society. The Ninth Edition, 8vo. 1727.

Philosophical Letters between the late learned Mr. Ray, and several of his ingenious Correspondents, Natives and Foreigners. To which are added, Those of Francis Willoughby, Esq; The whole consisting of many curious Discoveries and Improvements in the History of Quadrupeds, Birds, Fishes, Insects, Plants, Fossils, Fountains, &c. Published by the Rev. Mr. DERHAM. 8vo. 1718.

Synopsis Methodica Avium & Piscium, Opus posthumum, quod vivus recensuit & perfecit ipse insignissimus Author; in quo, multas species, in ipsius Ornithologia & Ichthyologia desideratas, adjecit, Methodumque suam Piscium naturæ magis convenientem reddit. Cum Appendice & Iconibus. Edente W. DERHAM. 8vo.

Dr. ROBERT HOOKE's Posthumous Works, in which the present Deficiency of Natural Philosophy is discours'd of; with the Methods of rendering it more certain and beneficial, &c. Published by RICHARD WALLER, Esq; Folio.

A Natural History of English Insects. Illustrated with a Hundred Copper-Plates, curiously engraven from the Life: And (for those who desire it) exactly coloured by the Author, ELIAZAR ALBIN, Painter. To which are added, large Notes, and many curious Observations, by W. DERHAM, F. R. S. 4to.